

Keep in Lake 6 This volume much not bave The herry, nor il ared in the building arbens the Suit's Carcent. fu. Jun 28. 73. B. \*\* + 275, 61a. Bequeathed to the Public Library of the City of Boston! Theodore Parker! Received 30th Colober, 1861. . So. 30,483.



Theodor ParMer.

This votreme centains corrections & additions in the hunderiting y her. Newton,





Parker 1 TRATIL

### ADVERTISEMENT.

PArt of the ensuing Discourse about Light was written at the desire of some Gentlemen of the Royal Society, in the Year 1675. and then Sent to their Secretary, and read at their Meetings, and the rest was added about Twelve Years after to complete the Theory; except the Third Book, and the last Proposition of the Second, which were since put together out of scattered Papers. To avoid being engaged in Disputes about these Matters, I have hitherto delayed the Printing, and Should still have delayed it, had not the importunity of Friends prevailed upon me. If any other Papers writ on this Subject are got out of my Hands they are imperfect, and were perhaps written before I had tried all the Experiments here set down, and fully satisfied my self about the Laws of Refractions and Composition of Colours. I have here Published what I think proper to come abroad, wishing that it may not be Translated into another Language without my Consent.

The Crowns of Colours, which fometimes appear about the Sun and Moon, I have endeavoured to give an Account of; but for want of sufficient Observations leave that Matter to be further examined. The Subject of the Third Book I have also left imperfect, not having tried all the ExpeExperiments which I intended when I was about these Matters, nor repeated some of those which I did try, until I had satisfied my self about all their Circumstances. To communicate what I have tried, and leave the rest to others for further Enquiry, is all my Design in publishing these Papers.

In a Letter written to Mr.Leibnitz in the Year 1676. and published by Dr. Wallis, I mentioned a Method by which I had found some general Theorems about squaring Curvilinear Figures, or comparing them with the Conic Sections, or other the simplest Figures with which they may be compared. And some Years ago I lent out a Manuscript containing such Theorems, and having since met with some Things copied out of it, I have on this Occasion made it publick, prefixing to it an Introduction and Subjoyning a Scholium concerning that Method. And I have joined with it another small Tract concerning the Curvilinear Figures of the Second Kind, which was also written many Tears ago, and made known to some Friends, who have solicited the making it publick.

I. N.

# [1]

# The FIRST BOOK

OF

# OPTICKS.

#### PART I.

Y Defign in this Book is not to explain the Properties of Light by Hypothefes, but to propole and prove them by Reafon and Experiments: In order to which, I shall premife the following Definitions and Axioms.

# DEFINITIONS.

#### DEFIN. I.

BY the Rays of Light I understand its least Parts, and those as well Successive in the fame Lines as Contemporary in feveral Lines. For it is manifest that Light confists of parts both Successive and Contemporary; because in the same place you may stop that which comes one moment, and let pass that which comes presently after; and in the same time you may stop it in any one place, and let it pass in any other. For that part of Light which is stopt cannot be the same with that which is let pass. The least Light or part of Light, which may be stopt alone without the rest of the Light, or propagated alone, or do or suffer any A thing thing alone, which the rest of the Light doth not or sufers not, I call a Ray of Light.

#### DEFIN. II.

Refrangibility of the Rays of Light, is their Disposition to be refracted or turned out of their Way in passing out of one transparent Body or Medium into another. And a greater or less Refrangibility of Rays, is their Disposition to be turned more or less out of their Way in like Incidences on the Same Medium. Mathematicians usually confider the Rays of Light to be Lines. reaching from the luminous Body to the body illuminated, and the refraction of those Rays to be the bending or breaking of those Lines in their passing out of one Medium into another. And thus may Rays and Refractions. be confidered, if Light be propagated in an instant. But by an Argument taken from the Æquations of the times of the Eclipses of Jupiter's Satellites it seems that Light is propagated in time, spending in its passage from the Sunto us about Seven Minutes of time : And therefore I have chosen to define Rays and Refractions in such general terms as may agree to Light in both cafes.

#### DEFIN. III.

Reflexibility of Rays, is their Disposition to be turned back into the fame Medium from any other Medium upon whose Surface they fall. And Rays are more or less reflexible, which are returned back more on less easily. As if Light pass out of Glass into Air, and by being inclined more and more to the common Surface of the Glass and Air, begins at length to be totally reflected by that Surface; those forts of Rays which, at like Incidences are reflected most copiously, or by inclining the Rays begin soonest to be totally reflected, are most reflexible. D E-

# [3]

#### DEFIN. IV.

The Angle of Incidence, is that Angle which the Line described by the incident Ray contains with the Perpendicular to the refle-Eting or refracting Surface at the Point of Incidence.

#### DEFIN. V.

The Angle of Reflexion or Refraction, is the Angle which the Line described by the reflected or refracted Ray containeth with the Perpendicular to the reflecting or refracting Surface at the Point of Incidence.

#### DEFIN. VI.

The Sines of Incidence, Reflexion, and Refraction, are the Sines of the Angles of Incidence, Reflexion, and Refraction.

#### DEFIN. VII.

The Light whose Rays are all alike Refrangible, I call Simple, Homogeneal and Similar; and that whose Rays are some more Refrangible than others, I call Compound, Heterogeneal and Dissimilar. The former Light I call Homogeneal, not because I would affirm it so in all respects; but because the Rays which agree in Refrangibility, agree at least in all those their other Properties. Which I confider in the following Discourse.

#### DEFIN. VIII.

The Colours of Homogeneal Lights, I call Primary, Homogeneal and Simple; and those of Heterogeneal Lights, Heterogeneal and Compound. For these are always compounded of the colours of Homogeneal Lights; as will appear in the following Discourse. A 2 AXI-

### [4]

# AXIOMS.

#### A X. I.

HE Angles of Incidence, Reflexion, and Refraction, lye in one and the fame Plane.

#### AX. II.

The Angle of Reflexion is equal to the Angle of Incidence.

#### A X. III.

If the refracted Ray be returned directly back to the Point of Incidence, it shall be refracted into the Line before described by the incident Ray.

#### A X. IV.

Refraction out of the rarer Medium into the denser, is made towards the Perpendicular; that is, so that the Angle of Refration be less than the Angle of Incidence.

#### A X. V.

The Sine of Incidence, is either accurately or very nearly in a given Ratio to the Sine of Refraction.

Whence if that Proportion be known in any one Inclination of the incident Ray, 'tis known in all the Inclinations, and thereby the Refraction in all cafes of Incidence on the fame refracting Body may be determined. Thus if the Refraction be made out of Air into Water, the Sine of Incidence of the red Light is to the Sine of its Refraction as 4 to 3. If out of Air into Glafs, the Sines are as as 17 to 11. In Light of other Colours the Sines have other Proportions : but the difference is so little that it need feldom be confidered.

Suppose therefore, that R S represents the Surface of Fig. 1. stagnating Water, and C is the point of Incidence in. which any Ray coming in the Air from A in the Line A C is reflected or refracted, and I would know whether this Ray shall go after Reflexion or Refraction : I erect upon the Surface of the Water from the point of Incidence the Perpendicular CP and produce it downwards to Q, and conclude by the first Axiom, that the Ray after Reflexion and Refraction, shall be found somewhere in the Plane of the Angle of Incidence A C P produced. I let fall therefore upon the Perpendicular C P the Sine of. Incidence A D, and if the reflected Ray be defired, I produce A D to B so that D B be equal to A D, and draw CB. For this Line CB shall be the reflected Ray; the Angle of Reflexion BCP and its Sine BD being equal to the Angle and Sine of Incidence, as they ought to be by the second Axiom. But if the refracted Ray be defired, I produce A D to H, so that D H may be to A D as the Sine of Refraction to the Sine of Incidence, that is as 3 to 4; and about the Center C and in the Plane A C P with the Radius C A describing a Circle A B E I draw Parallel to the Perpendicular CPQ, the Line HE cutting the circumference in E, and joyning CE, this Line CE shall be the Line of the refracted Ray. For if E F be let fall perpendicularly on the Line PQ, this Line EF shall be the Sine of Refraction of the Ray CE, the Angle of Refraction being ECQ; and this Sine EF is equal to DH, and confequently in Proportion to the Sine of Incidence A D as 3 to 4.

In

In like manner, if there be a Prism of Glass (that is a Glass bounded with two Equal and Parallel Triangular ends, and three plane and well polifhed Sides, which meet in three Parallel Lines running from the three Angles of one end to the three Angles of the other end) and if the Refraction of the Light in passing cross this Prism be desired : Let ACB represent a Plane cutting this Prism transversly to its three Parallel lines or edges there where the Light passeth through it, and let DE be the Ray incident upon the first side of the Prism A C where the Light goes into the Glass; And by putting the Proportion of the Sine of Incidence to the Sine of Refraction as 17 to 11 find E F the first refracted Ray. Then taking this Ray for the Incident Ray upon the second fide of the Glass B C where the Light goes out, find the next refracted Ray F G by putting the Proportion of the Sine of Incidence to the Sine of Refraction as 11 to 17. For if the Sine of Incidence out of Air into Glass be to the Sine of Refraction

as 17 to 11, the Sine of Incidence out of Glass into Air must on the contrary be to the Sine of Refraction as 11 to 17, by the third Axiom.

Fig. 3.

Much after the fame manner, if A C B D reprefent a Glafs fpherically Convex on both fides (ufually called a *Lens*, fuch as is a Burning-glafs, or Spectacle-glafs, or an Object-glafs of a Telefcope) and it be required to know how Light falling upon it from any lucid point Q fhall be refracted, let Q M reprefent a Ray falling upon any point M of its first spherical Surface A C B, and by erecting a Perpendicular to the Glafs at the point M, find the first refracted Ray M N by the Proportion of the Sines 17 to 11. Let that Ray in going out of the Glafs be incident upon N, and then find the fecond refracted Ray N q by the Proportion of the Sines 11 to 17. And after the fame

Fig. 2.

fame manner may the Refraction be found when the Lens is Convex on one fide and Plane or Concave on the other, or Concave on both Sides.

[7]

#### AX. VI.

Homogeneal Rays which flow from feveral Points of any Object, and fall almost Perpendicularly on any reflecting or refrating Plane or Spherical Surface, shall afterwards diverge from so many other Points, or be Parallel to so many other Lines, or converge to so many other Points, either accurately or without any sensible Error. And the same thing will happen, if the Rays be reflected or refracted successively by two or three or more Plane or spherical Surfaces.

The Point from which Rays diverge or to which they converge may be called their *Focus*. And the Focus of the incident Rays being given, that of the reflected or refracted ones may be found by finding the Refraction of any two Rays, as above; or more readily thus.

Caf. 1. Let ACB be a reflecting or refracting Plane, Fig. 43and Q the Focus of the incident Rays, and Q q C a perpendicular to that Plane. And if this perpendicular be produced to q, fo that q C be equal to QC, the point q fhall be the Focus of the reflected Rays. Or if q C be taken on the fame fide of the Plane with Q C and in Proportion to Q C as the Sine of Incidence to the Sine of Refraction, the point q fhall be the Focus of the refracted Rays.

Caf. 2. Let ACB be the reflecting Surface of any Fig. 5. Sphere whole Center is E. Bifect any Radius thereof (fuppofe E C) in T, and if in that Radius on the fame fide the point T you take the Points Q and q, fo that T Q, T E, and T q be continual Proportionals, and the point Q be the the Focus of the incident Rays, the point q shall be the Focus of the reflected ones.

Cal. 3. Let ACB be the refracting Surface of any Sphere whole Center is E. In any Radius thereof E C produced both ways take E T and Ct feverally in fuch Proportion to that Radius as the leffer of the Sines of Incidence and Refraction hath to the difference of those And then if in the fame Line you find any two Sines. Points Q and q, fo that TQ be to ET as Et to tq, taking t q the contrary way from t which T Q lieth from T, and if the Point Q be the Focus of any incident Rays, the Point q shall be the Focus of the refracted ones.

And by the fame means the Focus of the Rays after two or more Reflexions or Refractions may be found.

Cal. 4. Let ACBD be any refracting Lens, spheri-Wid. Whist and Mathcally Convex or Concave or Plane on either fide, and let CD be its Axis (that is the Line which cuts both its Surfaces perpendicularly, and passes through the Centers of the Spheres,) and in this Axis let F and f be the Foci of the refracted Rays found as above, when the incident Rays on both fides the Lens are Parallel to the fame Axis; and upon the Diameter F f bisected in E, describe a Circle. Suppose now that any Point Q be the Focus of any incident Rays. Draw QE cutting the faid Circle in T and t, and therein take t q in fuch Proportion to t E as t E or T E hath to T Q. Let t q lye the contrary way from t which T Q doth from T, and g shall be the Focus of the refracted Rays without any sensible Error, provided the Point Q be not so remote from the Axis, nor the Lens so broad as to make any of the Rays fall too obliquely on the refracting Surfaces.

> And by the like Operations may the reflecting or refracting Surfaces be found when the two Foci are given, and

Fig. 6.

Fig. 7.

p. 229.

God. Johist. Inal:

Mall. 9. 228.

and thereby a Lens be formed, which shall make the Rays flow towards or from what place you please.

So then the meaning of this Axiom is, that if Rays fall upon any Plane or Spherical Surface or Lens, and before their Incidence flow from or towards any Point Q, they shall after Reflexion or Refraction flow from or towards the Point q found by the foregoing Rules. And if the incident Rays flow from or towards feveral points Q, the reflected or refracted Rays shall flow from or towards fo many other Points q found by the fame Rules. Whether the reflected and refracted Rays flow from or towards the Point q is eafily known by the fituation of that Point. For if that Point be on the fame fide of the reflecting or refracting Surface or Lens with the Point Q, and the incident Rays flow from the Point Q, the reflected flow towards the Point q and the refracted from it; and if the incident Rays flow towards Q, the reflected flow from q, and the refracted towards it. And the contrary happens. when q is on the other fide of that Surface.

#### AX. VII.

Wherever the Rays which come from all the Points of any Object meet again in so many Points after they have been made to converge by Reflexion or Refraction, there they will make a Picture of the Object upon any white Body on which they fall.

So if PR reprefent any Object without Doors, and AB Fig. 3. be a Lens placed at a hole in the Window-flut of a dark Chamber, whereby the Rays that come from any Point Q of that Object are made to converge and meet again in the Point q; and if a Sheet of white Paper be held at qfor the Light there to fall upon it : the Picture of that Object PR will appear upon the Paper in its proper Shape B and and Colours. For as the Light which comes from the Point Q goes to the Point  $q_5$  fo the Light which comes from other Points P and R of the Object, will go to fo many other correspondent Points p and r (as is manifest by the fixth Axiom;) fo that every Point of the Object shall illuminate a correspondent Point of the Picture, and thereby make a Picture like the Object in Shape and Colour, this only excepted that the Picture shall be inverted. And this is the reason of that Vulgar Experiment of casting the Species of Objects from abroad upon a Wall or Sheet of white Paper in a dark Room.

Fig. 8.

In like manner when a Man views any Object PQR, the Light which comes from the feveral Points of the Object is so refracted by the transparent skins and humours of the Eye, (that is by the outward coat EFG called the Tunica Cornea, and by the crystalline humour AB which is beyond the Pupil mk) as to converge and meet again at fo many Points in the bottom of the Eye, and there to paint the Picture of the Object upon that skin (called the Tunica Retina) with which the bottom of the Eye is covered. For Anatomists when they have taken off from the bottom of the Eye that outward and most thick Coat called the Dura Mater, can then see through the thinner Coats the Pictures of Objects lively painted thereon. And these Pictures propagated by Motion along the Fibres of the Optick Nerves into the Brain, are the cause of Vision. For accordingly as these Pictures are perfect or imperfect, the Object is seen perfectly or imperfectly. If the Eye be tinged with any colour (as in the Disease of the Jaundise) so as to tinge the Pictures in the bottom of the Eye with that Colour, then all Objects appear tinged with the fame Colour. If the humours of the Eye by old Age decay, fo as by fhrinking to make the Cornea and Coat of the Crystalline

stalline humour grow flatter than before, the Light will not be refracted enough, and for want of a sufficient Refraction will not converge to the bottom of the Eye but to some place beyond it, and by consequence paint in the bottom of the Eye a confused Picture, and according to the indistinctness of this Picture the Object will appear confused. This is the reason of the decay of Sight in old Men, and shews why their Sight is mended by Spectacles. For those Convex-glasses supply the defect of plumpnels in the Eye, and by encreasing the Refraction make the Rays converge sooner to as to convene diffinctly at the bottom of the Eye if the Glass have a due degree of convexity. And the contrary happens in short-sighted Men whose Eyes are too plump. For the Refraction being now too great, the Rays converge and convene in the Eyes before they come at the bottom; and therefore the Picture made in the bottom and the Vision caused thereby will not be distinct, unless the Object be brought fo near the Eye as that the place where the converging Rays convene may be removed to the bottom, or that the plumpnels of the Eye be taken off and the Refractions diminished by a Concave-glass of a due degree of Concavity, or lastly that by Age the Eye grow flatter till it come to a due Figure : For short-sighted Men see remote Objects best in Old Age, and therefore they are accounted to have the most lasting Eyes.

#### A X. VIII.

An Object Seen by Reflexion or Refraction, appears in that place from whence the Rays after their last Reflexion or Refraction diverge in falling on the Spectator's Eye.

If the Object A be seen by Reflexion of a Looking-Fig. 9. glass m n, it shall appear, not in it's proper place A, but B 2 behind behind the Glass at *a*, from whence any Rays AB, AC, AD, which flow from one and the same Point of the Object, do after their Reflexion made in the Points B, C, D, diverge in going from the Glass to E, F, G, where they are incident on the Spectator's Eyes. For these Rays do make the same Picture in the bottom of the Eyes as if they had come from the Object really placed at *a* without the interposition of the Looking-glass; and all Vision is made according to the place and shape of that Picture. In like manner the Object D feen through a Prism ap-

Fig. 2. In like manner the Object D feen through a Prifm appears not in its proper place D, but is thence translated to fome other place d fituated in the last refracted Ray F G drawn backward from F to d.

Fig. 10,

And fo the Object Q feen through the Lens A B, appears at the place q from whence the Rays diverge in paffing from the Lens to the Eye. Now it is to be noted, that the Image of the Object at q is fo much bigger or leffer than the Object it felf at Q, as the diffance of the Image at q from the Lens AB is bigger or lefs than the diffance of the Object at Q from the fame Lens. And if the Object be feen through two or more fuch Convex or Concaveglaffes, every Glafs fhall make a new Image, and the Object fhall appear in the place and of the bignefs of the laft Image. Which confideration unfolds the Theory of Microfcopes and Telefcopes. For that Theory confifts in almoft nothing elfe than the defcribing fuch Glaffes as fhall make the laft Image of any Object as diffinct and large: and luminous as it can conveniently be made.

I have now given in Axioms and their Explications the fumm of what hath hitherto been treated of in Opticks. For what hath been generally agreed on I content my felf to affume under the notion of Principles, in order to what I have further to write. And this may fuffice for an IntroIntroduction to Readers of quick Wit and good Underftanding not yet versed in Opticks: Although those who are already acquainted with this Science, and have handled Glasses, will more readily apprehend what followeth.

## PROPOSITIONS.

#### PROP. I. Theor. I.

IGHTS which differ in Colour, differ also in Degrees of Refrangibility.

### The Proof by Experiments.

Exper. 1. I took a black oblong ftiff Paper terminated by Parallel Sides, and with a Perpendicular right Line drawn cross from one Side to the other, distinguished it into two equal Parts. One of these Parts I painted with a red Colour and the other with a blew. The Paper wasvery black, and the Colours intense and thickly laid on, that the Phænomenon might be more conspicuous. This Paper I viewed through a Prism of solid Glass, whose two Sides through which the Light passed to the Eye were plane and well polished, and contained an Angle of about Sixty Degrees : which Angle I call the refracting Angle of the Prism. And whilst I viewed it, I held it before a cig Prism. Window in fuch manner that the Sides of the Paper were parallel to the Prism, and both those Sides and the Prism parallel to the Horizon, and the crofs Line perpendicular to it; and that the Light which fell from the Window HPON

### upon the Paper made an Angle with the Baper, segual to that Angle which was made with the fame Paper by the Light reflected from it to the Eye. Beyond the Prism was the Wall of the Chamber under the Window covered over with black Cloth, and the Cloth was involved in Darkness that no Light might be reflected from thence, which in passing by the edges of the Paper to the Eye, might mingle it self with the Light of the Paper and obscure the Phænomenon thereof. These things being thus ordered, I found that if the refracting Angle of the Prism be turned upwards, so that the Paper may seem to be lifted upwards by the Refraction, its blew half will be lifted higher by the Refraction than its red half. But if the refracting Angle of the Prism be turned downward, so that the Paper may seem to be carried lower by the Refraction, its blew half will be carried something lower thereby than its red half. Wherefore in both cases the Light which comes from the blew half of the Paper through the Prism to the Eye, does in like Circumstances suffer a greater Refraction than the Light which comes from the red half, and by confequence is more refrangible.

Fig. 11. Illustration. In the Eleventh Figure, M N represents the Window, and DE the Paper terminated with parallel Sides DJ and HE, and by the transverse Line FG distinguished into two halfs, the one D G of an intenfely blew Colour, the other FE of an intensely red. And BACcab reprefents the Prism whose refracting Planes ABba and ACca *C* & transverse inter de la contraction de la c wards in such manner that the blew half DG is carried higher to dg than the red half F E is to fe, and therefore fuffers

fuffers a greater Refraction. If the edge of the refracting Angle be turned downward, the Image of the Paper will be refracted downward suppose to  $\delta \varepsilon$ , and the blew half will be refracted lower to  $\delta \gamma$  than the red half is to  $\varphi \varepsilon$ .

Exper. 2. About the aforefaid Paper, whose two halfs were painted over with red and blew, and which was stiff like thin Pastboard, I lapped several times a slender thred of very black Silk, in fuch manner that the feveral parts of the thred might appear upon the Colours like fo many black Lines drawn over them, or like long and flender dark Shadows caft upon them. I might have drawn black Lines with a Pen, but the threds were fmaller and better defined. This Paper thus coloured and lined I fet against a Wall perpendicularly to the Horizon, so that one of the Colours might stand to the right hand and the other to the left. Close before the Paper at the confine of the Colours below I placed a Candle to illuminate the Paper strongly : For the Experiment was tried in the Night. The flame of the Candle reached up to the lower edge of the Paper, or a very little higher. Then at the distance of Six Feet and one or two Inches from the Paper upon the Floor I crected a glass Lens four Inches and a quarter broad, which might collect the Rays coming from the several Points of the Paper, and make them converge towards so many other Points at the same distance of fix Feet and one or two Inches on the other fide of the Lens, and so form the Image of the coloured Paper upon a white Paper placed there; after the fame manner that a Lens at a hole in a Window casts the Images of Objects abroad upon a Sheet of white Paper in a dark Room. The aforefaid white Paper, erected perpendicular to the Horizon and to the Rays which fell upon it from the Lens, I moved sometimes towards the Lens, sometimes from it, to find the

the places where the Images of the blew and red parts of the coloured Paper appeared most distinct. Those places I easily knew by the Images of the black Lines which I had made by winding the Silk about the Paper. For the Images of those fine and slender Lines (which by reason of their blackness were like Shadows on the Colours) were confused and scarce visible, unless when the Colours on either side of each Line were terminated most distinctly. Noting therefore, as diligently as I could, the places where the Images of the red and blew halfs of the coloured Paper appeared most distinct, I found that where the red half of the Paper appeared distinct, the blew half appeared confused, so that the black Lines drawn upon it could scarce be seen; and on the contrary where the blew half appeared most distinct the red half appeared confused, so that the black Lines upon it were scarce visible. And between the two places where these Images appeared distinct there was the distance of an Inch and a half : the distance of the white Paper from the Lens, when the Image of the red half of the coloured Paper appeared most distinct, being greater by an Inch and an half than the distance of the same white Paper from the Lens when the Image of the blew half appeared most distinct. In like Incidences therefore of the blew and red upon the Lens, the blew was refracted more by the Lens than the red, so as to converge fooner by an Inch and an half, and therefore is more refrangible.

Fig. 12.

Illustration. In the Twelfth Figure, DE fignifies the coloured Paper, DG the blew half, FE the red half, MN the Lens, HJ the white Paper in that place where the red half with its black Lines appeared diftinct, and *bi* the fame Paper in that place where the blew half appeared diftinct. The place *bi* was nearer to the Lens MN than the place HJ by an Inch and an half. Scholium.

1 -

Scholium. The fame things fucceed notwithstanding that some of the Circumstances be varied : as in the first Experiment when the Prism and Paper are any ways inclined to the Horizon, and in both when coloured Lines are drawn upon very black Paper. But in the Description of these Experiments, I have set down such Circumstances by which either the Phænomenon might be rendred more conspicuous, or a Novice might more easily try them, or by which I did try them only. The fame thing I have often done in the following Experiments : Concerning all which this one Admonition may fuffice. Now from these Experiments it follows not that all the Light of the blew is more Refrangible than all the Light of the red ; For both Lights are mixed of Rays differently Refrangible, So that in the red there are some Rays not less Refrangible than those of the blew, and in the blew there are some Rays not more Refrangible than those of the red; But these Rays in Proportion to the whole Light are but few, and ferve to diminish the Event of the Experiment, but are not able to destroy it. For if the red and blew Colours were more dilute and weak, the diftance of the Images would be lefs than an Inch and an half; and if they were more intense and full, that distance would be greater, as will appear hereafter. These Experiments may suffice for the Colours of Natural Bodies. For in the Colours made by the Refraction of Prisms this Proposition will appear by the Experiments which are now to follow in the next Proposition.

-X-

### [ 81 ]

### PROP. II. Theor. II.

The Light of the Sun confifts of Rays differently Refrangible.

### The Proof by Experiments.

Exper. 3. N a very dark Chamber at a round hole about one third part of an Inch broad made in the Shut of a Window I placed a Glass Prism, whereby the beam of the Sun's Light which came in at that hole might be refracted upwards toward the opposite Wall of the Chamber, and there form a coloured Image of the Sun. The Axis of the Prifm (that is the Line passing through the middle of the Prism from one end of it to the other end Parallel to the edge of the Refracting Angle) was in this and the following Experiments perpendicular to the incident Rays. About this Axis I turned the Prism flowly, and faw the refracted Light on the Wall or coloured Image of the Sun first to descend and then to ascend. Between the Descent and Ascent when the Image feemed Stationary, I stopt the Prism, and fixt it in that Posture, that it should be moved no more. For in that posture the Refractions of the Light at the two sides of the Refracting Angle, that is at the entrance of the Rays into the Prism and at their going out of it, were equal to one another. So also in other Experiments as often as I would have the Refractions on both sides the Prism to be equal to one another, I noted the place where the Image of the Sun formed by the refracted Light ftood still between its two contrary Motions, in the common Period of its progrefs and egrefs; and when the Image fell upon that place, I made fast the Prism. And in this posture, as the

the most convenient, it is to be understood that all the Prisms are placed in the following Experiments, unless where some other posture is described. The Prism therefore being pla-ced in this posture, I let the refracted Light fall perpendicularly upon a Sheet of white Paper at the opposite Wall of the Chamber, and observed the Figure and Dimensions of the Solar Image formed on the Paper by that Light. This Image was Oblong and not Oval, but terminated with two Rectilinear and Parallel Sides, and two Semicircular Ends. On its Sides it was bounded pretty distinctly, but on its Ends very confusedly and indistinctly, the Light there decaying and vanishing by degrees. The breadth of this Image answered to the Sun's Diameter, and was about two Inches and the eighth part of an Inch, including the Penumbra. For the Image was eighteen Feet and an half distant from the Prism, and at this distance that breadth if diminished by the Diameter of the hole in the Window-shut, that is by a quarter of an Inch, subtended an Angle at the Prism of about half a Degree, which is the Sun's apparent Diameter. But the length of the Image was about ten Inches and a quarter, and the length of the Rectilinear Sides about eight Inches; And the refracting Angle of the Prifm whereby so great a length was made, was 64 degr. With a less Angle the length of the Image was less, the breadth remaining the fame. If the Prism was turned about its Axis that way which made the Rays emerge more obliquely out of the second refracting Surface of the Prism, the Image soon became an Inch or two longer, or more; and if the Prifm was turned about the contrary way, fo as to make the Rays tall more obliquely on the first refracting Surface, the Image foon became an Inch or two shorter. And therefore in trying this Experiment, I was as curious as I could be in placing the Prism by the above-mentioned Rule exactly in fuch

J.

fuch a polture that the Refractions of the Rays at their emergence out of the Prism might be equal to that at their incidence on it. This Prism had some Veins running along within the Glass from one end to the other, which scattered some of the Sun's Light irregularly, but had no sensible effect in encreasing the length of the coloured Spec-For I tried the same Experiment with other Prisms trum. with the fame Success. And particularly with a Prism which feemed free from fuch Veins, and whofe refracting Angle was 62<sup>1</sup> Degrees, I found the length of the Image 9<sup>3</sup> or 10 Inches at the diftance of  $18\frac{1}{2}$  Feet from the Prifm, the breadth of the hole in the Window-shut being  $\frac{1}{4}$  of an Inch as before. And because it is easie to commit a mistake in placing the Prism in its due posture, I repeated the Experiment four or five times, and always found the length of the Image that which is fet down above. With another Prism of clearer Glass and better Pollish, which feemed free from Veins and whofe refracting Angle was 63 Degrees, the length of this Image at the same distance of 18<sup>1</sup>/<sub>2</sub> Feet was also about 10 Inches, or 10<sup>1</sup>/<sub>8</sub>. Beyond these Measures for about  $\frac{1}{4}$  or  $\frac{1}{3}$  of an Inch at either end of the Spectrum the Light of the Clouds seemed to be a little tinged with red and violet, but fo very faintly that I sufpected that tincture might either wholly or in great measure arise from some Rays of the Spectrum scattered irregularly by fome inequalities in the Substance and Polish of the Glass, and therefore I did not include it in these Meafures. Now the different Magnitude of the hole in the Window-shut, and different thickness of the Prism where the Rays passed through it, and different inclinations of the Prism to the Horizon, made no sensible changes in the length of the Image. Neither did the different matter of the

the Prisms make any : for in a Vessel made of polished Plates of Glass cemented together in the shape of a Prism and filled with Water, there is the like Success of the Experiment according to the quantity of the Refraction. It is further to be observed, that the Rays went on in right Lines from the Prism to the Image, and therefore at their very going out of the Prism had all that Inclination to one another from which the length of the Image pro-ceeded, that is the Inclination of more than two Degrees and an half. And yet according to the Laws of Opticks vulgarly received, they could not possibly be so much in-clined to one another. For let EG represent the Window-Fig. 13. fhut, F the hole made therein through which a beam of the Sun's Light was transmitted into the darkned Chamber, and ABC a Triangular Imaginary Plane whereby the Prism is feigned to be cut transversly through the middle of the Light. Or if you please, let ABC represent the Prism it felf, looking directly towards the Spectator's Eye with its nearer end : And let XY be the Sun, MN the Paper upon which the Solar Image or Spectrum is caft, and P T the Image it felf whofe fides tovvards V and W are Rectilinear and Parallel, and ends tovvards P and T Semicircular. YKHP and XLJT are the two Rays, the first of which comes from the lower part of the Sun to the higher part of the Image, and is refracted in the Prism at K and H, and the latter comes from the higher part of the Sun to the lower part of the Image, and is refracted at L and J. Since the Refractions on both fides the Prism are equal to one another, that is the Refraction at K equal. to the Refraction at J, and the Refraction at L equal to the Refraction at H, so that the Refractions of the incident Rays at K and L taken together are equal to the Refractions of the emergent Rays at H and J taken together :

ther : it follows by adding equal things to equal things, that the Refractions at K and H taken together, are equal to the Refractions at J and L taken together, and therefore the two Rays being equally refracted have the fame Inclination to one another after Refraction which they had before, that is the Inclination of half a Degree answering to the Sun's Diameter. For so great was the Inclination of the Rays to one another before Refraction. So then, the length of the Image P T would by the Rules of Vulgar Opticks subtend an Angle of half a Degree at the Prism, and by consequence be equal to the breadth  $\nu w$ ; and therefore the Image would be round. Thus it would be were the two Rays XLJT and YKHP and all the rest which form the Image P w T v, alike Refrangible. And therefore seeing by Experience it is found that the Image is not round but about five times longer than broad, the Rays which going to the upper end P of the Image suffer the greatest Refraction, must be more Refrangible than those which go to the lower end T, unless the inequality of Refraction be casual.

This Image or Spectrum P T was coloured, being red at its leaft refracted end T, and violet at its most refracted end P, and yellow green and blew in the intermediate spaces. Which agrees with the first Proposition, that Lights which differ in Colour do also differ in Refrangibility. The length of the Image in the foregoing Experiments I measured from the faintest and outmost red at one end, to the faintest and outmost blew at the other end. I kein away w *Exper.* 4. In the Sun's beam which was propagated into the Room through the hole in the Window-flut, at the distance of some Feet from the hole, I held the Prism in such a posture that its Axis might be perpendicular to that beam. Then I looked through the Prism upon the hole,

with .

hole, and turning the Prifin to and fro about its Axis to make the Image of the hole afcend and defcend, when between its two contrary Motions it feemed flationary, I ftopt the Prifin that the Refractions on both fides of the refracting Angle might be equal to each other as in the former Experiment. In this Situation of the Prifin viewing through it the faid hole, I obferved the length of its refracted Image to be many times greater than its breadth, and that the most refracted part thereof appeared violet, \*the least refracted red, the middle parts blew green and yellow in order. The fame thing happened when I removed the Prifin out of the Sun's Light, and looked through it upon the hole finning by the Light of the Clouds beyond it. And yet if the Refraction were done regularly according to one certain Proportion of the Sines of Incidence and Refraction as is vulgarly supposed, the refracted Image ought to have appeared round.

So then, by these two Experiments it appears that in equal Incidences there is a confiderable inequality of Refractions : But whence this inequality arises, whether it be that some of the incident Rays are refracted more and others less, constantly or by chance, or that one and the fame Ray is by Refraction disturbed, shattered, dilated, and as it were split and spread into many diverging Rays, as Grimaldo supposes, does not yet appear by these Experiments, but will appear by those that follow.

Exper. 5. Confidering therefore, that if in the third Experiment the Image of the Sun should be drawn out into an oblong form, either by a Dilatation of every Ray, or by any other casual inequality of the Refractions, the same oblong Image would by a second Refraction made Sideways be drawn out as much in breadth by the like Dilatation of the Rays or other casual inequality of the Refractions. fractions Sideways, I tried what would be the Effects of fuch a fecond Refraction. For this end I ordered all things as in the third Experiment, and then placed a fecond Prism immediately after the first in a cross Position to it, that it might again refract the beam of the Sun's Light which came to it through the first Prism. In the first Prism this beam was refracted upwards, and in the fecond Sideways. And I found that by the Refraction of the fecond Prifm the breadth of the İmage was not increased, but its superior part which in the first Prism suffered the greater Refraction and appeared violet and blew, did again in the second Prism suffer a greater Refraction than its inferior part, which appeared red and yellow, and this without any Dilation of the Image in breadth.

Fig. 14. Illustration. Let S represent the Sun, F the hole in the Window, ABC the first Prism, DH the second Prism, Y the round Image of the Sun made by a direct beam of Light when the Prisms are taken away, PT the oblong Image of the Sun made by that beam passing through the first Prism alone when the second Prism is taken away, and pt the Image made by the cross Refractions of both Prisms together. Now if the Rays which tend towards the several Points of the round Image Y were dilated and spread by the Refraction of the first Prism, so that they should not any longer go in single Lines to single Points, but that every Ray being split, shattered, and changed from a Linear Ray to a Superficies of Rays diverging from the Point of Refraction, and lying in the Plane of the Angles of Incidence and Refraction, they should go in those Planes to so many Lines reaching almost from one end of the Image PT to the other, and if that Image should thence become oblong : those Rays and their feveral parts tending towards the feveral Points of the
the Image P T ought to be again dilated and spread Sideways by the transverse Refraction of the second Prism, so as to compose a foursquare Image, such as is represented at #7. For the better understanding of which, let the Image PT be diftinguished into five equal Parts PQK, KQRL, LRSM, MSVN, NVT. And by the same irregularity that the Orbicular Light Y is by the Refraction of the first Prism dilated and drawn out into a long Image P T, the the Light PQK which takes up a space of the same length and breadth with the Light Y ought to be by the Refra-Ation of the fecond Prism dilated and drawn out into the long Image  $\pi q kp$ , and the Light KQRL into the long Image k q r l, and the Lights LRSM, MSVN, NVT into fo many other long Images l r s m, m s v n, n v t1; and all these long Images would compose the foursquare Image  $\pi 1$ . Thus it ought to be were every Ray dilated by Re-fraction, and spread into a triangular Superficies of Rays diverging from the Point of Refraction. For the second Refraction would spread the Rays one way as much as the first doth another, and so dilate the Image in breadth as much as the first doth in length. And the fame thing ought to happen, were some Rays casually refracted more than others. But the Event is otherwise. The Image P T was not made broader by the Refraction of the second Prism, but only became oblique, as 'tis represented at p t, its upper end P being by the Refraction translated to a greater distance than its lower end T. So then the Light which went towards the upper end P of the Image, was, (at equal Incidences) more refracted in the fecond Prism than the Light which tended towards the lower end T, that is the blew and violet, than the red and yellow; and therefore was more Refrangible. The same Light was by the Refraction of the first Prism translated further from the

place

place Y to which it tended before Refraction; and therefore suffered as well in the first Prism as in the second a greater Refraction than the rest of the Light, and by consequence was more Refrangible than the rest, even before its incidence on the first Prism.

+

Sometimes I placed a third Prism after the second, and fometimes also a fourth after the third, by all which the Image might be often refracted fideways : but the Rays which were more refracted than the rest in the first Prism were also more refracted in all the rest, and that without any Dilatation of the Image fideways : and therefore those Rays for their constancy of a greater Refraction are defervedly reputed more Refrangible.

But that the meaning of this Experiment may more Fig. 15. clearly appear, it is to be confidered that the Rays which are equally Refrangible do fall upon a circle answering to the Sun's Disque. For this was proved in the third Experiment. By a circle I understand not here a perfect Geometrical Circle, but any Orbicular Figure whose length is. equal to its breadth, and which, as to fense, may seem circular. Let therefore A G represent the circle which all the most Refrangible Rays propagated from the whole Difque of the Sun, would illuminate and paint upon the opposite Wall if they were alone; E L the circle which all the least Refrangible Rays would in like manner illuminate and paint if they were alone; BH, CJ, DK, the circles which fo many intermediate forts of Rays would fuccefsively paint upon the Wall, if they were singly propagated from the Sun in successive Order, the rest being always intercepted; And conceive that there are other intermediate Circles without number which innumerable other intermediate forts of Rays would fucceffively paint upon the Wall if the Sun should successively emit every fort apart. And

[26]

And feeing the Sun emits all these forts at once, they must all together illuminate and paint innumerable equal circles, of all which, being according to their degrees of Refrangibility placed in order in a continual feries, that oblong Spectrum P T is composed which I described in the third Experiment. Now if the Sun's circular Image Y Fig: 14 & 15. which is made by an unrefracted beam of Light was by any dilatation of the fingle Rays, or by any other irregularity in the Refraction of the first Prism, converted into the Oblong Spectrum, PT: then ought every circle AG, BH, C J, Gc. in that Spectrum, by the cross Refraction of the fecond Prism again dilating or otherwife scattering the Rays as before, to be in like manner drawn out and transformed into an Oblong Figure, and thereby the breadth of the Image P T would be now as much augmented as the length of the Image Y was before by the Refraction of the first Prism; and thus by the Refractions of both Prisms together would be formed a foursquare Figure  $p \neq t$ ? as I described above. Wherefore fince the breadth of the Spectrum P T is not increased by the Refraction fideways, it is certain that the Rays are not split or dilated, or otherways irregularly scattered by that Refraction, but that every circle is by a regular and uniform Refraction translated entire into another place, as the circle AG by the greatest Refraction into the place ag, the circle BH by a less Refraction into the place bh, the circle CJ by a Refraction still less into the place ci, and so of the rest; by which means a new Spectrum pt inclined to the former PT is in like manner composed of circles lying in a right Line; and these circles must be of the same bigness with the former, because the breadths of all the Spectrums Y, PT and pt at equal distances from the Prisms are equal.

D 2

I con-

I confidered further that by the breadth of the hole F through which the Light enters into the Dark Chamber, there is a Penumbra made in the circuit of the Spectrum Y, and that Penumbra remains in the rectilinear Sides of the Spectrums P T and pt. I placed therefore at that hole a Lens or Object-glass of a Telescope which might cast the Image of the Sun distinctly on Y without any Penumbra at all, and found that the Penumbra of the Rectilinear Sides of the oblong Spectrums P T and pt was also thereby taken away, so that those Sides appeared as distinctly defined as did the Circumference of the first Image Y. Thus it happens if the Glass of the Prisms be free from veins, atd their Sides be accurately plane and well polished without those numberless waves or curles which ulually arife from Sand-holes a little smoothed in polishing with Putty. If the Glass be only well polished and free from veins and the Sides not accurately plane but a little Convex or Concave, as it frequently happens; yet may the three Spectrums Y, P T and pt want Penumbras, but not in equal distances from the Prisms. Now from this want of Penumbras, I knew more certainly that every one of the circles was refracted according to some most regular, uniform, and constant law. For if there were any irregularity in the Refraction, the right Lines A E and G L which all the circles in the Spectrum P T do touch, could not by that Refraction be translated into the Lines. a e and g l as diffinct and straight as they were before, but there would arife in those translated Lines some Penumbra or crookedness or undulation, or other sensible Perturbation contrary to what is found by Experience. Whatfoever Penumbra or Perturbation should be made in the circles by the cross Refraction of the second Prism, all that Penumbra or Perturbation would be conspicuous in the

the right Lines a e and g l which touch those circles. And therefore fince there is no such Penumbra or Perturbation in those right Lines there must be none in the circles. Since the distance between those Tangents or breadth of the Spectrum is not increased by the Refractions, the Diameters of the circles are not increased thereby. Since those Tangents continue to be right Lines, every circle which in the first Prism is more or less refracted, is exactly in the same Proportion more or less refracted in the second. And seeing all these things continue to succeed after the fame manner when the Rays are again in a third Prism, and again in a fourth refracted Sideways, it is evident that the Rays of one and the fame circle as to their degree of Refrangibility continue always Uniform and Homogeneal to one another, and that those of several circles do differ in degree of Refrangibility, and that in fome certain and constant Proportion. Which is the thing I was to prove.

There is yet another Circumstance or two of this Ex-Fig. 16. periment by which it becomes still more plain and convincing. Let the second Prism DH be placed not immeately after after the first, but at some distance from it; Suppose in the mid-way between it and the Wall on which the oblong Spectrum P T is cast, so that the Light from the first Prism may fall upon it in the form of an oblong Spectrum, #1 Parallel to this second Prism, and be refracted Sideways to form the oblong Spectrum pt upon the Wall. And you will find as before, that this Spectrum pt is inclined to that Spectrum P T, which the first Prism forms alone without the fecond; the blew ends P and p being further diftant from one another than the red ones T and  $t_1$ , and by confequence that the Rays which go to the blew end # of the Image #1 and which therefore suffer the greatest Refraction in the first Prism, are again in the second Prism more refracted than the reft. The

Fig. 17. The fame thing I try'd also by letting the Sun's Light into a dark Room through two little round holes F and o made in the Window, and with two Parallel Prisms ABC and «By placed at those holes (one at each) refracting those two beams of Light to the opposite Wall of the Chamber, in such manner that the two colour'd Images P T and MR which they there painted were joyned end to end and lay in one straight Line, the red end T of the one touching the blew end M of the other. For if these two refracted beams were again by a third Prism D H placed crost to the two first, refracted Sideways, and the Spectrums thereby translated to some other part of the Wall of the Chamber, suppose the Spectrum PT to pt and the Spectrum M N to mn, these translated Spectrums pt Grid. 9 33. and m n would not lie in one ftraight Line with their ends contiguous as before, but be broken off from one another and become Parallel, the blew end of the Image m n being by a greater Refraction translated farther from its former place M T, than the red end t of the other Image p t from the fame place MT which puts the Proposition past dispute. And this happens whether the third Prism DH be placed immediately after the two first or at a great distance from them, so that the Light refracted in the two first Prisms be either white and circular, or coloured and oblong when it falls on the third.

> *Exper.* 6. In the middle of two thin Boards I made round holes a third part of an Inch in Diameter, and in the Window-fhut a much broader hole, being made to let into my darkned Chamber a large beam of the Sun's Light; I placed a Prifm behind the Shut in that beam to refract it towards the oppofite Wall, and clofe behind the Prifm I fixed one of the Boards, in fuch manner that the middle of the refracted Light might pafs through the hole made

made in it, and the reft be intercepted by the Board. Then at the distance of about twelve Feet from the first Board I fixed the other Board, in such manner that the middle of the refracted Light which came through the hole in the first Board and fell upon the opposite Wall might pass through the hole in this other Board, and the rest being intercepted by the Board might paint upon it the coloured Spectrum of the Sun. And close behind this Board I fixed another Prism to refract the Light which came through the hole. Then I returned speedily to the first Prism, and by turning it slowly to and fro about its Axis, I cauled the Image which fell upon the fecond Board to move up and down upon that Board, that all its parts. might successively pass through the hole in that Board and fall upon the Prism behind it. And in the mean time, I noted the places on the opposite Wall to which that Light after its Refraction in the second Prism did pass; and by the difference of the places I found that the Light which being most refracted in the first Prism did go to the blew end of the Image, was again more refracted in the second Prism than the Light which went to the red end of that Image, which proves as well the first Proposition as the fecond. And this happened whether the Axis of the two Prisms were parallel, or inclined to one another and to the Horizon in any given Angles.

Illustration. Let F be the wide hole in the Window-fhut, Fig. 18. through which the Sun fhines upon the first Prism A B C, and let the refracted Light fall upon the middle of the Board D E, and the middle part of that Light upon the hole G made in the middle of that Board. Let this trajected part of the Light fall again upon the middle of the fecond Board de and there paint such an oblong coloured Image of the Sun as was described in the third Experiment. By [32]

By turning the Prism A B C flowly to and fro about its Axis this Image will be made to move up and down the Board de, and by this means all its parts from one end to the other may be made to pass successively through the hole g which is made in the middle of that Board. In the mean while another Prism abc is to be fixed next after that hole g to refract the trajected Light a second time. And these things being thus ordered, I marked the places M and N of the opposite Wall upon which the refracted Light fell, and found that whilft the two Boards and fecond Prism remained unmoved, those places by turning the first Prism about its Axis were changed perpetually. For when the lower part of the Light which fell upon the second Board de was cast through the hole g it went to a lower place M on the Wall, and when the higher part of that Light was cast through the same hole g, it went to a higher place N on the Wall, and when any intermediate part of the Light was cast through that hole it went to some place on the Wall between M and N. The unchanged Polition of the holes in the Boards, made the Incidence of the Rays upon the second Prism to be the same in all cases. And yet in that common Incidence some of the Rays were more refracted and others less. And those were more refracted in this Prism which by a greater Refraction in the first Prism were more turned out of the way, and therefore for their conftancy of being more refracted are defervedly called more Refrangible.

*Exper.* 7. At two holes made near one another in my Window-fhut I placed two Prifms, one at each, which might caft upon the oppofite Wall (after the manner of the third Experiment) two oblong coloured Images of the Sun. And at a little diftance from the Wall I placed a long flender Paper with ftraight and parallel edges, and ordered

Pr 10

ordered the Prisms and Paper so, that the red Colour of one Image might fall directly upon one half of the Paper, and the violet colour of the other Image upon the other half of the same Paper; so that the Paper appeared of two Colours, red and violet, much after the manner of the painted Paper in the first and second Experiments. Then with a black Cloth I covered the Wall behind the Paper, that no Light might be reflected from it to disturb the Experiment, and viewing the Paper through a third Prism . held parallel to it, I faw that half of it which was illuminated by the Violet-light to be divided from the other half by a greater Refraction, especially when I went a good way off from the Paper. For when I viewed it too near at hand, the two halfs of the Paper did not appear fully divided from one another, but seemed contiguous at one of their Angles like the painted Paper in the first Experiment. Which also happened when the Paper was too broad.

Sometimes instead of the Paper I used a white Thred, and this appeared through the Prism divided into two Parallel Threds as is represented in the 19th Figure, where Fig. 19. DG denotes the Thred illuminated with violet Light from D to E and with red Light from F to G, and de fgare the parts of the Thred seen by Refraction. If one half of the Thred be constantly illuminated with red, and the other half be illuminated with all the Colours successively, (which may be done by causing one of the Prisms to be turned about its Axis whilft the other remains unmoved) this other half in viewing the Thred through the Prism, will appear in a continued right Line with the first half when illuminated with red, and begin to be a little divided from it when illuminated with Orange, and remove turther from it when illuminated with Yellow, and still further E

further when with Green, and further when with Blew, and go yet further off when illuminated with Indigo, and furtheft when with deep Violet. Which plainly fhews, that the Lights of feveral Colours are more and more Refrangible one than another, in this order of their Colours, Red, Orange, Yellow, Green, Blew, Indigo, deep Violet; and fo proves as well the first Proposition as the fecond.

Fig. 17.

an free

I caufed alfo the coloured Spectrums P T and M N made in a dark Chamber by the Refractions of two Prifms to lye in a right Line end to end, as was defcribed above in the fifth Experiment, and viewing them through a third Prifm held Parallel to their length, they appeared no longer in a right Line, but became broken from one another, as they are reprefented at pt and mn, the violet end m of the Spectrum mn being by a greater Refraction tranflated further from its former place M T than the red end t of the other Spectrum pt.

Fig. 20. I further caufed those two Spectrums P T and MN to become co-incident in an inverted order of their Colours, the red end of each falling on the violet end of the other, as they are represented in the oblong Figure P T MN; and then viewing them through a Prism D H held Parallel to their length, they appeared not co-incident as when viewed with the naked Eye, but in the form of two diftinct Spectrums pt and mn croffing one another in the middle after the manner of the letter X. Which shews that the red of the one Spectrum and violet of the other, which were co-incident at P N and M T, being parted from one another by a greater Refraction of the violet to p and m than of the red to n and t, do differ in degrees of Refrangibility.

I illuminated also a little circular piece of white Paper all over with the Lights of both Prisms intermixed, and when

when it was illuminated with the red of one Spectrum and deep violet of the other, so as by the mixture of those Colours to appear all over purple, I viewed the Paper, first at a less distance, and then at a greater, through a third Prism; and as I went from the Paper, the refracted Image thereof became more and more divided by the unequal Refraction of the two mixed Colours, and at length parted into two distinct Images, a red one and a violet one, whereof the violet was furthest from the Paper, and therefore suffered the greatest Refraction. And when that Prism at the Window which cast the violet on the Paper was taken away, the violet Image disappeared; but when the other Prism was taken away the red vanished : which shews that these two Images were nothing else than the Lights of the two Prisms which had been intermixed on the purple Paper, but were parted again by their unequal Refractions made in the third Prism through which the Paper was viewed. This also was observable that if one of the Prisms at the Window, suppose that which cast the violet on the Paper, was turned about its Axis to make all the Colours in this order, Violet, Indigo, Blew, Green, Yellow, Orange, Red, fall successively on the Paper from that Prism, the violet Image changed Colour accordingly, and in changing Colour came nearer to the red one, until when it was also red they both became fully co-incident.

I placed also two paper circles very near one another, the one in the red Light of one Prism, and the other in the violet Light of the other. The circles were each of them an Inch in Diameter, and behind them the Wall was dark that the Experiment might not be disturbed by any Light coming from thence. These circles thus illuminated, I viewed through a Prism so held that the Refraction might be made towards the red circle, and as I went from them they

they came nearer and nearer together, and at length became co-incident; and afterwards when I went still further off, they parted again in a contrary order, the violet by a greater Refraction being carried beyond the red.

Exper. 8. In Summer when the Sun's Light uses to be strongest, I placed a Prism at the hole of the Windowshut, as in the third Experiment, yet so that its Axis might be Parallel to the Axis of the World, and at the opposite Wall in the Sun's refracted Light, I placed an open Book. Then going Six Feet and twoo Inches from the Book, I placed there the abovementioned Lens, by vvhich the Light reflected from the Book might be made to converge and meet again at the diftance of fix Feet and tvvo Inches behind the Lens, and there paint the Species of the Book upon a sheet of vvhite Paper much after the manner of the second Experiment. The Book and Lens being made fast, I noted the place vvhere the Paper vvas, vvhen the Letters of the Book, illuminated by the fullest red Light of the Solar Image falling upon it, did caft their Species on that Paper most distinctly'; And then I stay'd till by the Motion of the Sun and consequent Motion of his Image on the Book, all the Colours from that red to the middle of the blew pass'd over those Letters; and when those Letters were illuminated by that blew, I noted again the place of the Paper when they cast their Species most distinctly upon it : And I found that this last place of the Paper was nearer to the Lens than its former place by about two Inches and an half, or two and three quarters. So much sooner therefore did the Light in the violet end of the Image by a greater Refraction converge and meet, than the Light in the red end. But in trying this the Chamber was as dark as I could make it. For if these Colours be diluted and weakned by the mixture of any adventitious Light, the distance between

between the places of the Paper will not be fo great. This diffance in the fecond Experiment where the Colours of natural Bodies were made ufe of, was but an Inch and a half, by reafon of the imperfection of those Colours. Here in the Colours of the Prifm, which are manifestly more full, intense, and lively than those of natural Bodies, the distance is two Inches and three quarters. And were the Colours still more full, I question not but that the diftance would be confiderably greater. For the coloured Light of the Prism, by the interfering of the Circles defcribed in the 11th Figure of the fifth Experiment, and also by the Light of the very bright Clouds next the Sun's Body intermixing with these Colours, and by the Light fcattered by the inequalities in the polish of the Prism, was fo very much compounded, that the Species which those faint and dark Colours, the Indigo and Violet, cast upon the Paper were not distinct enough to be well observed.

Exper. 9. A Prism, whose two Angles at its Base were equal to one another and half right ones, and the third a right one, I placed in a beam of the Sun's Light let into a dark Chamber through a hole in the Window-fhut as in the third Experiment. And turning the Prism flowly about its Axis until all the Light which went through one of its Angles and was refracted by it began to be reflected by its Base, at which till then it went out of the Glass, I observed that those Rays which had suffered the greatest Refraction were sooner reflected than the rest. I conceived therefore that those Rays of the reflected Light, which were most Refrangible, did first of all by a total Reflexion become more copious in that Light than the rest, and that afterwards the rest also, by a total Reslexion, became as copious as these. To try this, I made the reflected Light pass through another Prism, and being refracted

Eted by it to fall afterwards upon a sheet of white Paper placed at some distance behind it, and there by that Refraction to paint the usual Colours of the Prism. And then causing the first Prism to be turned about its Axis as above, I observed that when those Rays which in this Prism had suffered the greatest Refraction and appeared of a blew and violet Colour began to be totally reflected, the blew and violet Light on the Paper which was most refracted in the fecond Prism received a sensible increase above that of the red and yellow, which was least refracted; and afterwards when the rest of the Light which was green, yellow and red began to be totally reflected in the first Prism, the light of those Colours on the Paper received as great an increase as the violet and blew had done before. Whence 'tis manifest, that the beam of Light reflected by the Base of the Prism, being augmented first by the more Refrangible Rays and afterwards by the less Refrangible ones, is compounded of Rays differently Refrangible. And that all such reflected Light is of the same Nature with the Sun's Light, before its Incidence on the Base of the Prism, no Man ever doubted : it being generally allowed, that Light by fuch Reflexions suffers no Alteration in its Modifications and Properties. I do not here take notice of any Refractions made in the Sides of the first Prism, because the Light enters it perpendicularly at the first Side, and goes out perpendicularly at the second Side, and therefore suffers none. So then, the Sun's incident Light being of the fame temper and conftitution with his emergent Light, and the last being compounded of Rays differently Refrangible, the first must be in like manner compounded.

Fig. 21.

1. Illustration. In the 21th Figure, ABC is the first Prism, BC its Base, B and C its equal Angles at the Base, each of 45 degrees, A its Rectangular Vertex, FM a beam of the Sun's Light let into a dark Room through a hole F one third part of an Inch broad, M its Incidence on the Base of the Prism, MG a less refracted Ray, MH a more refracted Ray, MN the beam of Light reflected from the Base, V X Y the fecond Prism by which this beam in passing through it is refracted, Nt the less refracted Light of this beam, and N p the more refracted part thereof. When the first Prism A B C is turned about its Axis according to the order of the Letters ABC, the Rays MH emerge more and more obliquely out of that Prism, and at length after their most oblique Emergence are reflected towards N, and going on to p do increase the number of the Rays N p. Aftervvards by continuing the motion of the first Prism, the Rays MG are also reflected to N and increase the number of the Rays N t. And therefore the Light M N admits into its Composition, first the more Refrangible Rays, and then the less Refrangible Rays, and yet after this Composition is of the same Nature vvith the Sun's immediate Light F M, the Reflexion of the specular Base BC causing no Alteration therein.

*Exper.* 10. Two Prifms, which were alike in fhape, I tied fo together, that their Axes and oppofite Sides being Parallel, they composed a Parallelopiped. And, the Sun fhining into my dark Chamber through a little hole in the Window-fhut, I placed that Parallelopiped in his beam at fome diftance from the hole, in fuch a posture that the Axes of the Prifms might be perpendicular to the incident Rays, and that those Rays being incident upon the first Side of one Prifm, might go on through the two contiguous Sides of both Prifms, and emerge out of the last Side of the first Side of the first Prifm. This Side being Parallel to the first Side of the first Prifm, caused the emerging Light to be Parallel

to.

to the Incident. Then, beyond these two Prisms I placed a third, which might refract that emergent Light, and by that Refraction cast the usual Colours of the Prism upon the opposite Wall, or upon a sheet of white Paper held at a convenient distance behind the Prism for that refracted Light to fall upon it. After this I turned the Parallelopiped about its Axis, and found that when the contiguous Sides of the two Prisms became so oblique to the incident Rays that those Rays began all of them to be reflected, those Rays which in the third Prism had suffered the greatest Refraction and painted the Paper with violet and blew, were first of all by a total Reflexion taken out of the transmitted Light, the rest remaining and on the Paper painting their Colours of Green, Yellow, Orange, and Red as before; and afterwards by continuing the motion of the two Prisms, the rest of the Rays also by a total Reflexion vanished in order, according to their degrees of Refrangibility. The Light therefore which emerged out of the two Prisms is compounded of Rays differently Refrangible, seeing the more Refrangible Rays may be taken out of it while the less Refrangible remain. But this Light being trajected only through the Parallel Superficies of the two Prisms, if it suffered any change by the Refraction of one Superficies it lost that impression by the contrary Refraction of the other Superficies, and so being restored to its pristine constitution became of the same nature and condition as at first before its Incidence on those Prisms; and therefore, before its Incidence, was as much compounded of Rays differently Refrangible as afterwards.

Fig. 22.

Illustration. In the 22th Figure A B C and B C D are the the two Prisms tied together in the form of a Parallelopiped, their Sides B C and C B being contiguous, and their Sides A B and C D Parallel. And H J K is the third Prism,

## [40]

Prism, by which the Sun's Light propagated through the hole F into the dark Chamber, and there passing through those fides of the Prisms AB, BC, CB and CD, is refracted at O to the white Paper P T, falling there partly upon P by a greater Refraction, partly upon T by a less Refraction, and partly upon R and other intermediate places by intermediate Refractions. By turning the Parallelopiped ACBD about its Axis, according to the order of the Letters A, C, D, B, at length when the contiguous Planes BC and CB become sufficiently oblique to the Rays F M, which are incident upon them at M, there will vanish totally out of the refracted Light OPT, first of all the most refracted Rays OP, (the reft OR and OT remaining as before) then the Rays OR and other intermediate ones, and lastly, the least refracted Rays OT. For when the Plane BC becomes sufficiently oblique to the Rays incident upon it, those Rays will begin to be totally reflected by it towards N; and first the most Refrangible Rays will be totally reflected (as was explained in the preceding experiment) and by confequence must first disappear at P, and afterwards the rest as they are in order totally reflected to N, they must disappear in the same order at R and T. So then the Rays which at O fuffer the greatest Refraction, may be taken out of the Light MO whilst the rest of the Rays remain in it, and therefore that Light MO is Compounded of Rays differently Refrangible. And because the Planes A B and C D are parallel, and therefore by equal and contrary Refractions destroy one anothers Effects, the incident Light FM must be of the same kind and nature with the emergent Light MO, and therefore doth also consist of Rays differently Refrangible. These two Lights FM and MO, before the most retrangible Rays are separated out of the emergent Light MO agree in Colour,

lour, and in all other properties so far as my observation reaches, and therefore are deservedly reputed of the same Nature and Constitution, and by consequence the one is compounded as well as the other. But after the most Re-frangible Rays begin to be totally reflected, and thereby separated out of the emergentLight MO, that Light changes its Colour from white to a dilute and faint yellow, a pretty good orange, a very full red fucceffively and then totally vanishes. For after the most Refrangible Rays which paint the Paper at P with a Purple Colour, are by a total reflexion taken out of the Beam of light MO, the rest of the Colours which appear on the Paper at R and T being mixed in the light MO compound there a faint yellow, and after the blue and part of the green which appear on the Paper between P and R are taken away, the rest which a appear between R and T (that is the Yellow, Orange, Red and a little Green) being mixed in the Beam MO compound there an Orange; and when all the Rays are by re-Hexiontaken out of the Beam MO, except the least Refrangible, which at T appear of a full Red, their Colour is the same in that Beam MO as afterwards at T, the Refraction of the Prism HJK serving only to separate the differently Refrangible Rays, without making any alteration in their Colours, as shall be more fully proved hereafter.... All which confirms as well the first Proposition as the fecond.

Scholium. If this Experiment and the former be conjoyned Fig. 22. and made one, by applying a fourth Prifm VXY to refract the reflected Beam M N towards tp, the conclusion will be clearer. For then the light Np which in the 4th Prifm is more refracted, will become fuller and stronger when the Light O P, which in the third Prifm HJK is more refracted, vanishes at P; and afterwards when the lefs refracted

refracted Light O T vanishes at T, the less refracted Light Nt will become encreased whilst the more refracted Light at p receives no further encrease. And as the trajected Beam MO in vanishing is always of such a Colour as ought to refult from the mixture of the Colours which fall upon the Paper PT, so is the reflected Beam MN always of such a Colour as ought to result from the mixture of the Colours which fall upon the Paper pt. For when the most refrangible Rays are by a total Reflexion taken out of the Beam MO, and leave that Beam of an Orange Colour, the excels of those Rays in the reflected Light, does not only make the Violet, Indigo and Blue at p more full, but also makes the Beam MN change from the yellowish Colour of the Sun's Light, to a pale white inclining to blue, and afterward recover its yellowish Colour again, so soon as all the rest of the transmitted light MOT is reflected.

Now feeing that in all this variety of Experiments, whether the trial be made in Light reflected, and that either from natural Bodies, as in the first and second Experiment, or Specular, as in the Ninth; or in Light refracted, and that either before the unequally refracted Rays are by diverging separated from one another, and losing their whiteness which they have altogether, appear severally of several Colours, as in the fifth Experiment; or after they are feparated from one another, and appear Coloured as in the fixth, feventh, and eighth Experiments; or in Light tra-jected through Parallel fuperficies, deftroying each others Effects as in the 10th Experiment; there are always found Rays, which at equal Incidences on the same Medium suffer unequal Refractions, and that without any splitting or dilating of single Rays, or contingence in the inequality of the Refractions, as is proved in the fifth and fixth Experiments ;

F 2

periments; and feeing the Rays which differ in Refrangibility may be parted and forted from one another, and that either by Refraction as in the third Experiment, or by Reflexion as in the tenth, and then the feveral forts apart at equal Incidences fuffer unequal Refractions, and those forts are more refracted than others after feparation, which were more refracted before it, as in the fixth and following Experiments, and if the Sun's Light be trajected through three or more crofs Prisms fucceffively, those Rays which in the first Prism are refracted more than others are in all the following Prisms, refracted more then others in the fame rate and proportion, as appears by the fifth Experiment; it's manifest that the Sun's Light is an Heterogeneous mixture of Rays, fome of which are constantly more Refrangible then. others, as was to be proposed.

### PROP. III. Theor. III.

The Sun's Light confifts of Rays differing in Reflexibility, and those Rays are more Reflexible than others which are more Refrangible.

HIS is manifest by the ninth and tenth Experiments: For in the ninth Experiment, by turning the Prism about its Axis, until the Rays within it which in going out into the Air were refracted by its Base, became so oblique to that Base, as to begin to be totally reflected thereby; those Rays became first of all totally reflected, which before at equal Incidences with the rest had suffered the greatest Refraction. And the same thing happens in the Reflexion made by the common Base of the two Prisms in the tenth Experiment.

PROP.

# [45]

## PROP. IV. Prob. I.

#### To Separate from one another the Heterogeneous Rays of Compound Light.

THE Heterogeneous Rays are in some measure sepa-rated from one another by the Refraction of the Prism in the third Experiment, and in the fifth Experiment by taking away the Penumbra from the Rectilinear fides of the Coloured Image, that separation in those very Rectilinear sides or straight edges of the Image becomes perfect. But in all places between those rectilinear edges, those innumerable Circles there described, which are severally illuminated by Homogeneral Rays, by interfering with one another, and being every where commixt, do render the Light sufficiently Compound. But if these Circles, whilst their Centers keep their distances and positions, could be made less in Diameter, their interfering one with another and by confequence the mixture of the Heterogeneous-Rays would be proportionally diminished. In the 23th Fig. 23. Figure let A G, B H, C J, D K, EL, F M be the Circles which so many forts of Rays flowing from the same Disque of the Sun, do in the third Experiment illuminate; of all which and innumerable other intermediate ones lying in a continual Series between the two Rectilinear and Paralleledges of the Sun's oblong Image P T, that Image is composed as was explained in the fifth Experiment. And let ag, bh, ci, dk, el, fm be so many less Circles lying in a like continual Series between two Parallel right Lines a f and gm with the fame distances between their Centers, and illuminated by the fame forts of Rays, that is the Circle ag with the fame fort by which the corresponding Circle

Circle AG was illuminated, and the Circle bh with the fame fort by which the corresponding Circle BH was illuminated, and the rest of the Circles ci, dk, el, fm respectively, with the fame forts of Rays by which the feveral correfponding Circles CJ, DK, EL, FM were illuminated. In the Figure PT composed of the greater Circles, three of those Circles A G, B H, CJ, are so expanded into one another, that the three forts of Rays by which those Circles are illuminated, together with other innumerable forts of intermediate Rays, are mixed at QR in the middle of the Circle B H. And the like mixture happens throughout almost the whole length of the Figure P T. But in the Figure pt composed of the less Circles, the three less Circles ag, bh, c i, which answer to those three greater, do not extend into one another; nor are there any where mingled so much as any two of the three forts of Rays by which those Circles are illuminated, and which in the Figure P T are all of them intermingled at B H.

Now he that fhall thus confider it, will eafily underftand that the mixture is diminifhed in the fame Proportion with the Diameters of the Circles. If the Diameters of the Circles whilft their Centers remain the fame, be made three times lefs than before, the mixture will be alfo three times lefs; if ten times lefs, the mixture will be ten times lefs, and fo of other Proportions. That is, the mixture of the Rays in the greater Figure P T will be to their mixture in the lefs pt, as the Latitude of the greater Figure is to the Latitude of the lefs. For the Latitudes of thefe Figures are equal to the Diameters of their Circles. And hence it eafily follows, that the mixture of the Rays in the refracted Spectrum pt is to the mixture of the Rays in the direct and immediate Light of the Sun, as the breadth of that Spectrum is to the difference between the length and breadth of the fame Spectrum. Widthird: MyMMAR Sed, So

So then, if we would diminish the mixture of the Rays, we are to diminish the Diameters of the Circles. Now these would be diminished if the Sun's Diameter to which they answer could be made less than it is, or (which comes to the fame purpose) if without Doors, at a great distance from the Prism towards the Sun, some opake body were placed, with a round hole in the middle of it, to intercept all the Sun's Light, excepting fo much as coming from the middle of his Body could pass through that hole to the Prism. For so the Circles AG, BH and the rest, would not any longer answer to the whole Disque of the Sun, but only to that part of it which could be feen from the Prism through that hole, that is to the apparent magnitude of that hole viewed from the Prism. But that these Circles may answer more distinctly to that hole a Lens is to be placed by the Prism to cast the Image of the hole, (that is, every one of the Circles A G, B H, Gc.) distinctly upon the Paper at P T, after such a manner as by a Lens placed at a Window the Species of Objects abroad are cast distinctly upon a Paper within the Room, and the Rectilinear Sides of the oblong folar Image in the fifth Experiment became distinct without any Penumbra. If this be done it will not be necessary to place that hole very far off, no not beyond the Window. And therefore instead of that hole, I used the hole in the Window-shut as follows.

Exper. 11. In the Sun's Light let into my darkned Chamber through a fmall round hole in my Windowfhut, at about 10 or 12 Feet from the Window, I placed a Lens, by which the Image of the hole might be diftinctly caft upon a fheet of white Paper, placed at the diftance of fix, eight, ten or twelve Feet from the Lens. For according to the difference of the Lenfes I ufed various diftances,

distances, which I think not worth the while to describe. Then immediately after the Lens I placed a Prism, by which the trajected Light might be refracted either upwards or fideways, and thereby the round Image which the Lens alone did cast upon the Paper might be drawn out into a long one with Parallel Sides, as in the third Experiment. This oblong Image I let fall upon another Paper at about the fame diftance from the Prism as before, moving the Paper either towards the Prism or from it, until I found the just distance where the Rectilinear Sides of the Image became most distinct. For in this case the circular Images of the hole which compose that Image after the fame manner that the Circles ag, bh, ci, &c. do Fig. 23. the Figure pt, were terminated most distinctly without any Penumbra, and therefore extended into one another the least that they could, and by consequence the mixture of the Heterogeneous Rays was now the least of all. By this Fig. 23, means I used to form an oblong Image (such as is pt) of and 24. circular Images of the hole (fuch as are ag, bh, ci, &c.) and by using a greater or less hole in the Window-shut, I made the circular Images ag, bh, ci, &c. of which it was formed, to become greater or less at pleasure, and thereby the mixture of the Rays in the Image pt to be as much or as little as I defired.

Fig. 24. Illustration. In the 24th Figure, F represents the circular hole in the Window-shut, M N the Lens whereby the Image or Species of that hole is cast distinctly upon a Paper at J, A B C the Prism whereby the Rays are at their emerging out of the Lens refracted from J towards another Paper at pt, and the round Image at J is turned into an oblong Image pt falling on that other Paper. This Image pt consists of Circles placed one after another in a Rectilinear order, as was sufficiently explained in the fifth Experiment;

Experiment; and these Circles are equal to the Circle I, and confequently answer in Magnitude to the hole F; and therefore by diminishing that hole they may be at pleasure diminished, whil'st their Centers remain in their places. By this means I made the breadth of the Image pt to be forty times, and sometimes fixty or seventy times less than its length. As for instance, if the breadth of the hole F be i of an Inch, and MF the diftance of the Lens from the hole be 12 Feet; and if p B or p M the distance of the Image pt from the Prism or Lens be 10 Feet, and the refracting Angle of the Prism be 62 degrees, the breadth of the Image p t will be  $\frac{1}{12}$  of an Inch and the length about fix Inches, and therefore the length to the breadth as 72 to 1, and by consequence the Light of this Image 71 times less compound than the Sun's direct Light. And Light \* thus far Simple and Homogeneal, is sufficient for trying all the Experiments in this Book about fimple Light. For the composition of Heterogeneal Rays is in this Light fo little that it is scarce to be discovered and perceived by sense, except perhaps in the Indigo and Violet; for these being dark Colours, do easily suffer a sensible allay by that little scattering Light which uses to be refracted irregularly by the inequaliteis of the Prism.

Yet instead of the circular hole F, 'tis better to substitute an oblong hole shaped like a long Parallelogram with its length Parallel to the Prism A B C. For if this hole be an Inch or two long, and but a tenth or twentieth part of an Inch broad or narrower : the Light of the Image pt will be as Simple as before or fimpler, and the Image will become much broader, and therefore more fit to have Experiments tried in its Light than before.

Instead of this Parallelogram-hole may be substituted a Triangular one of equal Sides, whose Base for instance is about

about the tenth part of an Inch, and its height an Inch or more. For by this means, if the Axis of the Prifm be Parallel to the Perpendicular of the Triangle, the Image Fig. 25. pt will now be formed of Equicrural Triangles ag, bh, ci, dk, el, fm, &cc. and innumerable other intermediate ones anfwering to the Triangular hole in fhape and bignefs, and lying one after another in a continual Series between two Parallel Lines af and gm. Thefe Triangles are a little intermingled at their Bafes but not at their Vertices, and therefore the Light on the brighter fide af of the Image where the Bafes of the Triangles are is a little compounded, but on the darker fide gm is altogether uncompounded, and in all places between the fides the Composition is Proportional to the diffances of the places from that obfcurer fide gm. And having a Spectrum pt of fuch a Composition, we may try Experiments either in its ftronger and lefs fimple Light near the fide af, or in its weaker and fimpler Light near the other fide lm, as it fhall feam moft convenient.

[50]

But in making Experiments of this kind the Chamber ought to be made as dark as can be, leaft any forreign Light mingle it felf with the Light of the Spectrum pt, and render it compound; efpecially if we would try Experiments in the more fimple Light next the fide gm of the Spectrum; which being fainter, will have a lefs Proportion to the forreign Light, and fo by the mixture of that Light be more troubled and made more compound. The Lens alfo ought to be good, fuch as may ferve for Optical Ufes, and the Prifm ought to have a large Angle, fuppofe of 70 degrees, and to be well wrought, being made of Glafs free from Bubbles and Veins, with its fides not a little Convex or Concave as ufually happens but truly Plane, and its pollifh elaborate, as in working Optickglaffes, glasses, and not such as is usually wrought with Putty, whereby the edges of the Sand-holes being worn away, there are left all over the Glass a numberless company of very little Convex polite rifings like Waves. The edges also of the Prism and Lens so far as they may make any irregular Refraction, must be covered with a black Paper glewed on. And all the Light of the Sun's beam let into the Chamber which is uselels and unprofitable to the Experiment, ought to be intercepted with black Paper or other black Obstacles. For otherwise the useless Light being reflected every way in the Chamber, will mix with the oblong Spectrum and help to difturb it. In trying these things so much Diligence is not altogether necessary, but it will promote the success of the Experiments, and by a very scrupulous Examiner of things deserves to be applied. It's difficult to get glass Prisms fit for this purpose, and and therefore I used sometimes Prismatick Vessels made with pieces of broken Looking-glasses, and filled with rain Water. And to increase the Refraction, I sometimes impregnated the Water strongly with Saccharum Saturni.

# PROP. V. Theor. IV.

Homogeneal Light is refracted regularly without any Dilatation Splitting or Shattering of the Rays, and the confused Vision of Objects seen through Refracting Bodies by Heterogeneal Light arises from the different Refrangibility of several sorts of Rays.

H E first Part of this Proposition has been already fufficiently proved in the fifth Experiment, and will further appear by the Experiments which follow. Exper. 12. In the middle of a black Paper I made a round hole about a fifth or fixth part of an Inch in Diameter. Upon this Paper I caufed the Spectrum of Homogeneal Light defcribed in the former Proposition, fo to fall, that some part of the Light might pass through the hole of the Paper. This transmitted part of the Light I refracted with a Prism placed behind the Paper, and letting this refracted Light fall perpendicularly upon a white Paper two or three Feet distant from the Prism, I found that the Spectrum formed on the Paper by this Light was not oblong, as when 'tis made (in the third Experiment) by Refracting the Sun's compound Light, but was (so far as I could judge by my Eye) perfectly circular, the length being no greater than the breadth. Which shews that this Light is refracted regularly without any Dilatation of the Rays.

*Exper.* 13. In the Homogeneal Light I placed a Circle of  $\frac{1}{4}$  of an Inch in Diameter, and in the Sun's unrefracted Heterogeneal white Light I placed another Paper Circle of the fame bignefs. And going from the Papers to the diftance of fome Feet, I viewed both Circles through a Prifm. The Circle illuminated by the Sun's Heterogeneal Light appeared very oblong as in the fourth Experiment, the length being many times greater than the breadth : but the other Circle illuminated with Homogeneal Light appeared Circular and diffinctly defined as when 'tis viewed with the naked Eye. Which proves the whole Proposition.

Exper. 14. In the Homogeneal Light I placed Flies and fuch like Minute Objects, and viewing them through a Prifm, I faw their Parts as diffinctly defined as if I had viewed them with the naked Eye. The fame Objects placed in the Sun's unrefracted Heterogeneal Light which was white I viewed alfo through a Prifm, and faw them moft confufedly

confusedly defined, so that I could not distinguish their smaller Parts from one another. I placed also the Letters of a fmall Print one while in the Homogeneal Light and then in the Heterogeneal, and viewing them through a Prifm, they appeared in the latter case so confused and indistinct that I could not read them ; but in the former they appeared fo diftinct that I could read readily, and thought I faw them as diffinct as when I viewed them with my naked Eye. In both cafes I viewed the fame Objects through the same Prism at the same distance from me and in the same Situation. There was no difference but in the Light by which the Objects were illuminated, and which in one cafe was Simple and in the other Compound, and therefore the diftinct Vision in the former case and confused in the latter could arise from nothing else than from that difference of the Lights. Which proves the whole Proposition.

And in these three Experiments it is further very remarkable, that the Colour of Homogeneal Light was neverchanged by the Refraction.

## PROP. VI. Theor. V.

The Sine of Incidence of every Ray confidered apart, is to its Sine of Refraction in a given Ratio.

THAT every Ray confidered apart is conftant to it felf in fome certain degree of Refrangibility, is fufficiently manifest out of what has been faid. Those Rays which in the first Refraction are at equal Incidences most refracted, are also in the following Refractions at equal Incidences most refracted; and so of the least Refrangible, and the rest which have any mean degree of RefranRefrangibility, as is manifest by the 5th, 6th, 7th, 8th, and 9th Experiments. And those which the first time at like Incidences are equally refracted, are again at like Incidences equally and uniformly refracted, and that whether they be refracted before they be separated from one another as in the 5th Experiment, or whether they be refracted apart, as in the 12th, 13th and 14th Experiments. The Refraction therefore of every Ray apart is regular, and what Rule that Refraction observes we are now to show.

The late Writers in Opticks teach, that the Sines of Incidence are in a given Proportion to the Sines of Refraction, as was explained in the 5th Axiom; and some by Instruments fitted for measuring Refractions, or otherwife experimentally examining this Proportion, do acquaint us that they have found it accurate. But whilst they, not understanding the different Refrangibility of several Rays, conceived them all to be refracted according to one and the same Proportion, 'tis to be presumed that they adapted their Measures only to the middle of the refracted Light; fo that from their Measures we may conclude only that the Rays which have a mean degree of Refrangibility, that is those which when separated from the rest appear green, are refracted according to a given Proportion of their Sines. And therefore we are now to fhew that the like given Proportions obtain in all the reft. That it should be so is very reasonable, Nature being ever conformable to her self: but an experimental Proof is desired. And fuch a Proof will be had if we can shew that the Sines of Refraction of Rays differently Refrangible are one to another in a given Proportion when their Sines of Incidence are equal. For if the Sines of Refraction of all the Rays are in given Proportions to the Sine of Refraction of

of a Ray which has a mean degree of Refrangibility, and this Sine is in a given Proportion to the equal Sines of Incidence, those other Sines of Refraction will also be in given Proportions to the equal Sines of Incidence. Now when the Sines of Incidence are equal, it will appear by the following Experiment that the Sines of Refraction are in a given Proportion to one another.

Exper. 15. The Sun shining into a dark Chamber through a little round hole in the Window-fhut, let S re-Fig. 26. present his round white Image painted on the opposite Wall by his direct Light, P T his oblong coloured Image made by refracting that Light with a Prism placed at the Window; and pt, or 2p 2t, or 3p 3t, hisoblong coloured Image made by refracting again the same Light sideways with a fecond Prism placed immediately after the first in a cross Position to it, as was explained in the fifth Experiment : that is to fay, pt when the Refraction of the second Prism is small, 2 p 2 t when its Refraction is greater, and 3 p 3 t when it is greatest. For such will be the diversity X of the Refractions if the refracting Angle of the second Prism be of various Magnitudes ; suppose of fifteen or twenty degrees to make the Image p t, of thirty or forty to make the Image 2 p 2 t, and of fixty to make the Image 3 p 3 t. But for want of solid Glass Prisms with Angles of convenient bignesses, there may be Vessels made of polished Plates of Glass cemented together in the form of Prisms and filled with Water. These things being thus ordered, I observed that all the solar Images or coloured Spectrums P T, pt, 2p 2t, 3p 3t did very nearly converge to the place S on which the direct Light of the Sun fell and painted his white round Image when the Prisms were taken away. The Axis of the Spectrum PT, that is the Line drawn through the middle of it Parallel to

its

its Rectilinear Sides, did when produced pass exactly through the middle of that white round Image S. And when the Refraction of the second Prism was equal to the Refraction of the first, the refracting Angles of them both being about 60 degrees, the Axis of the Spectrum 3 p 3 t made by that Refraction, did when produced pass also through the middle of the fame white round Image S. But when the Refraction of the fecond Prism was less than that of the first, the produced Axes of the Spectrums tp or 2t 2p made by that Refraction did cut the produced Axis of the Spe-Etrum TP in the Points m and n, a little beyond the Center of that white round Image S. Whence the Proportion of the Line 3t T to the Line 3p P was a little greater than the Proportion of 2tT to 2pP, and this Proportion a little greater than that of t T to pP. Now when the Light of the Spectrum P T falls perpendicularly upon the Wall, those Lines 3tT, 3pP, and 2tT, 2pP and tT, pP, are the Tangents of the Refractions; and therefore by this Experiment the Proportions of the Tangents of the Refractions are obtained, from whence the Proportions of the Sines being derived, they come out equal, so far as by viewing the Spectrums and using some Mathematical reasoning I could Estimate. For I did not make an Accurate Computation. So then the Proposition holds true in every Ray apart, so far as appears by Experiment. And that it is accurately true may be demonstrated upon this Supposition, That Bodies refract Light by acting upon its Rays in Lines Perpendicular to their Surfaces. But in order to this Demonstration, I must diftinguish the Motion of every Ray into two Motions, the one Perpendicular to the refracting Surface, the other Parallel to it, and concerning the Perpendicular Motion lay down the following Proposition.

-X-

If any Motion or moving thing whatfoever be incident With the Part matter with any velocity on any broad and thin Space terminated on both fides by two Parallel Planes, and in its paffage through that fpace be urged perpendicularly towards the further Plane by any force which at given diftances from the Plane is of given quantities; the perpendicular Velocity of that Motion or Thing, at its emerging out of that fpace, fhall be always equal to the Square Root of the Summ of the Square of the perpendicular Velocity of that Motion or Thing at its Incidence on that fpace; and of the Square of the perpendicular Velocity which that Motion or Thing would have at its Emergence, if at its Incidence its perpendicular Velocity was infinitely little.

And the same Proposition holds true of any Motion or Thing perpendicularly retarded in its passage through that space, if instead of the Summ of the two Squares you take their difference. The Demonstration Mathematicians will Wid. Mawl: Prin. Math. easily find out, and therefore I shall not trouble the Rea-

Suppose now that a Ray coming most obliquely in the Fig. 1. Line MC be refracted at C by the Plane RS into the Line CN, and if it be required to find the Line CE into which any other Ray AC shall be refracted; let MC, AD, be the Sines of incidence of the two Rays, and NG, EF, their Sines of Refraction, and let the equal Motions of the Incident Rays be represented by the equal Lines M C and AC, and the Motion MC being confidered as parallel to the refracting Plane, let the other Motion AC be distinguissed into two Motions AD and DC, one of which AD is parallel, and the other DC perpendicular to the refracting Surface. In like manner, let the Motions of the emering Rays be distinguissed into two, whereof the per-H pendicular

## [58]

perpendicular ones are  $\frac{MC}{NG}$  CG and  $\frac{AD}{EF}$  CF. And if the force of the refracting Plane begins to act upon the Rays either in that Plane or at a certain distance from it on the one fide, and ends at a certain distance from it on the other side, and in all places between those two Limits acts upon the Rays in Lines perpendicular to that rafracting Plane, and the Actions upon the Rays at equal diffances from the refracting Plane be equal, and at unequal ones either equal or unequal according to any rate whatever; that motion of the Ray which is Parallel to the refracting Plane will suffer no alteration by that force; and that motion which is perpendicular to it will be altered according to the rule of the foregoing Propolition. If therefore for the perpendicular Velocity of the emerging Ray CN you write  $\frac{MC}{NG}$  CG as above, then the perpendicular Velocity of any other emerging Ray CE which was  $\frac{AD}{EF}$  CF, will be equal to the square Root of  $CD_q + \frac{MC_q}{NG_q} CG_q$ . And by squaring these equals, and adding to them the Equals ADq and MCq---CDq, and dividing the Summs by the Equals CFq + EFq and CGq + NGq, you will have  $\frac{AD}{EFq}$  equal to  $\frac{MCq}{NGq}$ . Whence AD, the Sine of Incidence, is to EF the Sine of Refraction, as MC to NG, that is, in a given ratio. And this Demonstration being general, without determining what Light is, or by what kind of force it is refracted, or affuming any thing further than that the refracting Body acts upon the Rays in Lines perpendicular to its Surface; I take it to be a very convincing Argument of the full Truth of this Proposition.

So

So then, if the ratio of the Sines of Incidence and Refraction of any fort of Rays be found in any one Cafe, 'tis given in all Cafes; and this may be readily found by the Method in the following Proposition.

### PROP. VII. Theor. VI.

#### The Perfection of Telescopes is impeded by the different Refrangibility of the Rays of Light.

THE imperfection of Telescopes is vulgarly attri-buted to the spherical Figures of the Glasses, and therefore Mathematicians have propounded to Figure them by the Conical Sections. To fhew that they are mistaken, I have inferted this Proposition; the truth of which will appear by the measures of the Refractions of the several forts of Rays; and these measures I thus determine.

In the third experiment of the first Book, where the refracting Angle of the Prism was 621 degrees, the half of that Angle 31 deg. 15 min. is the Angle of Incidence of the Rays at their going out of the Glass into the Air; and the Sine of this Angle is 5188, the Radius being 10000. 5187. fere. When the Axis of this Prism was parallel to the Horizon, and the Refraction of the Rays at their Incidence on this Prism equal to that at their Emergence out of it, I observed with a Quadrant the Angle which the mean refrangible Rays (that is, those which went to the middle of the Sun's coloured Image) made with the Horizon and by this Angle and the Sun's altitude observed at the same time, I found the Angle which the emergent Rays contained with the incident to be 44 deg. and 40 min. and the half of this Angle added to the Angle of Incidence 31 deg. 15 min. makes the H 2 Angle

Angle of Refraction, which is therefore 53 deg. 35 min. and its Sine 8047. These are the Sines of Incidence and Refraction of the mean refrangible Rays, and their proportion in round numbers is 20 to 31. This Glass was of a colour inclining to green. The last of the Prisms mentioned in the third Experiment was of clear white Glass. Its refracting Angle 63<sup>1</sup>/<sub>2</sub> degrees. The Angle which the emergent Rays contained, with the incident 45 deg. 50 min. The Sine of half the first Angle 5262. The Sine of half the Summ of the Angles 8157. And their proportion in round numbers 20 to 31 as before.

From the Length of the Image, which was about 93 or ro Inches, subduct its Breadth, which was 2 1/8 Inches, and the Remainder 73 Inches would be the length of the Image were the Sun but a point, and therefore subtends the Angle which the most and least refrangible Rays, when inci-dent on the Prism in the same Lines, do contain with one another after their Emergence. Whence this Angle is 2 deg. 0. 7." For the diftance between the Image and the Prism where this Angle is made; was  $18\frac{1}{2}$  Feet, and at that distance the Chord 74 Inches subtends an Angle of 2 deg. 0.'7." Now half this Angle is the Angle which these emergent Rays contain with the emergent mean refrangible Rays, and a quarter thereof, that is 30. 2." may be accounted the Angle which they would contain which the same emergent mean refrangible Rays, were they co-incident to them within the Glass and suffered no other Refraction then that at their Emergence. For if two equal Refractions, the one at the incidence of the Rays on the Prism, the other at their Emergence, make half the Angle 2 deg. 0. 7." then one of those Refractions will make about a quarter of that Angle, and this quarter added to and
and fubducted from the Angle of Refraction of the mean refrangible Rays, which was 53 deg. 35', gives the Angles of Refraction of the most and least refrangible Rays 54 deg. 5' 2", and 53 deg. 4' 58", whose Sines are 8099 and 7995, the common Angle of Incidence being 31 deg. 15' and its Sine 5188; and these Sines in the least round numbers are in proportion to one another as 78 and 77 to 50.

Now if you fubduct the common Sine of Incidence 50° +from the Sines of Refraction 77 and 78, the remainders 27 and 28 fhew that in fmall Refractions the Refraction of the leaft refrangible Rays is to the Refraction of the most refrangible ones as 27 to 28 very nearly, and that the difference of the Refractions of the leaft refrangible and most refrangible Rays is about the  $27\frac{1}{2}$ th part of the whole Refraction of the mean refrangible Rays.

Whence they that are skilled in Opticks will eafily understand, that the breadth of the least circular space into which Object-Glasses of Telescopes can collect all forts of Parallel Rays, is about the 27<sup>t</sup>th part of half the aperture of the Glass, or 55th part of the whole aperture; and that the Focus of the most refrangible Rays is nearer to the Object-Glass than the Focus of the least refrangible ones, by about the 27<sup>t</sup>th part of the distance between the Object-Glass and the Focus of the mean refrangible ones.

And if Rays of all forts, flowing from any one lucid point in the Axis of any convex Lens, be made by the Refraction of the Lens to converge to points not too remote from the Lens, the Focus of the most refrangible Rays shall be nearer to the Lens than the Focus of the least refrangible ones, by a distance which is to the  $27\frac{1}{2}$ th part of the distance of the Focus of the mean refrangible Rays from the Lens as the distance between that Focus and the lucid point

F

Now to examine whether the difference between the Refractions which the most refrangible and the least refrangible Rays flowing from the same point suffer in the Object-Glasses of Telescopes and such like Glasses, be so great as is here described, I contrived the following Experiment.

Exper. 16. The Lens which I used in the second and eighth Experiments, being placed fix Feet and an Inch dif-tant from any Object, collected the Species of that Object by the mean refrangible Rays at the distance of fix Feet and an Inch from the Lens on the other fide. And therefore by the foregoing Rule it ought to collect the Species of that Object by the least refrangible Rays at the distance of fix Feet and  $3\frac{2}{3}$  Inches from the Lens, and by the most refrangible ones at the distance of five Feet and 103 Inches from it : So that between the two Places where these least and most refrangible Rays collect the Species, there may be the distance of about 5<sup>1</sup>/<sub>3</sub> Inches. For by that Rule, as fix Feet and an Inch ( the distance of the Lens from the lucid Object ) is to twelve Feet and two Inches ( the distance of the lucid Object from the Focus of the mean refrangible Rays) that is, as one is to two, fo is the  $27\frac{1}{2}$ th part of fix Feet and an Inch (the diftance between the Lens and the same Focus ) to the distance between the Focus of the most refrangible Rays and the Focus of the least refrangible ones, which is therefore  $5\frac{17}{55}$  Inches, that is very nearly 5<sup>1</sup>/<sub>3</sub> Inches. Now to know whether this measure was true, I repeated the second and eighth Experiment of this Book with coloured Light, which was less compounded than that I there made use of : For I now separaced the hetero-

h eterogeneous Rays from one another by the Method I described in the 11th Experiment, so as to make a coloured Spectrum about twelve or fifteen times longer than broad. This Spectrum I cast on a printed book, and placing the above-mentioned Lens at the distance of fix Feet and an Inch from this Spectrum to collect the Species of the illuminated Letters at the same distance on the other side, I found that the Species of the Letters illuminated with Blue were nearer to the Lens than those illuminated with deep Red by about three Inches or three and a quarter : but the Species of the Letters illuminated with Indigo and Violet appeared to confused and indistinct, that I could not read them : Whereupon viewing the Prism, I found it was full of Veins running from one end of the Glass to the other; fo that the Refraction could not be regular. I took another Prism therefore which was free from Veins, and instead of the Letters I used two or three Parallel black Lines a little broader than the stroakes of the Letters, and casting the Colours upon these Lines in fuch manner that the Lines ran along the Colours from one end of the Spectum to the other, I found that the Focus where the Indigo, or confine of this colour and Violet cast the Species of the black Lines most distinctly, to be about 4 Inches or 4' near-er to the Lens than the Focus where the deepest Red cast from is ys low. the Species of the fame black Lines most distinctly. The violet was fo faint and dark, that I could not discern the Species of the Lines distinctly by that Colour; and therefore confidering that the Prism was made of a dark coloured Glass inclining to Green, I took another Pism of clear white Glass; but the Spectrum of Colours which this Prism made had long white Streams of faint Light fhooting out from both ends of the Colours, which made me conclude that something was amiss; and view-

ing

ing the Prism, I found two or three little Bubbles in the Glass which refracted the Light irregularly. Wherefore I covered that part of the Glass with black Paper, and letting the Light pass through another part of it which was free from such Bubles, the Spectrum of Colours became free from those irregular Streams of Light, and was now fuch as I desired. But still I found the Violet so dark and faint, that I could scarce see the Species of the Lines by the Violet, and not at all by the deepelt part of it, which was next the end of the Spectrum. I suspected therefore that this faint and dark Colour might be allayed by that fcattering Light which was refracted, and reflected irregularly partly by some very small Bubbles in the Glasses and partly by the inequalities of their Polifh: which Light, tho' it was but little, yet it being of a White Colour, might suffice to affect the Sense so strongly as to difturb the Phænomena of that weak and dark Colour the Violet, and therefore I tried, as in the 12th, 13th, 14th Experiments, whether the Light of this Colour did not confift of a sensible mixture of heterogeneous Rays, but found it did not. Nor did the Refractions cause any other sensible Colour than Violet to emerge out of this Light, as they would have done out of White Light, and by consequence out of this Violet Light had it been sensi-bly compounded with White Light. And therefore I concluded, that the reason why I could not see the Species of the Lines distinctly by this Colour, was only the darkness of this Colour and Thinnels of its Light, and its diftance from the Axis of the Lens; I divided therefore those Parallel Black Lines into equal Parts, by which I might readily know the distances of the Colours in the Spectrum from one another, and noted the distances of the Lens from the Foci of fuch Colours as cast the Species of the Lines

Lines diffinctly, and then confidered whether the difference of those distances bear such proportion to 5 3-Inches, the greatest difference of the distances which the Foci of the deepest Red and Violet ought to have from the Lens, as the distance of the observed Colours from one another in the Spectrum bear to the like distance of the deepest Red and Violet measured in the rectilinear fides of the Spectrum, that is, to the length of those fides or excess of the length of the Spectrum above its breadth. And my Observations were as follows.

When I observed and compared the deepest sensible Red, and the Colour in the confine of Green and Blue, which at that rectilinear fides of the Spectrum was diftant from it half the length of those fides, the Focus where the confine of Green and Blue cast the Species of the Lines distinctly on the Paper, was nearer to the Lens then the Focus where the Red caft those Lines dictinctly on it by about  $2\frac{1}{2}$  or 2<sup>3</sup> Inches. For sometimes the Measures were a little greater, sometimes a little less, but seldom varied from one another above  $\frac{1}{3}$  of an Inch. For it was very difficult to define the Places of the Foci, without some little Errors. Now if the Colours diftant half the length of the Image, (measured at its rectilinear fides ) give  $2\frac{1}{2}$  or  $2\frac{3}{4}$  difference of the distances of their Foci from the Lens, then the Colours distant the whole length ought to give 5 or 52 Inches difference of those distances.

But here it's to be noted, that I could not see the Red to the full End of the Spectrum, but only to the Center of the Semicircle which bounded that End, or a little farther; and therefore I compared this Red not with that Colour which was exactly in the middle of the Spectrum, or confine of Green and Blue, but with that which verged a little more to the Blue than to the Green : And as I reckoned

oned the whole length of the Colours not to be the whole length of the Spectrum, but the length of its rectilinear fides, fo completing the Semicirlar Ends into Circles, when either of the obferved Colours fell within those Circles, I measured the distance of that Colour from the End of the Spectrum, and subducting half the distance from the meafured distance of the Colours, I took the remainder for their corrected distance; and in these Observations set down this corrected distance for the difference of their diftances from the Lens. For asthe length of the rectilinear fides of the Spectrum would be the whole length of all the Colours, were the Circles of which (as we shewed) that Spectrum consists contracted and reduced to Physical Points, fo in that Case this corrected distance would be the real distance of the observed Colours.

When therefore I further observed the deepest sensible Red, and that Blue whose corrected distance from it was  $\frac{7}{12}$  parts of the length of the rectilinear fides of the Spectrum, the difference of the distances of their Foci from the Lens was about 3<sup>-1</sup>/<sub>4</sub> Inches, and as 7 to 12 fo is 3<sup>-1</sup>/<sub>4</sub> to 5<sup>-4</sup>/<sub>5</sub>.

When I observed the deepest sensible Red, and that Indigo whose corrected distance was  $\frac{8}{12}$  or  $\frac{2}{3}$  of the length of the rectilinear sides of the Spectrum, the difference of the distances of their Foci from the Lens, was about 3  $\frac{2}{3}$  Inches, and as 2 to 3 so is 3  $\frac{2}{3}$ -to  $5\frac{1}{2}$ .

When I observed the deepest sensible Red, and that deep Indigo whose corrected distance from one another was  $\frac{9}{12}$  or  $\frac{3}{4}$  of the length of the rectilinear sides of the Spectum, the difference of the distances of their Foci from the Lens was about 4 Inches; and as 3 to 4 so 5  $\frac{1}{3}$ .

When I observed the deepest sensible Red, and that part of the Violet next the Indigo whose corrected distance from the Red was  $\frac{10}{12}$  or  $\frac{5}{6}$  of the length of the rectilinear fides of

the

the Spectrum, the difference of the distances of their Foci. from the Lens was about  $4\frac{1}{2}$  Inches; and as 5 to 6, fo is  $4\frac{1}{2}$  to  $5\frac{2}{5}$ . For fometimes when the Lens was advantagioufly placed, fo that its Axis respected the Blue, and all things else were well ordered, and the Sun shone clear, and I held my Eye very near to the Paper on which the Lens cast the Species of the Lines, I could see pretty distinctly the Species of those Lines by that part of the Violet which was next the Indigo; and sometimes I could see them by above half the Violet. For in making these Experiments I had observed, that the Species of those Colours only appeared distinct which were in or near the Axis of the Lens : So that if the Blue or Indigo were in the Axis, I could fee their Species diffinctly; and then the Red appeared much less distinct than before. Wherefore I contrived to make the Spectrum of Colours shorter than before, so that both its Ends might be nearer to the Axis of the Lens. And now its length was about  $2\frac{1}{2}$  Inches and breadth about  $\frac{1}{5}$  or of an Inch. Also instead of the black Lines on which the Spectrum was caft, I made one black Line broader than those, that I might see its Species more easily; and this Line I divided by short cross Lines into equal Parts, for measuring the distances of the observed Colours. And now I could fometimes fee the Species of this Line with its divifions almost as far as the Centers of the Semicircular Violet End of the Spectrum, and made these further Observations.

When I observed the deepest sensible Red, and that part of the Violet whose corrected distance from it was about  $\frac{8}{9}$  Parts of the rectilinear sides of the Spectrum the difference of the distances of the Foci of those Colours from the Lens, was one time  $4\frac{2}{3}$ , another time  $4\frac{3}{4}$ , another time  $4\frac{7}{81}$  Inches, and as 8 to 9, so are  $4\frac{2}{3}$ ,  $4\frac{3}{4}$ ,  $4\frac{7}{8}$ , to  $5\frac{1}{4}$ ,  $5\frac{11}{32}5\frac{31}{64}$  respectively. I 2 When I observed the deepest sensible Red, and deepest sensible Violet, (the corrected distance of which Colours when all things were ordered to the best advantage, and the Sun shone very clear, was about  $\frac{11}{12}$  or  $\frac{15}{16}$  parts of the length of the rectilinear fides of the coloured Spectrum, ) I found the difference of the distances of their Foci from the Lens sometimes  $4\frac{3}{4}$  fometimes  $5\frac{1}{4}$ , and for the most part 5 Inches or thereabouts : and as 11 to 12 or 15 to 16, so is five Inches to  $5\frac{1}{2}$  or  $5\frac{1}{2}$  Inches.

And by this progression of Experiments I satisfied my felf, that had the light at the very Ends of the Spectrum been strong enough to make the Species of the black Lines appear plainly on the Paper, the Focus of the deepest Violet would have been found nearer to the Lens, than the Focus of the deepest Red, by about 5<sup>1</sup>/<sub>5</sub> Inches at least. And this is a further Evidence, that the Sines of Incidence and Refraction of the several forts of Rays, hold the same proportion to one another in the state of Refractions which they do in the greatest.

My progrefs in making this nice and troublefome Experiment I have fet down more at large, that they that shall try it after me may be aware of the Circumspection requifite to make it fucceed well. And if they cannot make it fucceed fo well as I did, they may notwithstanding collect by the Proportion of the distance of the Colours in the Spectrum, to the difference of the distances of their Foci from the Lens, what would be the fucces in the more distant Colours by a better Trial. And yet if they use a broader Lens than I did, and fix it to a long streight Staff by means of which it may be readily and truly directed to the Colour whose Focus is defired, I question not but the Experiment will succeed better with them than it did with me. For I directed the Axis as nearly as I could to the middle middle of the Colours, and then the faint Ends of the Spectrum being remote from the Axis, cast their Species less distinctly on the Paper than they would have done had the Axis been successively directed to them.

-He-

Now by what has been faid its certain, that the Rays which differ in refrangibility do not converge to the fame Focus, but if they flow from a lucid point, as far from the Lens on one fide as their Foci are one the other, the Focus of the most refrangible Rays shall be nearer to the Lens than that of the least refrangible, by above the fourteenth part of the whole distance: and if they flow from a lucid point, so very remote from the Lens that before their Incidence they may be accounted Parallel, the Focus of the most refrangible Rays shall be nearer to the Lens than the Focus of the least refrangible, by about the 27th or 28th part of their whole distance from it. And the Diameter of the Circle in the middle space between those two Foci which they illuminate when they fall there on any Plane, perpendicular to the Axis (which Circle is the least into which they can all be gathered) is about the 55th part of the Diameter of the aperture of the Glass. So that 'tis a wonder that Telescopes represent Objects so distinct as they do. But were all the Rays of Light equally refrangible, the Error arising only from the sphericalness of the Figures of Glasses would be many hundred times less. For if the Object-Glass of a Telescope be Plano-convex, and the Plane side be turned towards the Object, and the Diameter of the Sphere whereof this Glass is a segment, be called D, and the Semidiameter of the aperture of the Glass be called S, and the Sine of Incidence out of Glass into Air, be to the Sine of Refraction as I to R: the Rays which come Parallel to the Axis of the Glafs, shall in the Place where the Image of the Object is most distinctly made, be scattered all over a little Circle

70

Circle whole Diameter is  $\frac{R}{I} \times \frac{S cub.}{D quad.}$  very nearly, as I gather by computing the Errors of the Rays by the method of infinite Series, and rejecting the Terms whose quantitities are inconfiderable. As for instance, if the Sine of Incidence I, be to the Sine of Refraction R, as 20 to 31, and if D the Diameter of the Sphere to which the Convex side of the Glass is ground, be 100 Feet or 1200 Inches, and S the Semidiameter of the aperture be two Inches, the Diameter of the little Circle ( that is  $\frac{R \times S \ cub}{I \times D \ quad}$  ) will be

 $\frac{31 \times 8}{20 \times 1200 \times 1200}$  (or  $\frac{31}{3600000}$ ) parts of an Inch. But the Diameter of the little Circle through which these Rays are scattered by unequal refrangibility, will be about the 55th part of the aperture of the Object-Glass which here is four Inches. And therefore the Error arising from the spherical Figure of the Glass, is to the Error arising from the diffe-rent Refrangibility of the Rays, as  $\frac{31}{3600000}$  to  $\frac{4}{55}$  that is as 1 to 8151 : and therefore being in Comparison so very little, deserves not to be considered.

But you will fay, if the Errors caused by the different refrangibility be so very great, how comes it to pass that Objects appear through Telescopes so distinct as they do ? I answer, 'tis because the erring Rays are not scattered uniformly over all that circular space, but collected infinitely more densely in the Center than in any other part of the Circle, and in the way from the Center to the Circumference grow continually rarer and rarer, fo as at the Circumference to become infinitely rare; and by reason of their rarity are not strong enough to be visible, unless in the Center and ve-Fig. 27. ry near it. Let ADE represent one of those Circles defcribed with the Center C and Semidiameter AC, and let BFG be a smaller Circle concentric to the former, cutting with

with its Circumference the Diameter AC in B, and befect AC in N, and by my reckoning the denfity of the Light in any place B will be to its denfity in N, as AB to BC; and the whole Light within the leffer Circle BFG, will be to the whole Light within the greater AED, as the Excess of the Square of AC above the Square of AB, is to the Square of AC. As if BC be the fifth part of AC, the Light will be four times denfer in B than in N, and the whole Light within the lefs Circle, will be to the whole Light within the greater, as nine to twenty five. Whence it's evident that the Light within the lefs Circle, mult ftrike the fenfe much more ftrongly, than that faint and dilated light round about between it and the Circumference of the greater.

But its further to be noted, that the most luminous of the prismatick Colours are the Yellow and Orange. These a mosium between yolds affect the Senses more strongly than all the rest together, and and red, y best ofto alour next to these in strength are the Red and Green. The Blue compared with these is a faint and dark Colour, and the Indigo and Violet are much darker and fainter, so that these compared with the stronger Colours are little to be regarded. The Images of Objects are therefore to be placed, not in the Focus of the mean refrangible Rays which are in the confine of Green and Blue, but in the Focus of those Rays which are in the middle of the Orange and Yellow; there where the Colour is most luminous and fulgent, that is in the brightest Yellow, that Yellow which inclines more to Orange than to Green. And by the Refraction of these Rays ( whole Sines of Incidence and Refraction in Glass are as 17 and 11) the Refraction of Glass and Crystal for optical uses is to be measured. Let us therefore place the Image of the Object in the Focus of these Rays, and all the Yellow and Orange will fall within a Circle, whofe Diameter is about the 250th part of the Diameter of the aperture

ture of the Glass. And if you add the brighter half of the Red, (that half which is next the Orange, and the brighter half of the Green, (that half which is next the Yellow,) about three fifth parts of the Light of these two Colours will fall within the same Circle, and two fifth parts will fall without it round about; and that which falls without will be spread through almost as much more space as that which falls within, and so in the gross be almost three times rarer. Of the other half of the Red and Green, (that is of the deep dark Red and Willow Green) about one quarter will fall within this Circle, and three quarters without, and that which falls without will be spread through about four or five times more space than that which falls within; and so in the gross be rarer, and if compared with the whole Light within it, will be about 25 times rarer than all that taken in the gross; or rather more than 30 or 40 times rarer, because the deep red in the end of the Spectrum of Colours made by a Prism is very thin and rare, and the Willow Green is fomething rarer than the Orange and Yellow. The Light of these Colours therefore being so very much rarer than that within the Circle, will scarce affect the Sense especially fince the deep Red and Willow Green of this Light, are much darker Colours then the rest. And for the same reason the Blue and Violet being much darker Colours than thefe, and much more rarified, may be neglected. For the dense and bright Light of the Circle, will obscure the rare and weak Light of these dark Colours round about it, and render them almost insensible. The sensible Image of a lucid point is therefore scarce broader than a Circle whole Diameter is the 250th part of the diameter of the aperture of the Object Glass of a good Telescope, or not much broader, if you except a faint and dark misty light round about it, which a Spectator will scarce regard. And therefore in a Telescope whole

whose aperture is four Inches, and length an hundred Feet, it exceeds not 2"45", or 3". And in a Telescope whose. aperture is two Inches, and length 20 or 30 Feet, it may be 5" or 6" and scarce above. And this Answers well to Experience : For some Aftronomers have found the Diameters of the fixt Stars, in Telescopes of between twenty and fixty Feet in length, to be about 4" or 5" or at most 6" in Diameter. But if the Eye-Glass be tincted faintly with the moke of a Lamp or Torch, to obscure the Light of the Star, the fainter Light in the circumference of the Star ceases to be visible, and the Star (if the Glass be sufficiently foiled with fmoke) appears something morelike a Mathematical Point. And for the fame reason, the enormous part of the Light in the Circumference of every lucid Point ought to be less discernable in shorter Telescopes than in

ought to be leis different refrangibility of the greatest Errors arifing from the greatest Errors arifing from the greatest fensible Errors ari- a Citte und be but 31/3600000
of the Glafs, would be to the greatest fensible Errors ari- a Citte und be but and be but and for the greatest fensible Errors ari- a Citte und be but 31/3600000 to  $\frac{4}{250}$  at most, that is only as 1 to 1826. And this fufficiently shews that it is not the spherical Figures of Glasses but the different refrangibility of the Rays which hinders the perfection of Telescopes.

There is another Argument by which it may appear that the different refrangibility of Rays, is the true Caufe of the imperfection of Telescopes. For the Errors of the Rays arifing from the spherical Figures of Object-Glasses, are as the the Cubes of the apertures of the Object-Glaffes; and thence to make Telescopes of various lengths, magnify with equal distinctness, the apertures of the Object-Glaffes, and the Charges or magnifying Powers, ought to be as the Cubes of the square Roots of their lengths; which doth not answer to Experience. But the errors of the Rays arising from the different refrangibility, are as the apertures of the Object-Glaffes, and thence to make Telescopes of various lengths, magnify with equal distinctness, their apertures and charges ought to be as the square Roots of their lengths; and this answers to experience as is well known. For instance, a Telescope of 64 Feet in length, with an aperture of  $2\frac{2}{3}$  Inches, magnifies about 120 times, with as much diftinctness as one of a Foot in length, with  $\frac{1}{3}$  of an Inch aperture, magnifies 15 times.

Now were it not for this different refrangibility of Rays, Telescopes might be brought to a greater Perfection than we have yet described, by composing the Object-Glass of two Glasses with Water between them. Let ADFC repre-Fig. 28. sent the Object-Glass composed of two Glasses ABED and and BEFC, alike convex on the outfides AGD and CHF, and alike concave on the infides BME, BNE, with Water in the concavity BMEN. Let the Sine of Incidence out of Glass into Air be as I to R and out of Water into Air as K to R, and by consequence out of Glass into Water, as I to K: and let the Diameter of the Sphere to which the convex fides AGD and CHF are ground be D, and the Diameter of the Sphere to which the concave sides BME and BNE are ground be to D, as the Cube Root of KK-KI to the Cube Root of RK-RI: and the Refractions on the concave fides of the Glasses, will very much correct the Errors: of the Refractions on the convex fides, so far as they arife from the sphericalness of the Figure. And by this means might

[74]

might Telescopes be brought to sufficient perfection, were it not for the different refrangibility of several fors of Rays. But by reason of this different refrangibility, I do not yet see any other means of improving Telescopes by Refractions alone than that of increasing their lengths, for which end the late  $\chi$ contrivance of *Hugenius* seems well accommodated. For very long Tubes are cumbersome, and scarce to be readily managed, and by reason of their length are very apt to bend, and shake by bending so as to cause a continual trembling in the Objects, whereby it becomes difficult to see them diffinctly : whereas by his contrivance the Glasses are readily manageable, and the Object-Glass being fixt upon a strong upright Pole becomes more steddy.

Seeing therefore the improvement of Telescopes of given lengths by Refractions is desperate; I contrived heretofore a Perspective by reflexion, using instead of an Object Glass a concave Metal. The diameter of the Sphere to which the Metal was ground concave was about 25 English Inches, and by confequence the length of the Inftrument about six Inches and a quarter. The Eye-Glass was plano-convex, and the Diameter of the Sphere to which the convex side was ground was about ; of an Inch, or a little lefs, and by confequence it magnified between 30 and 40 times. By another way of measuring I found that it magnified about 35 times. The Concave Metal bore an aperture of an Inch and a third part; but the aperture was limited not by an opake Circle, covering the Limb of the Metal round about, but by an opake circle placed between the Eye-Glass and the Eye, and perforated in the middle with a little round hole for the Rays to pass through to the Eye. For this Circle by being placed here, stopt much of the erroneous Light, which otherwise would have disturbed the Vision. By comparing it with a pretty good Perspective of four Feet in length, K 2.

length, made with a concave Eye-Glass, I could read at a greater distance with my own Instrument than with the Glass. Yet Objects appeared much darker in it than in the Glafs, and that partly becaufe more Light was lost by re-flexion in the Metal, then by refraction in the Glafs, and partly because my Instrument was overcharged. Had it magnified but 30 or 25 times it would have made the Object appear more brisk and pleafant. Two of these Imade about 16 Years ago, and have one of them still by me by which I can prove the truth of what I write. Yet it is not so good as at the first. For the concave has been divers times tarnished and cleared again, by rubbing it with very foft Leather. When I made these, an Artist in London undertook to imitate it; but using another way of polishing them than I did, he fell much short of what I had attained to, as l'afterwards understood by discoursing the under-Work-man he had imployed. The Polish I used was on thismanner. I had two round Copper Plates each fix Inches in Diameter, the one convex the other concave, ground very true to one another. On the convex I ground the Object-Metal or concave which was to be polifh'd, till it had taken the Figure of the convex and was ready for a Polish. Then I pitched over the convex very thinly, by dropping melted pitch upon it and warming it to keep the pitch soft, whilst I ground it with the concave Copper wetted to make it spread evenly all over the convex. Thus by working it well I made it as thin as a Groat, and after the convex was cold I ground it again to give it as true a Figure as I could. Then I took Putty which I had made very fine by washing it from all its groffer Particles, and laying a lit-tle of this upon the pitch, I ground it upon the Pitch with the concave Copper till it had done making a noise; and then upon the Pitch I ground the Object-Metal with a brisk Motion

Motion, for about two or three Minutes of time, leaning hard upon it. Then I put fresh Putty upon the Pitch and ground it again till it had done making a noise, and afterwards ground the Object Metal upon it as before. And this Work I repeated till the Metal was polifhed, grinding it the last time with all my strength for a good while toge-ther, and frequently breathing upon the Pitch to keep it moist without laying on any more fresh Putty. The Object-Metal was two Inches broad and about one third part of an Inch thick, to keep it from bending. I had two of these Metals, and when I had polished them both I tried which was best, and ground the other again to see if I could make it better than that which I kept. And thus by many Trials I learnt the way of polishing, till I made those two reflecting Pespectives I spake of above. For this Art of polifhing will be better learnt by repeated Practice than by my description. Before I ground the Object Metal on the Pitch, I always ground the Putty on it with the concave Copper till it had done making a noise, because if the Particles of the Putty were not by this means made to stick fast in the Pitch, they would by rolling up and down grate and fret the Object Metal and fill it full of little holes.

But becaufe Metal is more difficult to polifh than Glafs and is afterwards very apt to be fpoiled by tarnifhing, and reflects not fo much Light as Glafs quick-filvered over does: I would propound to use instead of the Metal, a Glafs ground concave on the forefide, and as much convex on the backfide, and quick-filvered over on the convex fide. The Glass must be every where of the fame thickness exactly. Otherwise it will make Objects look coloured and indistinct. By fuch a Glafs I tried about five or fix Years ago to make a reflecting Telescope of four Feet in length to magnify about 150 times, and I stissied my felf that there wants nothing

## [78]

thing but a good Artist to bring the design to Perfection. For the Glass being wrought by one of our London Artists after such a manner as they grind Glasses for Telescopes, tho it seemed as well wrought as the Object Glasses use to be, yet when it was quick-filvered, the reflexion discovered innumerable Inequalities all over the Glass. And by reason of these Inequalities, Objects appeared indistinct in this Instrument. For the Errors of reflected Rays caused by any Inequality of the Glass, are about six times greater than the Errors of refracted Rays caused by the like Inequalities. Yet by this Experiment I fatisfied my felf that the reflexion on the concave side of the Glass, which I feared would disturb the vision, did no sensible prejudice to it, and by consequence that nothing is wanting to perfect these Telescopes, but good Workmen who can grind and polish Glasses truly spherical. An Object-Glass of a fourteen Foot Telescope, made by one of our London Artificers, I once mended confiderably, by grinding it on Pitch with Putty, and leaning ve-ry eafily on it in the grinding, left the Putty fhould fcratch it. Whether this way may not do well enough for polifhing these reflecting Glasses, I have not yet tried. But he that shall try either this or any other way of polishing which he may think better, may do well to make his Glaffes ready for polishing by grinding them without that violence, wherewith our London Workmen press their Glasses in grinding. For by fuch violent pressure, Glasses are apt to bend a little in the grinding, and such bending will certainly spoil their Figure. To recommend therefore the confideration of these reflecting Glasses, to such Artists as are curious in figuring Glasses, I shall describe this Optical Instrument in the following Proposition.

## [79]

### PROP. VII. Prob. II.

#### To shorten Telescopes.

E T ABDC represent a Glass spherically concave on Fig. 29. the foreside AB, and as much convex on the backfide CD, so that it be every where of an equal thickness. Let it not be thicker on one side than on the other, lest it make Objects appear coloured and indiffinct, and let it be very truly wrought and quick-filvered over on the backfide; and fet in the Tube VXYZ which must be very black within. Let EFG represent a Prism of Glass or Crystal placed near the other end of the Tube, in the middle of it, by means of a handle of Brass or Iron FGK, to the end of which made flat it is cemented. Let this Prism be rectangular at E, and let the other two Angles at F and G be accurately equal to each other, and by confequence equal to half right ones, and let the plane fides FE and GE be square, and by consequence the third fideFG a rectangular parallelogram, whofe length is to its breath in a subduplicate proportion of two to one. Let it be so placed in the Tube, that the Axis of the Speculum may pass through the middle of the square fide EF perpendicularly, and by consequence through the middle of the fide F G at an Angle of 45 degrees, and let the fide EF be turned towards the Speculum, and the distance of this Prism from the Speculum be such that the Rays of the light PQ, RS, &c. which are incident upon the Speculum in Lines Parallel to the Axis thereof, may enter the Prism at the fide EF, and be reflected by the fide F G, and thence go out of it through the fide GE, to the point T which must be the common Focus of the Speculum ABDC, and of a Plano-convex Eye-Glass H, through which those Rays must pass to the Eye. And let the Rays at their coming QUE

out of the Glass pass through a small round hole, or aperture made in a little Plate of Lead, Brass, or Silver, wherewith the Glass is to be covered, which hole must be no bigger than is necessary for light enough to pass through. For so it will render the Object distinct, the Plate in which 'tis made intercepting all the erroneous part of the Light which comes from the Verges of the Speculum AB. Such an Instrument well made if it be 6 Foot long, (reckoning the length from the Speculum to the Prism, and thence to the Focus T) will bear an aperture of 6 Inches at the Speculum, and magnify between two and three hundred times. But the hole H here limits the aperture with more advantage, then if the aperture was placed at the Speculum. If the Instrument be made longer or shorter, the aperture must be in proportion as the Cube of the square Root of the length, and the magnifying as the aperture. But its convenient that the Speculum be an Inch or two broader than the aperture at the least, and that the Glass of the Speculum be thick, that it bend not in the working. The Prism EFG must be no bigger than is necessary, and its back fide FG must not be quick-filvered over. For without quick-filver it will reflect all the Light incident on it from the Speculum. In this Instrument the Object will be inverted, but may be erected by making the square fides EF and EG of the Prism EFG not plane but spherically convex, that the Rays may cross as well before they come at it as afterwards between it and the Eye-Glass. If it be defired that the Instrument bear a larger aperture, that may be also done by composing the Speculum of two Glasses with Water between poing the Speculum of two Galles with Water between them. But if all this at last, which indeed any one may saynost and propose 1899-915 could be made according to art; although there are certain limit, bekind which belescops can by no moans be made. For if dor norphulu hendles as we may see, from & hembling mohon of y thedows of Proscheles cast from high tow-ers, or from & sparkling of & field start. But thous here is made accord the there through Polescops, which have very wide (as & might be made accord the through the sile rent parts of this large of the side of a spiralely which have ing to this fashion would not sparkle. For & Parkeles of light, which have through the silerent parts of this large of the side of the sign of through the silerent parts of this large of y the ends and of the mode accord the is the porceidd. tohich not sparkle for a start of the site of a site of the side of this fashion would not sparkle for a start of the site of the ing to this fashion would not sparkle for a start of the site of the site of porceidd. the large of the ends and y temptime upon the rand for through the site of the starts of the large of the start of the site of the site of the search of y duicknow and confused on the motion they cannot in y least be porceidd. tohich north thus enlight and made on a large lucid point where the they than seem larger than they would are, and that without any thomas in the set of the at can be porceidd. Tohich grants to a, to take away y confusionely and and prove fighter c larger, but by no motion is a calm air, such as yorkars might be had epon y the of y highest mountuins.

[80]

Book J. Plate I. Part L.



















Book I. Plate III. Part I.







•

·

.

.\*

BOOK, I. Plate, IV. Part, I.









# [81] THE FIRST BOOK OF OPTICKS. PART II.

#### PROP. I. THEOR. I.

The Phanomena of Colours in refracted or reflected Light are not caused by new modifications of the Light variously impress, according to the various terminations of the Light and Shadow.

The Proof by Experiments.

#### EXPER. I.

FOR if the Sun shine into a very dark Chamber Fig. 1. through an oblong Hole F, whose breadth is the fixth or eighth part of an Inch, or something less; and his Beam FH do afterwards pass first through a very large Prism ABC, distant about 20 Feet from the L Hole, Hole, and parallel to it, and then (with its white part) through an oblong Hole H, whole breadth is about the fortieth or fixtieth part of an Inch, and which is made in a black opake Body GI, and placed at the diftance of two or three Feet from the Prilm, in a parallel fituation both to the Prism and to the former Hole, and if this white Light thus transmitted through the Hole H, fall afterwards upon a white Paper pt, placed after that Hole H, at the distance of three or four Feet from it, and there paint the usual Colours of the Prism, suppose red at t, yellow at s, green at r, blue at q, and violet at p; you may with an iron Wire, or any fuch like flender opake Body, whofe breadth is about the tenth part of an Inch, by intercepting the rays at k, l, m, n or o, take away any one of the Colours at t, s, r, q or p, whilft the other Colours remain up-on the Paper as before; or with an obftacle fomething bigger you may take away any two, or three, or four Co-lours together, the reft remaining: So that any one of the Colours as well as violet may become outmost in the confine of the shadow towards p, and any one of them as well as red may become outmost in the confine of the shadow towards t, and any one of them may also border upon the shadow made within the Colours by the obstacle R intercepting some intermediate part of the Light; and, lastly, any one of them by being left alone may border upon the fhadow on either hand ... All the Colours have themselves indifferently to any. confines of fhadow, and therefore the differences of theic Colours from one another, do not arise from the different confines of shadow, whereby Light is variously modified as has hitherto been the Opinion of Philofophers.

phers. In trying these things 'tis to be observed, that by how much the Holes F and H are narrower, and the intervals between them, and the Prisin greater, and the Chamber darker, by so much the better doth the Experiment succeed; provided the Light be not so far diminissed, but that the Colours at pt be sufficiently visible. To procure a Prism of solid Glass large enough for this Experiment will be difficult, and therefore a prismatick Vessel must be made of polished Glass-plates cemented together, and filled with Water. or clear oyl.

#### EXPER. II.

The Sun's Light let into a dark Chamber through Fig. 2. the round Hole F, half an Inch wide, passed first through the Prism ABC placed at the Hole, and then through a Lens PT fomething more than four Inches broad, and about eight Feet distant from the Prism, and thence converged to O the Focus of the Lens diftant from it about three Feet, and there fell upon a white Paper DE. If that Paper was perpendicular to that Light incident upon it, as 'tis represented in the posture DE, all the Colours upon it at O appeared white. But if the Paper being turned about an Axis parallel to the Prism, became very much inclined to the Light as 'tis reprefented in the politions de and e; the fame Light in the one cafe appeared yellow and red, in the other blue. Here one and the fame part of the Light in one and the same place, according to the various inclinations of the Paper, appeared in one cafe white, in another yellow or red, in a third blue, whilft the confine of Light and Shadow, L 2

Shadow, and the refractions of the Prism in all these cafes remained the same.

#### EXPER. III.

Fig. 3.

Such another Experiment may be more eafily tried as follows. Let a broad beam of the Sun's Light coming into a dark Chamber through a Hole in the Window shut be refracted by a large Prism ABC, whose re-fracting Angle C is more than 60 degrees, and so soon as it comes out of the Prism let it fall upon the white Paper DE glewed upon a stiff plane, and this Light, when the Paper is perpendicular to it, as 'tis represented in DE, will appear perfectly white upon the Paper. but when the Paper is very much inclined to it in fuch a manner as to keep always parallel to the Axis of the Prism, the whiteness of the whole Light upon the Paper will according to the inclination of the Paper this way, or that way, change either into yellow and red, as in the posture de, or into blue and violet, as in the posture of. And if the Light before it fall upon the Paper be twice refracted the fame way by two parallel Prifms, these Colours will become the more confpicuous. Here all the middle parts of the broad beam of white Light which fell upon the Paper, did without any confine of shadow to modify it, become coloured all over with one uniform Colour, the Colour being always the fame in the middle of the Paper as at the edges, and this Colour changed according the various obliquity of the reflecting Paper, without any change in the refractions or shadow, or in the Light which fell upon the Paper. And therefore these Colours are to to be derived from fome other cause than the new modifications of Light by refractions and shadows.

If it be asked; What then is their caufe? I anfwer,  $\bigstar$ That the Paper in the pofture de, being more oblique to the more refrangible rays than to the lefs refrangible ones, is more ftrongly illuminated by the latter than by the former, and therefore the lefs refrangible rays are predominant in the reflected Light. And wherever they are predominant in any Light they tinge it with red or yellow, as may in fome measure appear by the first Proposition of the first Book, and will more fully appear hereafter. And the contrary happens in the posture of the Paper  $d_{\pm}$ , the more refrangible rays being then predominant which always tinge Light with blues and violets.

#### EXPER. IV.

The Colours of Bubbles with which Children play are various, and change their fituation varioufly, without any refpect to any confine of fhadow. If fuch a Bubble be covered with a concave Glafs, to keep it from being agitated by any wind or motion of the Air, the Colours will flowly and regularly change their fituation, even whilft the Eye, and the Bubble, and all Bodies which emit any Light, or caft any fhadow, remain unmoved. And therefore their Colours arife from fome regular caufe which depends not on any confine of fhadow. What this caufe is will be fhewed in the next Book.

To these Experiments may be added the tenth Ex-periment of the first Book, where the Sun's Light in a dark Room being trajected through the parallel superficies of two Prisms tied together in the form of a Parallelopide, became totally of one uniform yellow or red Colour, at its emerging out of the Prisms. Here, in the production of these Colours, the confine of shadow can have nothing to do. For the Light changes from white to yellow, orange and red fucceffively, without any alteration of the confine of shadow: And at both edges of the emerging Light where the contrary confines of shadow ought to produce different effects, the Colour is one and the fame, whether it be white, yellow, orange or red : And in the middle of the emerging Light, where there is no confine of fhadow at all, the Colour is the very fame as at the edges, the whole Light at its very first emergence being of one uniform Colour, whether white, yellow, orange or red, and going on thence perpetually without any change of Colour, fuch as the confine of shadow is vulgarly supposed to work in refracted Light after its emergence. Neither can thefe Colours arife from any new modifications of the Light by refractions, because they change fuccessively from white to yellow, orange and red, while the refractions remain the fame, and also because the refractions are made contrary ways by parallel fuperficies which deftroy one anothers effects. They arise not therefore from any modifications of Light made by refractions and fhadows, but have fome other caufe. What that caufe is we shewed above in this tenth Experiment, and need not here repeat it.

There
There is yet another material circumstance of this Experiment. For this emerging Light being by a third Fig. 22.. Prism HIK refracted towards the Paper PT, and there Part 1. painting the usual Colours of the Prism, red, yellow, green, blue, violet : If these Colours arose from the refractions of that Prism modifying the Light, they would not be in the Light before its incidence on that Prism. And yet in that Experiment we found that when by turning the two first Prisms about their common Axis all the Colours were made to vanish but the red; the Light which makes that red being left alone, appeared of the very fame red Colour before its incidence on the third Prism. And in general we find by other Experiments that when the rays which differ in refrangibility are separated from one another, and any one fort of them is confidered apart, the Colour of the Light which they compose cannot be changed by any refraction or reflexion whatever, as it ought to be were Colours nothing else than modifications of Light caused by refractions, and reflexions, and shadows. This unchangeablenefs of Colour I am now to defcribe in the following Proposition.

## PROP. II. THEOR. II.

All homogeneal Light has its proper Colour anfwering to its degree of refrangibility, and that Colour cannot be changed by reflexions and refractions.

In the Experiments of the 4th Proposition of the first Book, when I had separated the heterogeneous rays from one another, the Spectrum pt formed by the separated rated rays, did in the progrefs from its end p, on which the most refrangible rays fell, unto its other end t, on which the least refrangible rays fell, appear tinged with this Series of Colours, violet, indico, blue, green, yellow, orange, red, together with all their intermediate degrees in a continual fucceffion perpetually varying: So that there appeared as many degrees of Colours, as there were forts of rays differing in refrangibility. for though all colours may be comprehended under soon principal onsy, yel of these these are unnumerable pagrees. E X P E R. V.

Now that these Colours could not be changed by refraction, I knew by refracting with a Prism sometimes one very little part of this Light, fometimes another very little part, as is defcribed in the 12th Experiment of the first Book. For by this refraction the Colour of the Light was never changed in the leaft. If any part of the red Light was refracted, it remained totally of the fame red Colour as before. No orange, no yellow, no green, or blue, no other new Colour was pro-duced by that refraction. Neither did the Colour any ways change by repeated refractions, but continued always the same red entirely as at first. The like conftancy and immutability I found also in the blue, green, and other Colours. So also if I looked through a Prism upon any body illuminated with any part of this homo-geneal Light, as in the 14th Experiment of the first Book is described; I could not perceive any new Co-lour generated this way. All Bodies illuminated with compound Light appear through Prisms confused (as was faid above) and tinged with various new Colours, but those illuminated with homogeneal Light appeared through

through Pritins neither lefs diftinct, nor otherwife coloured, than when viewed with the naked Eyes. Their Colours were not in the leaft changed by the refraction of the interpofed Prifm. I fpeak here of a fenfible change of Colour : For the Light which I here call homogeneal, being not abfolutely homogeneal, there ought to arife fome little change of Colour from its heterogeneity. But if that heterogeneity was fo little as it might be made, by the faid Experiments of the fourth Propofition, that change was not fenfible, and therefore, in Experiments where fenfe is judge, ought to be accounted none at all.

#### EXPER. VI.

And as thefe Colours were not changeable by refractions, io neither were they by reflexions. For all white, grey, red, yellow, green, blue, violet Bodies, as Paper, Afhes, red Lead, Orpiment, Indico, Bife, Gold, Silver, Copper, Grafs, blue Flowers, Violets, Bubbles of Water tinged with various Colours, Peacock's Feathers, the tincture of Lignum Nephriticum, and fuch like, in red homogeneal Light appeared totally red, in blue Light totally blue, in green Light totally green, and fo of other Colours. In the homogeneal Light of of any Colour they all appeared totally of that fame Colour, with this only difference, that fome of them reflected that Light more ftrongly, others more faintly. I never yet found any Body which by reflecting homogeneal Light could fenfibly change its Colour.

From

×

From all which it is manifeft, that if the Sun's Light confifted of but one fort of rays, there would be but one Colour in the whole World, nor would it be poffible to produce any new Colour by reflexions and refractions, and by confequence that the variety of Colours depends upon the composition of Light.

## DEFINITION.

malfremos

The homogeneal light and rays which appear red, or rather make Objects appear so, I call rubrific or red-making; those which make Objects appear yellow, green, blue and violet, I call yellow-making, green-making, blue-making, violet-making, and fo of the reft. And if at any time I fpeak of light and rays as coloured or endued with Colours, I would be underftood to speak not philofophically and properly, but groily, and accor-ding to fuch conceptions as vulgar People in feeing all these Experiments would be apt to frame. For the rays to fpeak properly are not coloured. In them there is nothing else than a certain power and disposition to stir up a sensation of this or that Colour. For as found in a Bell or mufical String or other founding Body, is nothing but a trembling Motion, and in the Air nothing but that. Motion propagated from the Object, and in the Senforium 'tis a sense of that Motion under the form of found ; fo Colours in the Object are nothing but a disposition to reflect this or that fort of rays more copioully than the reft; in the rays they are nothing but their difpolitions to propagate

gate this or that Motion into the Senforium, and in the Senforium they are fenfations of those Motions under the forms of Colours.  $\downarrow$ 

## PROP. III. PROB. I.

To define the refrangibility of the feveral forts of homogeneal Light answering to the feveral Colours.

For determining this Problem I made the following Experiment.

#### EXPER. VII.

When I had caufed the rectilinear line fides AF, GM, Fig. 4. of the Spectrum of Colours made by the Prifm to be diftinctly defined, as in the fifth Experiment of the first Book is described, there were found in it all the homogeneal Colours in the fame order and fituation one among another as in the Spectrum of fimple Light, described in the fourth Experiment of that Book. For the Circles of which the Spectrum of compound Light PT is composed, and which in the middle parts of the Spectrum interfere and are intermixt with one another, are not intermixt in their outmost parts where they touch those rectilinear fides AF and GM. And therefore in those rectilinear fides when diffinctly defined, there is no new Colour generated by refraction. I observed also, that if any where between the two outmost Circles TMF and PGA a right line, as  $\gamma \sigma$ , was crofs to the Spectrum, fo as at both ends to fall perpendicularly upon its rectilinear fides, there appeared  $M_2$ one

one and the fame Colour and degree of Colour from one end of this line to the other. I delineated therefore in a Paper the perimeter of the Spectrum FAPGMT, and in trying the third Experiment of the first Book, I held the Paper so that the Spectrum might fall upon this delineated Figure, and agree with it exactly, whilft an Affistant whose Eyes for distinguishing Colours were more critical than mine, did by right lines ab, yob, c, or. drawn crois the Spectrum, note the confines of the Colours that is of the red  $M \propto \beta F$  of the orange  $\alpha \gamma \wedge \beta$ , of the yellow yeso, of the green end, of the blue nixe, of the indico  $i \times \mu_R$ , and of the violet  $\times GA\mu$ . And this operation being divers times repeated both in the fame and in feveral Papers, I found that the Obfervations agreed well enough with one another, and that the rectilinear fides MG and FA were by the faid erofs lines divided after the manner of a mufical Chord. Let GM be produced to X, that MX may be equal to G M, and conceive G X, XX, X, X, X, X, X, X, X, MX, to be in proportion to one another, as the numbers  $I, \frac{8}{9}, \frac{5}{6}, \frac{3}{4}, \frac{2}{3}, \frac{3}{5}, \frac{9}{16}, \frac{1}{2}$ , and fo to reprefent the. Chords of the Key, and of a Tone, a third Minor, a fourth, a fifth, a fixth Major, a feventh, and an eighth. above that Key: And the intervals Ma, ay, ye, en, ni,  $^{1\lambda}$ , and  $^{\lambda}G$ , will be the spaces which the several Colours (red, orange, yellow, green, blue, indico, violet). take up.

Now these intervals or spaces subtending the differences of the refractions of the rays going to the limits. of those Colours, that is, to the points M,  $\alpha$ ,  $\gamma$ , s, n, t,  $\lambda$ , G, may without any sensible Error be accounted proportional to the differences of the fines of refraction of those rays

rays having one common fine of incidence, and therefore fince the common fine of incidence of the most and least refrangible rays out of Glass into Air was, (by a method described above ) found in proportion to their fines of refraction, as 50 to 77 and 78, divide the dif-ference between the fines of refraction 77 and 78, as the line G M is divided by those intervals, you will have 77,  $77_{\overline{s}}^{1}$ ,  $77_{\overline{s}}^{1}$ ,  $77_{\overline{s}}^{1}$ ,  $77_{\overline{s}}^{1}$ ,  $77_{\overline{s}}^{2}$ ,  $77_{\overline{s}}^{2}$ ,  $77_{\overline{s}}^{2}$ , 78, the fines of refraction of those rays out of Glass into Air, their common fine of incidence being 50. So then the fines of the incidences of all the red-making rays out of Glafs into Air, were to the fines of their refractions, not greater than 50 to 77, nor lefs than 50 to  $77_{s}^{t}$ , but varied from one another according to all interme-diate Proportions. And the fines of the incidences of the green-making rays [were to the fines of their refractions in all proportions from that of 50 to  $77_{5}^{t}$ , unto that of 50 to  $77_{2}^{t}$ . And by the like limits above-mentioned were the refractions of the rays be-longing to the reft of the Colours defined the fines of longing to the reft of the Colours defined, the fines of the red-making rays extending from 77 to  $77\frac{1}{3}$ , those of the orange-making from  $77\frac{1}{3}$  to  $77\frac{1}{3}$ , those of the yel-low-making from  $77\frac{1}{5}$  to  $77\frac{1}{3}$ , those of the green-making from  $77\frac{1}{5}$  to  $77\frac{1}{2}$ , those of the blue-making from  $77\frac{1}{2}$  to  $77\frac{2}{3}$ , those of the indico-making from  $77\frac{2}{3}$  to  $77\frac{2}{9}$ , and those of the violet from  $77\frac{2}{9}$  to 78. These are the Laws of the refractions made out of

[93]

These are the Laws of the refractions made out of Glass into Air, and thence by the three Axioms of the first Book the Laws of the refractions made out of Air into Glass are easily derived.

#### EXPER.

# L94]

### EXPER. VIII.

I found moreover that when Light goes out of Air through feveral contiguous refracting Mediums as through Water and Glass, and thence goes out again into Air, whether the refracting fuperficies be parallel or inclined to one another, that Light as often as by contrary refractions 'tis fo corrected, that it emergeth in lines parallel to those in which it was incident, continues ever after to be white. But if the emergent rays be inclined to the incident, the whitenefs of the emerging Light will by degrees in paffing on from the place of emergence, become tinged in its edges with Colours. This I tryed by refracting Light with Prifms of Glass within a prismatick Veffel of Water. Now those Colours argue a diverging and separation of the heterogeneous rays from one another by means of their unequal refractions, as in what follows will more fully appear. And, on the contrary, the permanent whitenels argues, that in like incidences of the rays there is no fuch feparation of the emerging rays, and by confequence no inequality of their whole refractions. Whence I feem to gether the two following Theorems.

I. The Exceffes of the fines of refraction of feveral forts of rays above their common fine of incidence when the refractions are made out of divers denfer mediums immediately into one and the fame rarer medium, are to one another in a given Proportion. I thought at first  $\frac{1}{4}$ when the angles of Incidence out of air into some other medium usere go or as near as could be go degrees, they Cotan out The the angles of Refection were in a given Proportion. But when the angles of Refection were in a given Proportion. But when the angles of Refection were in a given Proportion. But when the angles of Refection were in a given Proportion. But when the angles of Refection were in a given Proportion. But 2. The Proportion of the fine of incidence to the fine of refraction of one and the fame fort of rays out of one medium into another, is composed of the Proportion of the fine of incidence to the fine of refraction out of the first medium into any third medium, and of the Proportion of the fine of incidence to the fine of refraction out of that third medium into the second medium.

By the first Theorem the refractions of the rays of every fort made out of any medium into Air are known by having the refraction of the rays of any one fort. Asfor instance, if the refractions of the rays of every fort out of Rain-water into Air be defired, let the common. fine of incidence out of Glass into Air be subducted from the fines of refraction, and the Exceffes will be 27,  $27_{\overline{5}}^{1}$ ,  $27_{\overline{5}}^{1}$ ,  $27_{\overline{5}}^{1}$ ,  $27_{\overline{5}}^{1}$ ,  $27_{\overline{5}}^{2}$ ,  $27_{\overline{5}}^{2}$ ,  $27_{\overline{9}}^{2}$ , 28. Suppose now that the fine of incidence of the least refrangible rays be to their fine of refraction out of Rain-water into Air as three to four, and fay as I the difference of those fines is to 3 the fine of incidence, fo is 27 the least of the. Exceffes above-mentioned to a fourth number 81; and. 81 will be the common fign of incidence out of Rainwater into Air, to which fine if you add all the abovementioned Exceffes you will have the defired fines of the refractions 108, 108; 108; 108; 108; 108; 108;  $108^{\frac{7}{2}}$ , 109.

By the latter Theorem the refraction out of one medium into another is gathered as often as you have the refractions out of them both into any third medium. As if the fine of incidence of any ray out of Glais into Air be to its fine of refraction as 20 to 31, and the fine of incidence of the fame ray out of Air into Water, be to to its fine of refraction as four to three; the fine of incidence of that ray out of Glafs into Water will be to its fine of refraction as 20 to 31 and 4 to 3 joyntly, that is, as the Factum of 20 and 4 to the Factum of 31 and 3, or as 80 to 93.

And these Theorems being admitted into Opticks, there would be fcope enough of handling that Science voluminoully after a new manner; not only by teaching those things which tend to the perfection of vision, but alio by determining mathematically all kinds of Phænomena of Colours which could be produced by refractions. For to do this, there is nothing elfe requifite than to find out the feparations of heterogeneous rays, and their various mixtures and proportions in every mixture. By this way of arguing I invented almost all the Phænomena defcribed in these Books, beside some others lefs neceffary to the Argument; and by the fucceffes I met with in the tryals, I dare promife, that to him who shall argue truly, and then try all things with good Glaffes and fufficient circumspection, the expected event will not be wanting. But he is first to know what Colours will arife from any others mixt in any affigned Proportion.

#### PROP. IV. THEOR. III.

\* Colours may be produced by composition which shall be like to the Colours of homogeneal Light as to the appearance of Colour, but not as to the immutability of Colour and constitution of Light. And those Colours by how much they are more compounded by so much are they less full and inteuse, and by too much composition they may be diluted diluted and weakened till they ceafe. There may be also Colours produced by composition, which are not fully

like any of the Colours of homogeneal Light.

For a mixture of homogeneal red and yellow com- \* pounds an orange, like in appearance of Colour to that orange which in the feries of unmixed prifmatick Colours lies between them; but the Light of one orange is homogeneal as to refrangibility, that of the other is heterogeneal, and the Colour of the one, if viewed through a Prism, remains unchanged, that of the other is changed and refolved into its component Colours red and yellow. And after the fame manner other neighbouring homogeneal Colours may compound new Colours, like the intermediate homogeneal ones, as yellow and green, the Colour between them both, and afterwards, if blue be added, there will be made a green the middle Colour of the three which enter the compofition. For the yellow and blue on either hand, if they are equal in quantity they draw the intermediate green equally towards themfelves in composition, and so keep it as it were in equilibrio, that it verge not more to the yellow on the one hand, than to the blue on the other, but by their mixt actions remain still a middle Colour. To this mixed green there may be further added fome red and violet, and yet the green will not prefently ceafe but only grow lefs full and vivid, and by increafing the red and violet it will grow more and more dilute, until by the prevalence of the added Colours it be overcome and turned into whitenefs, or fome other Colour. So if to the Colour of any homogeneal Light, the Sun's white Light composed of all forts of rays be N added,

added, that Colour will not vanish or change its species but be diluted, and by adding more and more white it will be diluted more and more perpetually. Lastly, if red and violet be mingled, there will be generated according to their various Proportions various Purples, fuch as are not like in appearance to the Colour of any homogeneal Light, and of these Purples mixt with yellow and blue may be made other new Colours.

### PROP. V. THEOR. IV.

Whiteness and all grey Colours between white and black, may be compounded of Colours, and the whiteness of the Sun's Light is compounded of all the primary Colours mixt in a due proportion.

## The Proof by Experiments.

#### EXPER. IX.

Fig. 5.

The Sun fhining into a dark Chamber through a little round Hole in the Window flut, and his Light being there refracted by a Prifm to caft his coloured. Image PT upon the oppofite Wall : I held a white Paper V to that Image in fuch manner that it might be illuminated by the coloured Light reflected from thence, and yet not intercept any part of that Light in its paffage from the Prifm to the Spectrum. And I found that when the Paper was held nearer to any Colour than to the reft, it appeared of that Colour to which it approached neareft; but when it was equally or almoft equally

equally diftant from all the Colours, so that it might be equally illuminated by them all it appeared white. And in this last situation of the Paper, if some Colours were intercepted, the Paper loft its white Colour, and appeared of the Colour of the reft of the Light which was not intercepted. So then the Paper was illuminated with Lights of various Colours, namely, red, yellow, green, blue and violet, and every part of the Light re-tained its proper Colour, until it was incident on the Paper, and became reflected thence to the Eye ; fo that if it had been either alone (the reft of the Light being intercepted) or if it had abounded most and been predominant in the Light reflected from the Paper, it would have tinged the Paper with its own Colour; and yet being mixed with the reft of the Colours in a due proportion, it made the Paper look white, and therefore by a composition with the rest produced that Colour. The feveral parts of the coloured Light reflected from the Spectrum, whilft they are propagated from thence thro' the Air, do perpetually retain their proper Colours, because wherever they fall upon the Eyes of any Specta-tor, they make the several parts of the Spectrum to appear under their proper Colours. They retain there-fore their proper Colours when they fall upon the Paper V, and so by the confusion and perfect mixture of those Colours compound the whiteness of the Light reflected from thence.

## EXPER. X.

Let that Spectrum or folar Image PT fall now upon Fig. 6. the Lens MN above four Inches broad, and about fix Feet

 $N_2$ 

Feet diftant from the Prifin ABC, and fo figured that it may cause the coloured Light which divergeth from the Prism to converge and meet again at its Focus G, about fix or eight Feet diftant from the Lens, and there to fall perpendicularly upon a white Paper DE. And if you move this Paper to and fro, you will perceive that near the Lens, as at de, the whole folar Image (fuppofe at pt) will appear upon it intenfly coloured after the manner above-explained, and that by receding from the Lens those Colours will perpetually come towards one another, and by mixing more and more dilute one another continually, until at length the Paper come to the Focus G, where by a perfect mixture they will wholly vanish and be converted into whiteness, the whole Light appearing now upon the Paper like a little white Circle. And afterwards by receding further from the Lens, the rays which before converged will now cross one another in the Focus G, and diverge from thence, and thereby make the Colours to appear again, but yet in a contrary order ; fuppole at de, where the red t is now above which before was below, and the violet p is below which before was above.

Let us now ftop the Paper at the Focus G where the Light appears totally white and circular, and let us confider its whitenefs. I fay, that this is composed of the converging Colours. For if any of those Colours be intercepted at the Lens, the whitenefs will cease and degenerate into that Colour which ariseth from the composition of the other Colours which are not intercepted. And then if the intercepted Colours be let pass and fall upon that compound Colour, they mix with it, and by their mixture reftore the whitenefs. So So if the violet, blue and green be intercepted, the remaining yellow, orange and red will compound upon the Paper an orange, and then if the intercepted Colours be let pafs they will fall upon this compounded orange, and together with it decompound a white. So alfo if the red and violet be intercepted, the remaining yellow, green and blue, will compound a green upon the Paper, and then the red and violet being let pafs will fall upon this green, and together with it decompound a white. And that in this composition of white the feveral rays do not fuffer any change in their colorific qualities by acting upon one another, but are only mixed, and by a mixture of their Colours produce white, may further appear by thefe Arguments.

If the Paper be placed beyond the Focus G, fuppofe at  $\delta_i$ , and then the red Colour at the Lens be alternately intercepted, and let pafs again, the violet Colour on the Paper will not fuffer any change thereby, as it ought to do if the feveral forts of rays acted upon one another in the Focus G, where they crofs. Neither will the red upon the Paper be changed by any alternate ftopping, and letting pafs the violet which croffeth it.

And if the Paper be placed at the Focus G, and the white round Image at G be viewed through the Prifm HIK, and by the refraction of that Prifm be translated to the place rv, and there appear tinged with various Colours, namely, the violet at v and red at r, and others between, and then the red Colour at the Lens be often ftopt and let pass by turns, the red at r will accordingly difappear and return as often, but the violet at v will not thereby fuffer any change. And so by ftopping and letting pass alternately the blue at the Lens, Lens, the blue at r will accordingly difappear and return, without any change made in the red at r. The red therefore depends on one fort of rays, and the blue on another fort, which in the Focus G where they are commixt do not act on one another. And there is the fame reafon of the other Colours.

I confidered further, that when the most refrangible rays P p, and the least refrangible ones T t, are by con-verging inclined to one another, the Paper, if held very oblique to those rays in the Focus G, might reflect one fort of them more copioully than the other fort, and by that means the reflected Light would be tinged in that Focus with the Colour of the predominant rays, provided those rays feverally retained their Colours or colorific qualities in the composition of white made by them in that Focus. But if they did not retain them in that white, but became all of them feverally endued there with a disposition to strike the sense with the perception of white, then they could never lose their white-ness by fuch reflexions. I inclined therefore the Paper to the rays very obliquely, as in the fecond Experiment of this Book, that the most refrangible rays might be more copioully reflected than the reft, and the whitenefs at length changed fucceffively into blue, indico and violet. Then I inclined it the contrary way, that the most refrangible rays might be more copious in the reflected Light than the reft, and the whitenefs turned

fucceffively to yellow, orange and red. Laftly, I made an Inftrument X Y in fashion of a Comb, whose Teeth being in number fixteen were about an Inch and an half broad, and the intervals of the Teeth about two Inches wide. Then by interposing fuc-

fucceffively the Teeth of this Inftrument near the Lens,. I intercepted part of the Colours by the interpofed Tooth, whilft the reft of them went on through the in-terval of the Teeth to the Paper DE, and there painted a round solar Image. But the Paper I had first placed so, that the Image might appear white as often as the Comb was taken away; and then the Comb being as was faid interposed, that whiteness by reason of the intercepted part of the Colours at the Lens did always change into the Colour compounded of those Colours which were not intercepted, and that Colour was by the motion of the Comb perpetually varied fo, that in the paffing of every Tooth over the Lens all. these Colours red, yellow, green, blue and purple, did always fucceed one another. I caused therefore all the Teeth to pass fucceffively over the Lens, and when the motion was flow, there appeared a perpetual fucceffion of the Colours upon the Paper : But if I fo much acce-lerated the motion, that the Colours by reafon of their quick succession could not be distinguished from one another, the appearance of the fingle Colours ceafed. There was no red, no yellow, no green, no blue, nor purple to be seen any longer, but from a confusion of them all there arose one uniform white Colour. Of the Light which now by the mixture of all the Colours appeared white, there was no part really white. One part was red, another yellow, a third green, a fourth blue, a fifth purple, and every part retains its proper Colour till it ftrike the Senforium. If the impreffions follow one another flowly, fo that they may be feverally perceived, there is made a diftinct fenfation of all the Colours one after another in a continual fucceffion. But

But if the imprefiions follow one another fo quickly that they cannot be feverally perceived, there arifeth out of them all one common sensation, which is neither of this Colour alone nor of that alone, but hath it felf indifferently to 'em all, and this is a fensation of whitenefs. By the quicknefs of the fucceffions the impreffions of the feveral Colours are confounded in the Senforium, and out of that confusion ariseth a mixt senfation. If a burning Coal be nimbly moved round in a Circle with Gyrations continually repeated, the whole Circle will appear like fire; the reason of which is, that the fenfation of the Coal in the feveral places of that Circle remains imprest on the Senforium, until the Coal return again to the fame place. And fo in a quick confecution of the Colours the impression of every Colour remains in the Sensorium, until a revolution of all the Colours be compleated, and that first Colour return again. The impressions therefore of all the successive Colours are at once in the Senforium, and joyntly ftir up a fenfation of them all; and fo it is manifest by this Experiment, that the commixt impreffions of all the Colours do ftir up and beget a fensation of white, that is, that whitenefs is compounded of all the Colours.

-X-

And if the Comb be now taken away, that all the Colours may at once pais from the Lens to the Paper, and be there intermixed, and together reflected thence to the Spectators Eyes ; their imprefions on the Senforium being now more fubtily and perfectly commixed there, ought much more to ftir up a fenfation of whitenefs.

You

You may inftead of the Lens use two Prisms HIK and LMN, which by refracting the coloured Light the contrary way to that of the first refraction, may make the diverging rays converge and meet again in G, as you see it represented in the seventh Figure. For Fig. 7. where they meet and mix they will compose a white Light as when a Lens is used.

### EXPER. XI.

Let the Sun's coloured Image PT fall upon the Wall Fig. 8. of a dark Chamber, as in the third Experiment of the first Book, and let the fame be viewed through a Prifm abc, held parallel to the Prifm ABC, by whofe refra-Ction that Image was made, and let it now appear lower than before, suppose in the place S over against the redcolour T. And if you go near to the Image PT, the Spectrum S will appear oblong and coloured like the Image PT; but if you recede from it, the Colours of the Spectrum S will be contracted more and more, and at length vanish, that Spectrum S becoming perfectly round and white; and if you recede yet further, the Colours will emerge again, but in a contrary order. Now that Spectrum S appears white in that cafe when the rays of feveral forts which converge from the feveral parts of the Image PT, to the Prisin abc, are fo refracted unequally by it, that in their paffage from the Prifin to the Eye they may diverge from one and the fame point of the Spectrum S, and fo fall afterwards upon one and the fame point in the bottom of the Eye, and there be mingled.

0

And

And further, if the Comb be here made use of, by whose Teeth the Colours at the Image PT may be fucceffively intercepted ; the Spectrum S when the Comb is moved flowly will be perpetually tinged with fucceffive Colours : But when by accelerating the motion of the Comb, the fucceffion of the Colours is fo quick that they cannot be severally seen, that Spectrum S, by a confused and mixt sensation of them all, will appear white.

#### EXPER. XII.

Fig. 9.

The Sun shining through a large Prism ABC upon a Comb X Y, placed immediately behind the Prifm, his Light which passed through the interstices of the Teeth fell upon a white Paper DE. The breadths of the Teeth were equal to their interffices, and feven Teeth. together with their interffices took up an Inch in breadth. Now when the Paper was about two or three Inches distant from the Comb, the Light which paffed through its feveral interffices painted fo many ranges of Colours kl, mn, op, qr, &c. which were parallel to one another and contiguous, and without any mixture of white. And these ranges of Colours, if the Comb was moved continually up and down with a reciprocal motion, ascended and descended in the Paper, and when the motion of the Comb was fo quick, that the Colours could not be diffinguished from one another, the whole Paper by their confusion and mixture in the Senforium appeared white.

Let

## [107]

Let the Comb now reft, and let the Paper be removed further from the Prifin, and the feveral ranges of Colours will be dilated and expanded into one another more and more, and by mixing their Colours will dilute one another, and at length, when the diffance of the Paper from the Comb is about a Foot, or a little more (fuppofe in the place 2D 2E) they will to far dilute one another as to become white.

With any Obftacle let all the Light be now ftopt which paffes through any one interval of the Teeth, fo that the range of Colours which comes from thence may be taken away, and you will fee the Light of the reft of the ranges to be expanded into the place of the range taken away, and there to be coloured. Let the intercepted range pafs on as before, and its Colours falling upon the Colours of the other ranges, and mixing with them, will reftore the whitenefs.

Let the Paper 2D 2E be now very much inclined to the rays, fo that the most refrangible rays may be more copiously reflected than the reft, and the white Colour of the Paper through the excess of those rays will be changed into blue and violet. Let the Paper be as much inclined the contrary way, that the least refrangible rays may be now more copiously reflected than the reft, and by their excess the whiteness will be changed into yellow and red. The feveral rays therefore in that white Light do retain their colorific qualities, by which those of any fort, when-ever they become more copious than the reft, do by their excess and predominance cause their proper Colour to appear.

And

# [ 801 ]

And by the fame way of arguing, applied to the third Experiment of this Book, it may be concluded, that the white Colour of all refracted Light at its very first emergence, where it appears as white as before its incidence, is compounded of various Colours.

### EXPER. XIII.

In the foregoing Experiment the feveral intervals of the Teeth of the Comb do the office of so many Prisms, every interval producing the Phænomenon of one Prifm. Whence instead of those intervals using several Prisms, I try'd to compound whiteness by mixing their Colours, and did it by using only three Prilins, as also by using only Fig. 10. two as follows. Let two Prisms ABC and a bc, whose refracting Angles B and b are equal, be fo placed parallel to one another, that the refracting Angle B of the one may touch the Angle c at the base of the other, and their planes CB and cb, at which the rays emerge, may lye in directum. Then let the Light trajected through them fall upon the Paper MN, distant about 8 or 12 Inches from the Prifins. And the Colours generated by the interior limits B and c of the two Prisms, will: be mingled at PT, and there compound white. For if either Prism be taken away, the Colours made by the other will appear in that place PT, and when the Prisin is reftored to its place again, fo that its Colours may there fall upon the Colours of the other, the mixture of them both will reftore the whitenels.

This

This Experiment fucceeds alfo, as I have tryed, when the Angle b of the lower Prifm, is a little greater than the Angle B of the upper, and between the interior Angles B and c, there intercedes some space B c, as is represented in the Figure, and the refracting planes BC and bc, are neither in directum, nor parallel to one another. For there is nothing more requifite to the fuccefs of this Experiment, than that the rays of all forts may be uniformly mixed upon the Paper in the place PT. If the most refrangible rays coming from the fuperior Prism take up all the space from M to P, the rays of the fame fort which come from the inferior Prifin ought to begin at P, and take up all the reft of the fpace from thence towards N. If the leaft refrangible rays coming from the fuperior Prism take up the space MT, the rays of the fame kind which come from the other Prism ought to begin at T, and take up the remaining space TN. If one fort of the rays which have intermediate degrees of refrangibility, and come from the fuperior Prism be extended through the space MQ, and another fort of those rays through the space MR, and a third fort of them through the space MS, the same forts of rays coming from the lower Prism, ought to illuminate the remaining spaces QN, RN, SN respe-Stively. And the fame is to be underftood of all the other forts of rays. For thus the rays of every fort will be scattered uniformly and evenly through the whole space MN, and so being every where mixt in the same proportion, they must every where produce the fame Colour. And therefore fince by this mixture they produce white in the exterior fpaces MP and TN, they must also produce white in the interior space PT. This is

## [ 110 ]

is the reafon of the composition by which whiteness was produced in this Experiment, and by what other way soever I made the like composition the result was whiteness.

Laftly, If with the Teeth of a Comb of a due fize, the coloured Lights of the two Prifms which fall upon the fpace PT be alternately intercepted, that fpace PT, when the motion of the Comb is flow, will always appear coloured, but by accelerating the motion of the Comb fo much, that the fucceflive Colours cannot be diftinguished from one another, it will appear white.

### EXPER. XIV.

-Ye

Hitherto I have produced whitenefs by mixing the Colours of Prifms. If now the Colours of natural Bodies are to be mingled, let Water a little thickned with Soap be agitated to raife a froth, and after that froth has ftood a little, there will appear to one that fhall view it intently various Colours every where in the furfaces of the feveral Bubbles; but to one that fhall go fo far off that he cannot diftinguifh the Colours from one another, the whole froth will grow white with a perfect whitenefs.

### EXPER. XV.

Laftly, in attempting to compound a white by mixing the coloured Powders which Painters ufe, I confidered that all coloured Powders do fupprefs and ftop in them a very confiderable part of the Light by which they

they are illuminated. For they become coloured by reflecting the Light of their own Colours more copioufly, and that of all other Colours more sparingly, and yet they do not reflect the Light of their own Colours fo copioully as white Bodies do. If red Lead, for instance, and a white Paper, be placed in the red Light of the coloured Spectrum made in a dark Chamber by the refraction of a Prisin, as is described in the third Eperiment of the first Book ; the Paper will appear more lucid than the red Lead, and therefore reflects the redmaking rays more copioufly than red Lead doth. And if they be held in the Light of any other Colour, the Light reflected by the Paper will exceed the Light re-flected by the red Lead in a much greater proportion. And the like happens in Powders of other Colours. And therefore by mixing fuch Powders we are not to expect a ftrong and full white, fuch as is that of Paper, but some dusky obscure one, such as might arise from a mixture of light and darkness, or from white and black, that is, a grey, or dun, or ruffet brown, fuch as are the Colours of a Man's Nail, of a Moufe, of Ashes, of ordinary Stones, of Mortar, of Dust and Dirt in Highways, and the like. And fuch a dark white I have often produced by mixing coloured Powders. For thus one part of red Lead, and five parts of Viride Æris, composed a dun Colour like that of a Mouse. For these two Colours were feverally fo compounded of others, that in both together were a mixture of all Colours; and. there was less red Lead used than Viride Æris, because of the fulness of its Colour. Again, one part of red Lead, and four parts of blue Bife, composed a dun Colour verging a little to purple, and by adding to this a certain.

certain mixture of Orpiment and Viridi Æris in a due proportion, the mixture loft its purple tincture, and be-came perfectly dun. But the Experiment fucceeded beft without Minium thus. To Orpiment I added by little and little a certain full bright purple, which Painters use until the Orpiment ceased to be yellow, and became of a pale red. Then I diluted that red by adding a little Viride Æris, and a little more blue Bise than Viridi Æris, until it became of fuch a grey or pale white, as verged to no one of the Colours more than to another. For thus it became of a Colour equal in whiteneis to that of Afhes or of Wood newly cut, or of a Man's Skin. The Orpiment reflected more Light than did any other of the Powders, and therefore conduced more to the whitenefs of the compounded Colour than they. To affign the proportions accurately may be difficult, by reason of the different goodness of Powders of the fame kind. Accordingly as the Colour of any Powder is more or lefs full and luminous, it ought to be used in a less or greater proportion.

Now confidering that these grey and dun Colours may be also produced by mixing whites and blacks, and by confequence differ from perfect whites not in Species of Colours but only in degree of luminoufnefs, it is ma-nifest that there is nothing more requisite to make them perfectly white than to increase their Light sufficiently; and, on the contrary, if by increasing their Light they can be brought to perfect whiteness, it will thence also follow, that they are of the same Species of Colour with the best whites, and differ from them only in the quantity of Light. And this I tryed as follows. I took the third of the above-mentioned grey mixtures (that . . .

## [113]

(that which was compounded of Orpiment, Purple, Bife and Viride Æris) and rubbed it thickly upon the tloor of my Chamber, where the Sun shone upon it through the opened Cafement; and by it, in the fhadow, I laid a piece of white Paper of the same bigness. Then going from them to the distance of 12 or 18 Feet, fo that I could not discern the unevenness of the surface of the Powder, nor the little shadows let fall from the gritty particles thereof; the Powder appeared intenfly white, so as to transcend even the Paper it self in whitenefs, especially if the Paper were a little shaded from the Light of the Clouds, and then the Paper compared with the Powder appeared of fuch a grey Colour as the Powder had done before. But by laying the Paper where the Sun shines through the Glass of the Window, or by fhutting the Window that the Sun might fhine through the Glass upon the Powder, and by such other fit means of increasing or decreasing the Lights wherewith the Powder and Paper were illuminated, the Light wherewith the Powder is illuminated may be made ftronger in fuch a due proportion than the Light wherewith the Paper is illuminated, that they shall both appear exactly alike in whiteness. For when I was trying this, a Friend coming to vifit me, I ftopt him at the door, and before I told him what the Colours were, or what I was doing; I askt him, Which of the two whites were the best, and wherein they differed ? And after he had at that diffance viewed them well, he answered, That they were both good whites, and that he could not fay which was best, nor wherein their Colours differed. Now if you confider, that this white of the Powder in the Sun-shine was compounded of the Colours P

Colours which the component Powders (Orpinent, Purple, Bife, and Viride Æris) have in the fame Sunfhine, you must acknowledge by this Experiment, as well as by the former, that perfect whiteness may be compounded of Colours.

From what has been faid it is alfo evident, that the whitenefs of the Sun's Light is compounded of all the Colours wherewith the feveral forts of rays whereof that Light confifts, when by their feveral refrangibilities they are feparated from one another, do tinge Paper or any other white Body whereon they fall. For those Colours by Prop. 2. are unchangeable, and whenever all those rays with those their Colours are mixt again, they reproduce the fame white Light as before.

#### PROP. VI. PROB. II.

In a mixture of primary Colours, the quantity and quality of each being given, to know the Colour of the compound.

Fig. II.

With the Center O and Radius OD defcribe a Circle ADF, and diffinguish its circumference into feven parts DE, EF, FG, GA, AB, BC, CD, proportional to the feven mufical Tones or Intervals of the eight Sounds, Sol, la, fa, fol, la, mi, fa, fol, contained in an Eight; that is, proportional to the numbers  $\frac{1}{9}$ ,  $\frac{1}{10}$ ,  $\frac{1}{9}$ ,  $\frac{1}{10}$ ,  $\frac{1}{16}$ ,  $\frac{1}{10}$ . Let the first part DE represent a red Colour, the fecond EF orange, the third FG yellow, the fourth GH green, the fifth AB blue, the fixth BC indico, and the feventh CD violet. And conceive that these are all the Colours of uncompounded. Light gradually paffing

paffing into one another, as they do when made by Prisins; the circumference DEFGABCD, representing the whole feries of Colours from one end of the Sun's coloured Image to the other, fo that from D to E be all degrees of red, at E the mean Colour between red and orange, from E to F all degrees of orange, at F the mean between orange and yellow, from F to G all degrees of yellow, and fo on. Let p be the center of gravity of the Arch DE, and q, r, s, t, v, x, the centers of gravity of the Arches EF, FG, GA, AB, BC and CD refpectively, and about those centers of gra-vity let Circles proportional to the number of rays of each Colour in the given mixture be described; that is, the circle p proportional to the number of the red-ma-king rays in the mixture, the Circle q proportional to the number of the orange-making rays in the mixture, and fo of the reft. Find the common center of gravity of all those Circles p, q, r, s, t, v, x. Let that center be Z; and from the center of the Circle ADF, through Z to the circumference, drawing the right line OY, the place of the point Y in the circumference shall shew the Colour arising from the composition of all the Colours in the given mixture, and the line OZ shall be proportional to the fulness or intenseness of the Colour, that is, to its diftance from whitenefs. As if Y fall in the middle between F and G, the compounded Colour shall be the best yellow; if Y verge from the middle towards F or G, the compounded Colour shall accordingly be a yellow, verging towards orange or green. If Z fall upon the circumference the Colour shall be intense and florid in the highest degree; if it fall in the mid way between the circumference and center, it shall be P 2 but

but half so intense, that is, it shall be such a Colour as would be made by diluting the intenfeft yellow with an equal quantity of whiteness; and if it fall upon the center O, the Colour shall have lost all its intensenes, and become a white. But it is to be noted, That if the point Z fall in or near the line OD, the main ingredients being the red and violet, the Colour compounded shall not be any of the prismatic Colours, but a purple, inclining to red or violet, accordingly as the point Z lieth on the fide of the line DO towards E or towards C, and in general the compounded violet is more bright and more fiery than the uncompounded. Also if only two of the primary Colours which in the Circle are oppofite to one another be mixed in an equal proportion, the point Z shall fall upon the center O, and yet the Colour compounded of those two shall not be perfectly white, but some faint anonymous Colour. For I could never yet by mixing only two primary Colours produce a perfect white. Whether it may be compounded of a mixture of three taken at equal diftances in the circumference I do not know, but of four or five I do not much question but it may. But these are curiosities of little or no moment to the understanding the Phænomena of nature. For in all whites produced by nature, there afes to be a mixture of all forts of rays, and by confequence a composition of all Colours.

To give an inftance of this Rule; fuppofe a Colour is compounded of thefe homogeneal Colours, of violet 1 part, of indico 1 part, of blue 2 parts, of green 3 parts, of yellow 5 parts, of orange 6 parts, and of red 10 parts. Proportional to thefe parts I defcribe the Circles x, v, t, s, r, q, p refpectively, that is, fo that if the Circle x be be 1, the Circle v may be 1, the Circle t 2, the Circle's 3, and the Circles r, q and p, 5, 6 and 10. Then I find Z the common center of gravity of these Circles, and through Z drawing the line OY, the point Y falls upon the circumference between E and F, fome thing nearer to E than to F, and thence I conclude, that the Colour compounded of these ingredients will be an orange, verging a little more to red than to yellow. Also I find that OZ is a little less than one half of OY, and thence I conclude, that this orange hath a little less than half the fulnels or intensents of an uncompounded orange; that is to fay, that it is fuch an orange as may be made by mixing an homogeneal orange with a good white in the proportion of the line OZ to the line ZY, this proportion being not of the quantities of mixed orange and white powders, but of the quantities of the lights reflected from them.

This Rule I conceive accurate enough for practife, though not mathematically accurate ; and the truth of it may be fufficiently proved to fenfe, by ftopping any of the Colours at the Lens in the tenth Experiment of this Book. For the reft of the Colours which are not ftopped, but pafs on to the Focus of the Lens, will there compound either accurately or very nearly fuch a Colour as by this Rule ought to refult from their mixture.

# [118]

# PROP. VII. THEOR. V.

All the Colours in the Universe which are made by Light, and depend not on the power of imagination, are either the Colours of homogeneal Lights, or compounded of these and that either accurately or very nearly, according to the Rule of the foregoing Problem.

For it has been proved ( in Prop.1. Lib.2.) that the changes of Colours made by refractions do not arife from any new modifications of the rays imprest by those refractions, and by the various terminations of light and fhadow, as has been the conftant and general opinion of Philosophers. It has also been proved that the feveral Colours of the homogeneal rays do constantly answer to their degrees of refrangibility, (Prop. 1. Lib. 1. and Prop.2. Lib.2.) and that their degrees of refrangibility cannot be changed by refractions and reflexions, (Prop.2. *Lib.*1.) and by confequence that those their Colours are likewise immutable. It has also been proved directly by refracting and reflecting homogeneal Lights apart, that their Colours cannot be changed, (Prop.2. Lib.2.) It has been proved also, that when the feveral forts of rays are mixed, and in croffing pafs through the fame space, they do not act on one another to as to change each others colorifick qualities, (Exper. 10. Lib.2.) but by mixing their actions in the Senforium beget a fenfation differing from what either would do apart, that is a fensation of a mean Colour between their proper Colours; and particularly when by the concourfe and mixtures of all forts of rays, a white Colour

Colour is produced, the white is a mixture of all the Colours which the rays would have apart, ( Prop. 5. Lib. 2. ) The rays in that mixture do not lofe or alter their feveral colorifick qualities, but by all their various kinds of actions mixt in the Senforium, beget a fenfation of a middling Colour between all their Colours which is whitenefs. For whitenefs is a mean between all Colours, having it felf indifferently to them all, for as with equal facility to be tinged with any of them. A red Powder mixed with a little blue, or a blue with a little red, doth not presently lose its Colour, but an white Powder mixed with any Colour is prefently tinged with that Colour, and is equally capable of being tinged with any Colour what-ever. It has been shewed alfo, that as the Sun's Light is mixed of all forts of rays, fo its whitenefs is a mixture of the Colours of all forts of rays; those rays having from the beginning their fe-veral colorific qualities as well-as-their feveral refrangibilities, and retaining them perpetually unchang'd notwithstanding any refractions or reflexions they may at any time fuffer, and that when-ever any fort of the Sun's rays is by any means (as by reflexion in Exper.  $\gamma$ . and 10. Lib. 1. or by refraction as happens in all refractions) feparated from the reft, they then manifest their proper Colours. Theie things have been proved, and the fum of all this amounts to the Proposition here to be proved. For if the Sun's Light is mixed of feveral forts of rays, each of which have originally their feveral refrangibilities and colorifick qualities, and notwithstanding their refractions and reflections, and their various feparations or mixtures, keep those their ori-ginal properties perpetually the fame without alteration ;

tion; then all the Colours in the World must be fuch as constantly ought to arife from the original colorific qualities of the rays whereof the Lights confife by which those Colours are seen. And therefore if the reason of any Colour what-ever be required, we have nothing elfe to do then to confider how the rays in the Sun's Light have by reflexions or refractions, or other causes been parted from one another, or mixed together; or otherwife to find out what forts of rays are in the Light by which that Colour is made, and in what proportion; and then by the last Problem to learn the Colour which ought to arife by mixing those rays (or their Colours) in that proportion. I fpeak here of Colours fo far as they arife from Light. For they appear fometimes by other causes, as when by the power of phantafy we fee Colours in a Dream, or a mad Man fees things before him which are not there; or when we fee Fire by ftriking the Eye, or fee Colours like the Eye of a Peacock's Feather, by preffing our Eyes in either corner whilft we look the other way. Where these and fuch like caufes interpofe not, the Colour always answers to the fort or forts of the rays whereof the Light confifts, as I have conftantly found in what-ever Phænomena of Colours I have hitherto been able to examin. I shall in the following Propositions give instances of this in the Phænomena of chiefest note.

-f-

PROP.

# 

## PROP. VIII. PROB. III.

By the discovered Properties of Light to explain the Colours made by Prisms.

Let ABC reprefent a Prism refracting the Light of Fig. 12. the Sun, which comes into a dark Chamber through a Hole F & almost as broad as the Prism, and let M N reprefent a white Paper on which the refracted Light is caft, and fuppose the most refrangible or deepest violet making rays fall upon the fpace  $P_{\pi}$ , the leaft refrangible or deepeft red-making rays upon the fpace T7, the middle fort between the Indico-making aud bluemaking rays upon the fpace  $Q_{\chi}$ , the middle fort of the green-making rays upon the space Re, the middle fort between the yellow-making and orange-making rays upon the fpace  $S_{\sigma}$ , and other intermediate forts upon intermediate spaces. For so the spaces upon which the feveral forts adequately fall will by reafon of the different refrangibility of those forts be one lower than another. Now if the Paper MN be fo near the Prifin that the fpaces P T and m do not interfere with one another, the diftance between them T  $\pi$  will be illuminated by all the forts of rays in that proportion to one another which they have at their very first coming out of the Prifm, and confequently be white. But the fpaces PT and  $\pi^{\uparrow}$ . on either hand, will not be illuminated by them all, and therefore will appear coloured. And particularly at P, where the outmost violet-making rays fall alone, the Colour must be the deepest violet. At Q where the violet-making and indico-making rays are mixed, it muft

muft be a violet inclining much to indico. At R where the violet-making, indico-making, blue-making, and one half of the green-making rays are mixed, their Colours muft (by the conftruction of the fecond Problem) compound a middle Colour between indico and blue. At S where all the rays are mixed except the red-making and orange-making, their Colours ought by the fame Rule to compound a faint blue, verging more to green than indie. And in the progress from S to T, this blue will grow more and more faint and dilute, till at T, where all the Colours begin to be mixed, it end in whitenefs.

So again, on the other fide of the white at T, where the leaft refrangible or utmost red-making rays are alone the Colour must be the deepest red. At  $\sigma$  the mixture of red and orange will compound a red inclining to orange. At  $\epsilon$  the mixture of red, orange, yellow, and one half of the green must compound a middle Colour between orange and yellow. At  $\chi$  the mixture of all Colours but violet and indico will compound a faint yellow, verging more to green than to orange. And this yellow will grow more faint and dilute continually in its progress from  $\chi$  to  $\pi$ , where by a mixture of all forts of rays it will become white.

These Colours ought to appear were the Sun's Light perfectly white: But because it inclines to yellow, the excess of the yellow-making rays whereby 'tis tinged with that Colour, being mixed with the faint blue between S and T, will draw it to a faint green. And so the Colours in order from P to T ought to be violet, indico, blue, very faint green, white, faint yellow, orange, red. Thus it is by the computation : And they that please to view.
view the Colours made by a Prism will find it so in Nature.

Thefe are the Colours on both fides the white when the Paper is held between the Prifm, and the point X where the Colours meet, and the interjacent white vanifhes. For if the Paper be held ftill farther off from the Prifm, the moft refrangible and leaft refrangible rays will be wanting in the middle of the Light, and the reft of the rays which are found there, will by mixture produce a fuller green than before. Alfo the yellow and blue will now become lefs compounded, and by confequence more intenfe than before. And this alfo agrees with experience.

And if one look through a Prifm upon a white Object encompassed with blackness or darkness, the reason of the Colours arifing on the edges is much the fame, as will appear to one that shall a little confider it. If a black Object be encompassed with a white one, the Colours which appear through the Prisin are to be derived from the Light of the white one, spreading into the Regions of the black, and therefore they appear in a contrary order to that, in which they appear when a white Object is furrounded with black. And the fame is to be underftood when an Object is viewed, whofe parts are some of them less luminous than others. For in the Borders of the more and lefs luminous parts, Colours ought always by the fame Principles to arife from the excess of the Light of the more luminous, and to be of the fame kind as if the darker parts were black, but yet to be more faint and dilute.

What

Q 2

What is faid of Colours made by Prifms may be eafily applied to Colours made by the Glaffes of Telefcopes, or Microfcopes, or by the humours of the Eye. For if the Object-glafs of a Telefcope be thicker on one fide than on the other, or if one half of the Glafs, or one half of the Pupil of the Eye be covered with any opake fubftance : the Object-glafs, or that part of it or of the Eye which is not covered, may be confidered as a Wedge with crooked fides, and every Wedge of Glafs, or other pellucid fubftance, has the effect of a Prifm in refracting the Light which paffes through it.

How the Colours in the 9th and 10th Experiments of the first Part arise from the different reflexibility of Light, is evident by what was there faid. But it is observable in the 9th Experiment, that whilst the Sun's direct Light is yellow, the excess of the blue-making rays in the reflected Beam of Light MN, fuffices only to bring that yellow to a pale white inclining to blue, and not to tinge it with a manifestly blue Colour. To obtain therefore a better blue, I used instead of the yellow Light of the Sun the white Light of the Clouds, by varying a little the Experiment as follows.

#### EXPER. XVI.

Fig. 13. Let HFG reprefent a Prifm in the open Air, and S the Eye of the Spectator, viewing the Clouds by their. Light coming into the Prifm at the plane fide FIGK, and reflected in it by its bafe HEIG, and thence going out through its plain fide HEFK to the Eye. And when the Prifm and Eye are conveniently placed, fo that the Angles of incidence and reflexion at the bafe may

may be about 40 degrees, the Spectator will fee a Bow-MN of a blue Colour, running from one end of the bafe to the other, with the concave fide towards him, and the part of the base IMNG beyond this Bow will be brighter than the other part EMNH on the other side of it. This blue Colour MN being made by no-thing elfe than by reflexion of a fpecular fuperficies, feems fo odd a Phænomenon, and fo unaccountable for by the vulgar Hypothesis of Philosophers, that I could not but think it deferved to be taken notice of. Now for understanding the reason of it, suppose the plane. ABC to cut the plane fides and bafe of the Prisin perpendicularly. From the Eye to the line BC, wherein that plane cuts the base, draw the lines Sp and St, in the-Angles Spc 50 degr. ;, and Stc 49 degr. 1/28, and the point p will be the limit beyond which none of the most refrangible rays can pass through the base of the Prism, and be refracted, whose incidence is such that they may be reflected to the Eye; and the point t will be the like limit for the least refrangible rays, that is, beyond which none of them can pass through the base, whose incidence is fuch that by reflexion they may come to the Eye. And the point r taken in the middle way between p and t, will be the like limit for the meanly refrangible rays. And therefore all the refrangible rays which fall. upon the base beyond t, that is, between t and B, and. can come from thence to the Eye will be reflected thither : But on this fide t, that is, between t and c, many, of these rays will be transmitted through the base. And all the most refrangible rays which fall upon the base beyond p, that is, between p and B, and can by reflexion come from thence to the Eye, will be reflected thither

thither, but every where between t and c, many of thefe rays will get through the bafe and be refracted ; and the fame is to be underftood of the meanly refrangible rays on either fide of the point r. Whence it follows, that the bale of the Prism must every where between t and B, by a total reflexion of all forts of rays to the Eye, look white and bright. And every where between P and C, by reason of the transmission of many rays of every fort, look more pale, obfcure and dark. But at r, and in other places between p and t, where all the more refrangible rays are reflected to the Eye, and many of the lefs refrangible are transmitted, the excess of the most refrangible in the reflected Light will tinge that Light with their Colour, which is violet and blue. And this happens by taking the line Cprt B any where between the ends of the Prifin HG and EI.

### PROP. IX. PROB. IV.

# By the discovered Properties of Light to explain the Colours of the Rain-bow.

This Bow never appears but where it Rains in the Sun-fhine, and may be made artificially by fpouting up Water which may break aloft, and fcatter into Drops, and fall-down like Rain. For the Sun fhining upon thefe Drops certainly caufes the Bow to appear to a Spectator ftanding in a due polition to the Rain and Sun. And hence it is now agreed upon, that this Bow is made by refraction of the Sun's Light in Drops of falling Rain. This was underftood by fome of the Ancients, and of late more fully difcovered and explained by the Famous Antonius

# 

Antonius de Dominis Archbishop of Spilato; in his Book. De Radiis Visus & Lucis, published by his Friend Bartolus at Venice, in the Year 1611, and written above twenty Years before. For he teaches there how the interior Bow is made in round Drops of Rain by two refractions of the Sun's Light, and one reflexion between them, and the exterior by two refractions and two forts of reflexions between them in each Drop of Water, and proves his Explications by Experiments made with a Phial full of Water, and with Globes of Glafsfilled with Water, and placed in the Sun to make the Colours of the two Bows appear in them. The fame Explication Des-Cartes hath purfued in his Meteors, and mended that of the exterior Bow. But whilft they understood not the true origin of Colours, it's necessary to purfue it here a little further. For understanding therefore how the Bow is made, let a Drop of Rain or any other fpherical transparent Body be represented by the Sphere BNFG, described with the Center C, and Fig. 14-Semi-diameter CN. And let AN be one of the Sun's rays incident upon it at N, and thence refracted to F, where let it either go out of the Sphere by refraction towards V, or be reflected to G ; and at G let it either go. out by refraction to R, or be reflected to H; and at H let it go out by refraction towards S, cutting the incident ray in Y; produce A N and R G, till they meet in X, and upon AX and NF let fall the perpendiculars. CD and CE, and produce CD till it fall upon the circumference at L. Parallel to the incident ray A N draw. the Diameter BQ, and let the fine of incidence out of Air into Water be to the fine of refraction as I to R. Now if you suppose the point of incidence N to. move

[ 128 ] move from the point B, continually till it come to L, the Arch QF will first increase and then decrease, and fo will the Angle AXR which the rays AN and GR contain; and the Arch QF and Angle AXR will be biggest when ND is to CN as VII-RR to V3 RR, in which cafe NE will be to ND as 2 R to I. Alfo the Angle AYS which the rays A N and HS contain will first decrease, and then increase and grow least when ND is to CN as VII-RR to V8 RR, in which cafe NE will be to ND as 3 R to I. And fo the Angle which the next emergent ray ( that is, the emergent ray after three reflexions) contains with the incident ray AN will come to its limit when ND is to CN as VII-RR to  $V_{15}$  RR, in which cafe N E will be to N D as 4 R to I, and the Angle which the ray next after that emergent, that is, the ray emergent after four reflexions, contains with the incident will come to its limit, when ND is to  $CN_{\ell}$  as  $\sqrt{11-RR}$  to  $\sqrt{24}$  RR, in which cafe NE will be to ND as 5R to I; and fo on infinitely, the numbers 3, 8, 15, 24, &c. being gathered by continual addition of the terms of the arithmetical progression 3,5,7,9, Sc. The truth of all this Mathematicians will eafily examine. Vid. Whish. Thy. Math. Prol. P 231

Now it is to be observed, that as when the Sun comes to his Tropicks, days increase and decrease but a very little for a great while together ; fo when by increasing the diftance CD, these Angles come to their limits, they vary their quantity but very little for some time together, and therefore a far greater number of the rays which fall upon all the points N in the Quadrant BL, shall emerge in the limits of these Angles, then in any other inclinations. And further it is

to

to be observed, that the rays which differ in refrangibility will have different limits of their Angles of emergence, and by confequence according to their different degrees of refrangibility emerge. most copiously in different Angles, and being separated from one another appear each in their proper Colours. And what those Angles are may be easily gathered from the foregoing Theorem by computation.

For in the leaft refrangible rays the fines I and R (as was found above) are 108 and 81, and thence by computation the greateft Angle AXR will be found 42 degrees and 2 minutes, and the leaft Angle AYS, 50 degr. and 57 minutes. And in the most refrangible rays the fines I and R are 109 and 81, and thence by computation the greatest Angle AXR will be found 4.0 degrees and 17 minutes, and the least Angle AYS 54 degrees and 7 minutes.

Suppofe now that O is the Spectator's Eye, and OP a line Fig. 15. drawn parallel to the Sun's rays, and let POE, POF, POG, POH, be Angles of 40 degr. 17 min. 42 degr. 2 min. 50 degr. 57 min. and 54 degr. 7 min. refpectively, and these Angles turned about their common fide OP, fhall with their other fides OE, OF; OG, OH dedefcribe the verges of two Rain-bows AFBE and CHDG. For if E, F, G, H, be Drops placed any where in the conical fuperficies defcribed by OE, OF, OG, OH, and be illuminated by the Sun's rays SE, SF, SG, SH; the Angle SEO being equal to the Angle POE or 40 degr. 17 min. fhall be the greates Angle in which the most refrangible rays can after one reflexion be refracted to the Eye, and therefore all the Drops in the line OE fhall fend the most refrangible R rays rays most copiously to the Eye, and thereby strike the fenses with the deepest violet Colour in that region. And in like manner the Angle SFO being equal to the Angle POF, or 42 deg. 2 min. Stall be the greatest in which the least refrangible rays after one reflexion can emerge out of the Drops, and therefore those rays shall come most copiously to the Eye from the Drops inthe line OF, and strike the fenses with the deepest red Colour in that region. And by the fame argument, the rays which have intermediate degrees of refrangibility shall come most copiously from Drops between E and F, and strike the fenses with the intermediate Colours in the order which their degrees of refrangibility require, that is, in the progress from E to F, or from the infide of the Bow to the outfide in this order, violet, indico, blue, green, yellow, orange, red. But the violet, by the mixture of the white Light of the Clouds, will appear faint and incline to purple.

Again, the Angle SGO being equal to Angle POG, or 50 gr. 51 min. Ihall be the leaft Angle in which the leaft refrangible rays can after two reflexions emerge out of the Drops, and therefore the leaft refrangible rays Ihall come most copiously to the Eye from the Drops in the line OG, and ftrike the fense with the deepest red in that region. And the Angle SHO being equal to the Angle POH or 54 gr. 7 min. Ihall be the least Angle in which the most refrangible rays after two reflections can emerge out of the Drops, and therefore those rays Ihall come most copiously to the Eye from the Drops in the line OH, and ftrike the fenses with the deepest violet in that region. And by the fame argument, the Drops in the regions between G and H Ihall ftrike the fense with the

the intermediate Colours in the order which their degrees of refrangibility require, that is, in the progrefs from G to H, or from the infide of the Bow to the outfide in this order, red, orange, yellow, green, blue, indico, violet. And fince theie four lines OE, OF, OG. OH, may be fituated any where in the above-mentioned conical fuperficies, what is faid of the Drops and Colours in these lines is to be understood of the Drops and Colours every where in those fuperficies.

Thus shall there be made two Bows of Colours, an interior and ftronger, by one reflexion in the Drops, and an exterior and fainter by two; for the Light be-comes fainter by every reflexion. And their Colours fhall ly in a contrary order to one another, the red of both Bows bordering upon the fpace GF which is be-tween the Bows. The breadth of the interior Bow EOF measured cross the Colours shall be 1 degr. 45 min. and the breadth of the exterior GOH shall be 3 degr. 10 min. and the diftance between them GOF, shall be 8 gr. 55 min. the greatest Semi-diameter of the innermost, that is, the Angle POF being 42 gr. 2 min. and the least Semi-diameter of the outermost POG, being 50 gr. 57 min. Thefe are the meafures of the Bows, as they would be were the Sun but a point; for by the breadth of his Body the breadth of the Bows will be increased and their distance decreased by half a degree, and fo the breadth of the interior Iris will be 2 degr. 15 min. that of the exterior 3 degr. 40 min. their distance 8 degr. 25 min. the greatest Semi-diameter of the interior Bow 42 degr. 17 min. and the least of the exterior 50 degr. 42 min. And fuch are the dimensions of the Bows in the Heavens found to be very nearly, R 2 when

# [132]

when their Colours appear strong and perfect. Foronce, by fuch means as I then had, I meafured the greatest Semi-diameter of the interior Iris about 42 degrees, the breadth of the red, yellow and green in that Iris 63 or 64 minutes, befides the outmost faint red obscured by brightness of the Clouds, for which we may allow 3 or 4 minutes more. The breadth of the blue was about 40 minutes more befides the violet, which was fo much obscured by the brightness of the Clouds, that I could not measure its breadth. But fupposing the breadth of the blue and violet together to equal that of the red, yellow and green together, the whole breadth of this Iris will be about  $2\frac{1}{4}$  degrees as above. The least distance between this Iris and the exterior Iris was about 8 degrees and 30 minutes. The exterior Iris was broader than the interior, but so faint, especially on the blue fide, that I could not measure its breadth diftinctly. At another time when both Bows appeared more diftinct, I measured the breadth of the interior Iris 2 gr. 10', and the breadth of the red, yel-low and green in the exterior Iris, was to the breadth of the fame Colours in the interior as 3 to 2.

This Explication of the Rain-bow is yet further confirmed by the known Experiment (made by Antonius de Dominis and Des-Cartes) of hanging up any where in the Sun-fhine a Glass-Globe filled with Water, and viewing it in fuch a pofture that the rays which come from the Globe to the Eye may contain with the Sun's rays an Angle of either 42 or 50 degrees. For if the Angle be about 42 or 43 degrees, the Spectator (fuppofe at O) fhall fee a full red Colour in that fide of the Globe oppofed to the Sun as 'tis reprefented at F, and

if

if that Angle become lefs (fuppofe by depreffing the Globe to E) there will appear other Colours, yellow, green and blue fucceffively in the fame fide of the Globe. But if the Angle be made about 50 degrees (fuppofe by lifting up the Globe to G) there will appear a red Colour in that fide of the Globe towards the Sun, and if the Angle be made greater (fuppofe by lifting up the Globe to H) the red will turn fucceffively to the other Colours yellow, green and blue. The fame thing I have tried by letting a Globe reft, and raifing or depreffing the Eye, or otherwife moving it to make the Angle of a juft magnitude.

I have heard it reprefented, that if the Light of a Candle be refracted by a Prism to the Eye; when the blue Colour falls upon the Eye the Spectator shall fee red in the Prifm, and when the red falls upon the Eye he shall see blue; and if this were certain, the Colours of the Globe and Rain-bow ought to appear in a contrary order to what we find. But the Colours of the Candle being very faint, the mistake seems to arise from the difficulty of difcerning what Colours fall on the Eye. For, on the contrary, I have fometimes had occafion to obferve in the Sun's Light refracted by a Prifm, that the Spectator always fees that Colour in the Prifm. which falls upon his Eye. And the fame I have found true alfo in Candle-Light. For when the Prifm is moved flowly from the line which is drawn directly from the Candle to the Eye, the red appears first in the Prism and then the blue, and therefore each of them is feen when it falls upon the Eye. For the red paffes over the Eye first, and then the blue.

The

The Light which comes through Drops of Rain by two refractions without any reflexion, ought to appear ftrongest at the distance of about 26 degrees from the Sun, and to decay gradually both ways as the diftance from him increases and decreases. And the same is to be understood of Light transmitted through spherical Hail-stones. And if the Hail be a little flatted, as it often is, the Light transmitted may grow io ftrong at a little less distance than that of 26 degrees, as to form a Halo about the Sun or Moon; which Halo, as often as the Hail-ftones are duly figured may be coloured, and then it must be red within by the least refrangible rays, and blue without by the most refrangible ones, especially if the Hail-ftones have opake Globules of Snow in their center to intercept the Light within the Halo (as Hugenius has observed) and make the infide thereof more diffinctly defined than it would otherwife be. For fuch Hail-stones, though spherical, by terminating the Light by the Snow, may make a Halo red within and colourlefs without, and darker in the red than without, as Halos use to be. For of those rays which pass close by the Snow the rubriform will be least refracted, and fo come to the Eye in the directeft lines.

The Light which paffes through a Drop of rain after two refractions, and three or more reflexions, is fearce ftrong enough to caufe a fenfible Bow ; but in those Cylinders of Ice by which *Hugenius* explains the *Parbelia*, it may perhaps be fenfible.

PROP.

# [135]

# PROP. X. PROB. V.

# By the discovered Properties of Light to explain the permanent Colours of natural Bodies.

Thefe Colours arife from hence, that fome natural Bodies reflect fome forts of rays, others other forts more copioufly than the reft. Minium reflects the leaft refrangible or red-making rays most copioufly, and thence appears red. Violets reflect the most refrangible, most copioufly, and thence have their Colour, and to of other Bodies. Every Body reflects the rays of its own Colour more copioufly than the reft, and from their excels and predominance in the reflected Light has its Colour.

#### EXPER. XVII.

For if the homogeneal Lights obtained by the folution of the Problem proposed in the 4th Proposition of the first Book you place Bodies of feveral Colours, you will find, as I have done, that every Body looks most splendid and luminous in the Light of its own Colour. Cinnaber in the homogeneal red Light is most resplendent, in the green Light it is manifestly less resplendent, and in the blue Light still less. Indico in the violet blue Light is most resplendent, and its splendor is gradually diminissed as it is removed thence by degrees through the green and yellow Light to the red. By a Leek the green Light, and next that the blue and yellow which compound green, are more strongly reflected

flected than the other Colours red and violet, and fo of the reft. But to make these Experiments the more manifest, fuch Bodies ought to be chosen as have the fullest and most vivid Colours, and two of those Bodies are to be compared together. Thus, for inftance, if Cinnaber Camleum Cyprium and ultra marine blue, or fome other full blue be ved ultra mannum held together in the homogeneal Light, they will both appear red, but the Cinnaber will appear of a ftrongly luminous and refplendent red, and the ultra marine blue of a faint obscure and dark red; and if they be held together in the blue homogeneal Light they will both appear blue, but the ultra marine will appear of a ftrongly luminous and resplendent blue, and the Cinnaber of a faint and dark blue? Which puts it out of difpute, that the Cinnaber reflects the red Light much more copiously than the ultra marine doth, and the ultra marine reflects the blue Light much more copioufly than the Cinnaber doth. The fame Experiment may be tryed fuccesfully with red Lead and Indico, or with any other two coloured Bodies, if due allowance be made for the different ftrength or weakness of their Colour and Light.

And as the reafon of the Colours of natural Bodies is evident by thefe Experimenrs, to it is further confirmed and put paft difpute by the two firft Experiments of the firft Book, whereby 'twas proved in fuch Bodies that the reflected Light which differ in Colours do differ alfo in degrees of refrangibility. For thence it's certain, that fome Bodies reflect the more refrangible, others the lefs refrangible rays more copioufly.

And

And that this is not only a true reafon of these Colours, but even the only reason may appear further from this confideration, that the Colour of homogeneal Light cannot be changed by the reflexion of natural Bodies.

For if Bodies by reflexion cannot in the leaft change the Colour of any one fort of rays, they cannot appear coloured by any other means than by reflecting those which either are of their own Colour, or which by mixture must produce it.

But in trying Experiments of this kind care must be had that the Light be fufficiently homogeneal. For if Bodies be illuminated by the ordinary prifmatick Colours, they will appear neither of their own day-light Colours, nor of the Colour of the Light caft on them, but of 10me middle Colour between both, as I have found by Experience. Thus red Lead (for inftance) illuminated with the ordinary prifmatick green will not appear either red or green, but orange or yellow, or between yellow and green accordingly, as the green Light by which 'tis' illuminated is more or lefs compounded. For becaufe red Lead appears red when il-luminated with white Light, wherein all forts of rays are equally mixed, and in the green Light all forts of rays are not equally mixed, the excess of the yellowmaking, green-making and blue-making rays in the incident green Light, will cause those rays to abound fo much in the reflected Light as to draw the Colour from red towards their Colour. And becaufe the red Lead reflects the red-making rays most copiously in proportion to their number, and next after them the orange-making and yellow-making rays; thefe rays in the

[138]

the reflected Light will be more in proportion to the Light than they were in the incident green Light, and thereby will draw the reflected Light from green towards their Colour. And therefore the red Lead will appear neither red nor green, but of a Colour between both. In transparently coloured Liquors 'tis observable, 97 that their Colour uses to vary with their thickness. Thus, for instance, a red Liquor in a conical Glass held between the Light and the Eye, looks of a pale and dilute yellow at the bottom where 'tis thin, and a little higher where 'tis thicker grows orange, and where 'tis still thicker becomes red, and where 'tis thickess the red is deepest and darkest. For it is to be conceived that fuch a Liquor stops the indico-making and violetmaking rays most easily, the blue making rays more difficultly, the green-making rays still more difficultly, and the red-making most difficultly : And that if the thickness of the Liquor be only so much as suffices to ftop a competent number of the violet-making and indico-making rays, without diminishing much the number of the reft, the reft must (by Prop. 6. Lib. 2.) compound a pale yellow. But if the Liquor be so much thicker as to ftop also a great number of the blue-making rays, and fome of the green-making, the reft muft compound an orange; and where it is fo thick as to ftop alfo a great number of the green-making and a confiderable number of the yellow-making, the reft must begin to compound a red, and this red must grow deeper and darker as the yellow making and orange-making rays are more and more ftopt by increasing the thicknels of the Liquor, fo that few rays befides the redmaking can get through.

Of

Of this kind is an Experiment lately related to me by Mr. Halley, who, in diving deep into the Sea, found in a clear Sun-fhine day, that when he was funk many Fathoms deep into the Water, the upper part of his Hand in which the Sun shone directly through the Water looked of a red Colour, and the under part of his Hand illuminated by Light reflected from the Water below looked green. For thence it may be gathered, that the Sea-water reflects back the violet and bluemaking rays most easily, and lets the red-making rays pass most freely and copiously to great depths. For thereby the Sun's direct Light at all great depths, by reation of the predominating red-making rays, must appear red; and the greater the depth is, the fuller and intenfer must that red be. And at fuch depths as the violet-making rays fcarce penetrate unto, the bluemaking, green-making and yellow-making rays being reflected from below more copioufly than the red-making ones, must compound a green.

Now if there be two Liquors of full Colours, fuppofe a red and a blue, and both of them fo thick as fuffices to make their Colours fufficiently full; though either Liquor be fufficiently transparent apart, yet will you not be able to see through both together. For if only the red-making rays pass through one Liquor, and only the blue-making through the other, no rays can pass through both. This Mr. Hook tried cafually with Glass-wedges filled with red and blue Liquors, and was surprized at the unexpected event, the reason of it being then unknown; which makes me trust the more to his Experiment, though I have not tryed it my self. But he that would repeat it, must take care the Liquors be of very good and full Colours.

Now

# [140]

Now whilft Bodies become coloured by reflecting or transmitting this or that fort of rays more copioully than the reft, it is to be conceived that they ftop and fifle in themselves the rays which they do not reflect or transmit. For if Gold be foliated and held between your Eye and the Light, the Light looks blue, and therefore maffy Gold lets into its Body the blue-making rays to be reflected to and fro within it till they be ftopt and ftifled, whilft it reflects the yellow-making outwards, and thereby looks yellow. And much after the fame manner that Leaf-gold is yellow by reflected, and blue by transmit-ted Light, and maffy Gold is yellow in all positions of the Eye; there are some Liquors as the tincture of Lignum Nephriticum, and some forts of Glass which transmit one fort of Light most copiously, and reflect another fort, and thereby look of feveral Colours, according to the polition of the Eye to the Light. But if these Liquors or Glasses were to thick and massy that no Light could get through them, I question not but that they would like all other opake Bodies appear of one and the same Colour in all positions of the Eye, though this I cannot yet affirm by experience. For all coloured Bodies, so far as my Observation reaches, may be feen through if made fufficiently thin, and therefore are in some measure transparent, and differ only in degrees of transparency from tinged transparent Liquors; these Liquors, as well as those Bodies, by a sufficient thickness becoming opake. A transparent Body which looks of any Colour by transmitted Light, may also look of the fame Colour by reflected Light, the Light of that Colour being reflected by the further furface of the Body, or by the Air beyond it. And then the reflected Colour will be diminished, and perhaps cease, by making

making the Body very thick, and pitching it on the back-fide to diminifh the reflexion of its further furface, fo that the Light reflected from the tinging particles may predominate. In fuch cafes, the Colour of the reflected Light will be apt to vary from that of the Light transmitted. But whence it is that tinged Bodies and Liquors reflect fome fort of rays, and intromit or transmit other forts, shall be faid in the next Book. In this Proposition I content my felf to have put it past dispute, that Bodies have such Properties, and thence appear coloured.

# PROP. XI. PROB. VI.

By mixing coloured Lights to compound a Beam of Light of the Jame Colour and Nature with a Beam of the Sun'sdirect Light, and therein to experience the truth of the foregoing Propositions.

Let A B Ca bc reprefent a Prifm by which the Sun's Fig. 16: Light let into a dark Chamber through the Hole F, may, be refracted towards the Lens M N, and paint upon it at p, q, r, s and t, the ufual Colours violet, blue, green, yellow and red, and let the diverging rays by the refraction of this Lens converge again towards X, and there, by the mixture of all those their Colours, compound, a white according to what was shewn above. Then let another Prism DEGdeg, parallel to the former, be placed at X, to refract that white Light upwards towards Y. Let the refracting Angles of the Prisms, and their distances from the Lens be equal, fo that the rays which converged from the Lens towards X, and without refraction, would there have crossed and diverged again, may by the refraction of the fecond Prism be reduced.

# [142]

reduced into Parallelism and diverge no more. For then those rays will recompose a Beam of white Light XY. If the refracting Angle of either Prism be the bigger, that Prism must be so much the nearer to the Lens. You will know when the Prisms and the Lens are well fet together by obferving if the Beam of Light X Y which comes out of the fecond Prifm be perfectly white to the very edges of the Light, and at all diftan-ces from the Prilm continue perfectly and totally white like a Beam of the Sun's Light. For till this happens, the polition of the Prisms and Lens to one another must be corrected, and then if by the help of a long Beam of Wood, as is represented in the Figure, or by a Tube, or some other such instrument made for that purpose, they be made fast in that situation, you may try all the fame Experiments in this compounded Beam of Light-XY, which in the foregoing Experiments have been made in the Sun's direct Light. For this compounded Beam of Light has the fame appearance, and is endowed with all the fame Properties with a direct Beam of the Sun's Light, fo far as my Obfervation reaches. And in trying Experiments in this Beam you may by ftopping any of the Colours p, q, r, s and t, at the Lens, fee how the Colours produced in the Experiments are no other than those which the rays had at the Lens before they entered the composition of this Beam : And by confequence that they arife not from any new modifications of the Light by refractions and reflexions, but from the various separations and mixtures of the rays originally endowed with their colour-making qualities.

So, for inftance, having with a Lens  $4\frac{1}{4}$  lnches broad, and two Prifms on either Hand  $6\frac{1}{4}$  Feet diftant from the Lens, made fuch a Beam of compounded Light : to examin

examin the reason of the Colours made by Prisins, I refracted this compounded Beam of Light XY with another Prifm HIKkh, and thereby caft the ufual prifmatick Colours PQRST upon the Paper LV placed behind. And then by stopping any of the Colours p, q, r, s, t, at the Lens, I found that the same Colour would vanish at the Paper. So if the purple P was stopped at the Lens, the purple P upon the Paper would vanish, and the reft of the Colours would remain unaltered, unless perhaps the blue, so far as some purple latent in it at the Lens might be separated from it by the following refractions. And fo by intercepting the greenupon the Lens, the green R upon the Paper would va-nish, and so of the rest; which plainly shews, that as the white Beam of Light XY was compounded of feve Lights varioufly coloured at the Lens, fo the Co-lours which afterwards emerge out of it by new refractions are no other than those of which its whiteness was compounded. The refraction of the Prifin HIK kh generates the Colours PQRST upon the Paper, not by changing the colorific qualities of the rays, but by feparating the rays which had the very fame colorific qualities before they entered the composition of the refracted Beam white of Light XY. For otherwise the rays. which were of one Colour at the Lens might be of another upon the Paper, contrary to what we find.

So again, to examin the reafon of the Colours of natural Bodies, I placed fuch Bodies in the Beam of Light X Y, and found that they all appeared there of those their own Colours which they have in Day-light, and that those Colours depend upon the rays which had the fame Colours at the Lens before they entred the compofition.

fition of that Beam. Thus, for instance, Cinnaber illuminated by this Beam appears of the fame red Colour as in Day-light; and if at the Lens you intercept the green-making and blue-making rays, its rednefs will become more full and lively : But if you there intercept the redmaking rays, it will not any longer appear red, but become yellow or green, or of fome other Colour, according to the forts of rays which you do not intercept. So Gold in this Light XY appears of the same yellow Colour as in Day-light, but by intercepting at the Lens a due quantity of the yellow-making rays it will appear white like Silver (as I have tryed) which shews that its yellowness arifes from the excess of the intercepted rays tinging that whiteness with their Colour when they are let pass. So the infusion of Lignum Nephriticum (as I have also tryed) when held in this Beam of Light XY, looks blue by the reflected part of the Light, and yellow by the transmitted part of it, as when 'tis viewed in Daylight, but if you intercept the blue at the Lens the infufion will lose its reflected blue Colour, whilft its transmitted red remains perfect and by the lois of some bluemaking rays wherewith it was allayed becomes more intenfe and full. And, on the contrary, if the red and orangemaking rays be intercepted at Lens, the infufion will lole its transmitted red, whilst its blue will remain and become more full and perfect. Which shews, that the infusion does not tinge the rays with blue and yellow, but only transmit those most copiously which were red-making before, and reflects those most copiously which were blue-making before. And after the fame manner may the reafons of other Phænomena be examined, by trying them in this artificial Beam of Light XY.

THE





Book I. Part II. Plate II.









Book I. Part II. Plate N.





# THE SECOND BOOK OF OPTICKS. PART I.

# Observations concerning the Reflexions, Refractions, and Colours of thin transparent Bodies.

I has been obferved by others that transparent Subfances, as Glafs, Water, Air, &c. when made very thin by being blown into Bubbles, or otherwife formed into Plates, do exhibit various Colours according to their various thinnefs, although at a greater thicknefs they appear very clear and colourlefs. In the former Book I forbore to treat of these Colours, because they seemed of a more difficult confideration, and were not neceffary for establishing the Properties of Light there discoursed of. But because they may conduce to further discoveries for completing the Theory of Light, especially as to the constitution of the parts of natural Bodies, on which their Colours or Transparency depend; I have here set down an account of them. To render this Discourse short and distinct, I have first described the principal of my A a ObserObservations, and then confidered and made use of them. The Observations are these.

## OBS. I.

Compressing two Prisms hard together that their Sides (which by chance were a very little convex)might fomewhere touch one another: I found the place in which they touched to become absolutely transparent, as if they had there been one continued piece of Glass. For when the Light fell fo obliquely on the Air, which in other places was between them, as to be all reflected ; it feemed in that place of contact to be wholly tranfmitted, infomuch that when looked upon, it appeared like a black or dark Spot, by reason that little or no sensible Light was reflected from thence, as from other places; and when looked through it feemed (as it were) a hole in that Air which was formed into a thin Plate, by being compressed between the Glasses. And through this hole Objects that were beyond might be feen diftinctly, which could not at all be feen through other parts of the Glaffes where the Air was interjacent. Although the Glaffes were a little convex, yet this tranfparent Spot was of a confiderable breadth, which breadth feemed principally to proceed from the yielding inwards of the parts of the Glasses, by reason of their mutual pressure. For by pressing them very hard together it would become much broader than otherwife.

OBS.

[3]

# OBS. II.

When the Plate of Air, by turning the Prisms about their common Axis, became fo little inclined to the incident Rays, that fome of them began to be transmitted, there arose in it many slender Arcs of Colours which at first were shaped almost like the Conchoid, as you see them delineated in the first Figure. And Fig. 1. by continuing the motion of the Prisms, these Arcs increafed and bended more and more about the faid tranf--parent Spot, till they were completed into Circles or Rings incompaffing it, and afterwards continually grew more and more contracted.

These Arcs at their first appearance were of a violet and blue Colour, and between them were white Arcs of Circles, which prefently by continuing the motion of the Prisms became a little tinged in their inward Limbs with red and yellow, and to their outward Limbs the blue was adjacent. So that the order of these Colours from the central dark Spot, was at that time white, blue, violet ; black ; red, orange, yellow, white, blue, violet, &c. But the yellow and red were much fainter than the blue and violet.

The motion of the Prisms about their Axis being continued, these Colours contracted more and more, shrinking towards the whiteness on either fide of it, until they totally vanished into it. And then the Circles in those parts appeared black and white, without any other Colours intermixed. But by further moving the Prifms about, the Colours again emerged out of the whitenefs, the violet and blue as its inward Limb, and at its outward

#### OBS. III.

When the Rings or fome parts of them appeared only black and white, they were very diffinct and well defined, and the backnefs feemed as intenfe as that of the central Spot. Alfo in the borders of the Rings, where the Colours began to emerge out of the whitenefs, they were pretty diffinct, which made them vifible to a very great Multitude. I have fometimes numbred above thirty Succeffions (reckoning every black and white Ring for one Succeffion) and feen more of them, which by reafon of their fmalnefs I could not number. But in other Politions of the Prifms, at which the Rings appeared of many Colours, I could not diffinguifh above eight or nine of them, and the exterior of thofe were very confufed and dilute.

In these two Observations to see the Rings diftinct, and without any other Colour than black and white, I found it neceffary to hold my Eye at a good distance from them. For by approaching nearer, although in the same inclination of my Eye to the plane of the Rings, there emerged a blueiss Colour out of the white, which by dilating it felf more and more into the black rendred the Circles less distinct, and less the white a little tinged with red and yellow. I found also by looking through a flit or oblong hole, which was narrower than the Pupil of my Eye, and held close to it it parallel to the Prifms, I could fee the Circles much diftincter and visible to a far greater number than otherwife.

#### OBS. IV.

To observe more nicely by the order of the Colours which arofe out of the white Circles as the Rays became lefs and lefs inclined to the plate of Air; I took two Object Glaffes, the one a Plano-convex for a fourteen-foot Telescope, and the other a large double convex for one of about fifty-foot; and upon this laying the other with its its plane-fide downwards, I preffed them flowly together, to make the Colours fucceffively emerge in the middle of the Circles, and then flowly lifted the upper Glass from the lower to make them succef-fively vanish again in the same place. The Colour, which by preffing the Glasses together emerged last in the middle of the other Colours, would upon its first appearance look like a Circle of a Colour almost uniform from the circumference to the center, and by compreffing the Glaffes still more, grow continually broader until a new Colour emerged in its center, and thereby it became a Ring encompassing that new Colour. And by compreffing the Glaffes still more, the Diameter of this Ring would encrease, and the breadth of its Orbit or Perimeter decrease until another new Colour emerged in the center of the last : And fo on until a third, a fourth, a fifth, and other following new Colours fucceffively emerged there, and became Rings encompaffing the innermost Colour, the last of which was the black Spot. And, on the contrary, by lifting

lifting up the upper Glafs from the lower, the diameter of the Rings would decreafe, and the breadth of their Orbit encreafe, until their Colours reached fucceffively to the center ; and then they being of a confiderable breadth, I could more eafily difcern and diftinguifh their Species than before. And by this means I obferved their Succeffion and Quantity to be as followeth.

Next, to the pellucid central Spot made by the contact of the Glasses fucceeded blue, white, yellow, and red, the blue was fo little in quantity that I could not difcern it in the circles made by the Prisms, nor could I well diftinguish any violet in it, but the yellow and red were pretty copious, and seemed about as much in extent as the white, and four or five times more than the blue. The next Circuit in order of Colours immediately encompaffing these were violet, blue, green, yellow, and red, and these were all of them copious and vivid, excepting the green, which was very little in quantity, and feemed much more faint and dilute than the other Colours. Of the other four, the violet was the leaft in extent, and the blue lefs than the yellow or red. The third Circuit or Order was purple, blue, green, yellow, and red ; in which the purple feemed more reddifh than the violet in the former Circuit, and the green was much more confpicuous, being as brifque and copious as any of the other Colours, except the yellow; but the red began to be a little faded, inclining very much to purple. After this fucceeded the fourth Circuit of green and red. The green was very copious and lively, inclining on the one fide to blue, and on the other fide to yellow. But in this
this fourth Circuit there was neither violet, blue, nor yellow, and the red was very imperfect and dirty. Alfo the fucceeding Colours became more and more imperfect and dilute, till after three or four Revolutions they ended in perfect whitenefs. Their Form, when the Glaffes were most compressed fo as to make the black Spot appear in the Center, is delineated in the Second Figure; where a, b, c, d, e: f, g, b, i, k: l, m, n, o, p:q, r: Fig. 2.s, t: v, x: y denote the Colours reck'ned in order from the center, black, blue, white, yellow, red : violet, blue, green, yellow, red : purple, blue, green, yellow, red : green, red : greenish blue, red : greenish blue, pale red : greenish blue, 
#### OBS. V.

To determine the interval of the Glaffes, or thicknefs of the interjacent Air, by which each Colour was produced, I meafured the Diameters of the first fix Rings at the most lucid part of their Orbits, and squaring them, I found their Squares to be in the Arithmetical Progression of the odd Numbers, 1.3.5.7.9.11. And fince one of these Glaffes was Plain, and the other Spherical, their Intervals at those Rings must be in the fame Progression. I measured also the Diameters of the dark or faint Rings between the more lucid Colours, and found their Squares to be in the Arithmetical Progression of the even Numbers, 2.4.6.8.10.12. And it being very nice and difficult to take these measures exactly; I repeated them at divers times at divers parts of the Glaffes, that by their Agreement I might be confirmed in them. And the fame Method I used in deterdetermining fome others of the following Obfervations.

#### OBS. VI.

The Diameter of the fixth Ring at the most lucid part of its Orbit was 100 parts of an Inch, and the Diameter of the Sphere on which the double convex Object-Glass was ground was about 102 Feet, and hence I gathered the thickness of the Air or Aereal Interval of the Glasses at that Ring. But some time after, sufpecting that in making this Observation I had not determined the Diameter of the Sphere with sufficient accurateness, and being uncertain whether the Planoconvex Glass was truly plain, and not something concave or convex on that fide which I accounted plain; and whether I had not preffed the Glaffes together, as I often did, to make them touch. (for by preffing fuch Glaffes together their parts eafily yield inwards, and the Rings thereby become fenfibly broader than they would be, did the Glaffes keep their Figures.) I repeated the Experiment, and found the Diameter of the fixth lucid Ring about 15 parts of an Inch. I repeated the Experiment alfo with fuch an Object-Glafs of another Telescope as I had at hand. This was a double convex ground on both fides to one and the fame Sphere, and its Focus was diftant from it 83<sup>2</sup> Inches. And thence, if the Sines of incidence and refraction of the bright yellow Light be affumed in proportion as 11 to 17, the Diameter of the Sphere to which the Glass was figured will by computation be found 182 Inches. This Glafs I laid upon a flat one, fo that the black

black Spot appeared in the middle of the Rings of Colours without any other preffure than that of the weight of the Glafs. And now measuring the Diameter of the fifth dark Circle as accurately as I could, I found it the fifth part of an Inch precifely. This measure was taken with the points of a pair of Compasses on the upper furface on the upper Glais, and my Eye was about eight or nine Inches diftance from the Glais, almost perpendicularly over it, and the Glass was a of an Inch thick, and thence it is eafy to collect that the true Diameter of the Ring between the Glasses was greater than its measured Diameter above the Glasses in the proportion of 80 to 79 or thereabouts, and by confequence equal to  $\frac{16}{79}$  parts of an Inch, and its true Semi-diameter equal to <sup>8</sup>/<sub>79</sub> parts. Now as the Diameter of the Sphere (182 Inches) is to the Semi-diameter of this fifth dark Ring  $\left(\frac{8}{79} \text{ parts of an Inch}\right)$  fo is this Semi-diameter to the thicknefs of the Air at this fifth dark Ring; which is therefore  $\frac{32}{5(7931)}$  or  $\frac{10}{1774534}$  parts of an Inch, and the fifth part thereof; viz. the  $\frac{1}{88727}$ th part of an Inch, is the thickness of the Air at the first of these dark Rings.

The fame Experiment I repeated with another double convex Object-glafs ground on both fides to one and the fame Sphere. Its Focus was diftant from it 168<sup>1</sup>/<sub>2</sub> Inches, and therefore the Diameter of that Sphere was 184 Inches. This Glafs being laid upon the fame plain Glafs, the Diameter of the fifth of the dark Rings, when the black Spot in their center appeared plainly without preffing the Glaffes, was by the meafure of the Compafies upon the upper Glafs  $\frac{11}{600}$  parts of an Inch, and by confequence between the Glaffes it was  $\frac{1122}{6000}$ . For the upper Glafs was  $\frac{1}{8}$  of an Inch thick, B b and my Eye was diftant from it 8 Inches. And a third proportional to half this from the Diameter of the Sphere is  $\frac{1}{888570}$  parts of an Inch. This is therefore the thickness of the Air at this Ring, and a fifth part thereof, viz. the  $\frac{1}{888570}$ th part of an Inch is the thickness thereof at the first of the Rings as above.

I tryed the fame thing by laying these Object-Glaffes upon flat pieces of a broken Looking-glass, and found the fame measures of the Rings : Which makes me rely upon them till they can be determined more accurately by Glaffes ground to larger Spheres, though in such Glaffes greater care must be taken of a true plain.

These Dimensions were taken when my Eye was placed almost perpendicularly over the Glasses, being about an Inch, or an Inch and a quarter, distant from the incident rays, and eight Inches diftant from the Glafs; fo that the rays were inclined to the Glafs in an Angle of about 4 degrees. Whence by the following, Observation you will understand, that had the rays been perpendicular to the Glaffes, the thickness of the Air at these Rings would have been less in the proportion of the Radius to the fecant of 4 degrees, that is of 10000. Let the thickneffes found be therefore diminifhed in this proportion, and they will become  $\frac{1}{83940}$  and  $\frac{\tau}{89003}$ , or ( to use the nearest round number ) the  $\frac{\tau}{89000}$ th part of an Inch. This is the thickness of the Air at the darkest part of the first dark Ring made by perpendicular rays, and half this thickness multiplied by the progreffion, 1, 3, 5, 7, 9, 11, & c. gives the thickneffes of the Air at the most luminous parts of all the brightest Rings, viz.  $\frac{1}{178000}$ ,  $\frac{3}{178000}$ ,  $\frac{5}{178000}$ ,  $\frac{7}{178000}$ ,  $\Im c$ . their arithmetical means

### [ II ]

means  $\frac{2}{178000}$ ,  $\frac{4}{178000}$ ,  $\frac{6}{178000}$ ,  $\Im c$ . being its thickneffes at the darkeft parts of all the dark ones.

#### OBS. VII.

The Rings were leaft when my Eye was placed perpendicularly over the Glaffes in the Axis of the Rings: And when I viewed them obliquely they became bigger, continually fwelling as I removed my Eye further from the Axis. And partly by meafuring the Diameter of the fame Circle at feveral obliquities of my Eye, partly by other means, as alfo by making ufe of the two Prifms for very great obliquities. I found its Diameter, and confequently the thicknefs of the Air at its perimeter in all those obliquities to be very nearly in the proportions expressed in this Table.

	Angle idence	of In-	Angle of fraction	of Re-	Diameter of the Ring.	Thickness of the Air.
1	Air.		the Air.	•		
-	deg.	min.	-			
	.00	00	00	00	IO	IO
	06	.26	10	00	$IO_{13}^{T}$	$I \bigcirc \frac{2}{13}$
	12	45	20	00	$IO_{\overline{3}}^{T}$	$IO_{\overline{3}}^{2}$
	18	49	30	00	$IO_{\frac{2}{4}}$	$II\frac{\tilde{I}}{2}$
	24	30	40	00	$II_{\frac{2}{3}}$	13
	29	37	50	00	$I2\frac{i}{2}$	$I 5^{\frac{1}{2}}$
	33	58	60	00	14	20
	35	47	65	00	$15\frac{1}{4}$	$23\frac{1}{4}$
	37	19	70	00	164	284
	. 38	33	75	00	$19^{\frac{1}{4}}$	37
	39	27	80	00	$22\frac{6}{7}$	$52\frac{1}{4}$
	.40	00	85	00	29	84-
	40	II	90	00	35	1221

**BP 3** 

In

In the two first Columns are expressed the obliquities of the incident and emergent rays to the plate of the Air, that is, their angles of incidence and refraction. In the third Column the Diameter of any coloured Ring at those obliquities is expressed in parts, of which ten constitute that Diameter when the rays are perpendicular. And in the fourth Column the thickness of the Air at the circumference of that Ring is expressed in parts of which also ten constitute that thickness when the rays are perpendicular.

And from thefe meafures I feem to gather this Rule : That the thicknefs of the Air is proportional to the fecant of an angle, whofe Sine is a certain mean proportional between the Sines of incidence and refraction. And that mean proportional, fo far as by thefe meafures I can determine it, is the first of an hundred and fix arithmetical mean proportionals between those Sines counted from the Sine of refraction when the refraction is made out of the Glass into the plate of Air, or from the Sine of incidence when the refraction ismade out of the plate of Air into the Glass.

#### O.B.S. VIII.

The dark Spot in the middle of the Rings increased alfo by the obliquation of the Eye, although almost infensibly. But if instead of the Object-Glasses the Prisms were made use of, its increase was more manifest when viewed so obliquely that no Colours appeared about it. It was least when the rays were incident most obliquely on the interjacent Air, and as the obliquity decreased it increased more and more until the coloured Rings appeared,

peared, and then decreafed again, but not fo much as it increased before. And hence it is evident, that the transparency was not only at the absolute contact of the Glasses, but also where they had some little interval. I have fometimes observed the Diameter of that Spot to be between half and two fifth parts of the Diameter of the exterior circumference of the red in the first circuit or revolution of Colours when viewed almost perpendicularly; whereas when viewed obliquely it hath wholly vanished and become opake and white like the other parts of the Glass; whence it may be collected that the Glasses did then scarcely, or not at all, touch one another, and that their interval at the perimeter of that Spot when viewed perpendicularly was about a fifth or fixth part of their interval at the circumference of the faid red.

#### OBS. IX.

By looking through the two contiguous Object-Glaffes, I found that the interjacent Air exhibited Rings of Colours, as well by transmitting Light as by reflecting it. The central Spot was now white, and from it the order of the Colours were yellowish red; black; violet, blue, white, yellow, red; violet, blue, green, yellow, red,  $\Im c$ . But these Colours were very faint and dilute unless when the Light was trajected very obliquely through the Glasses: For by that means they became pretty vivid. Only the first yellowish red, like the blue in the fourth Observation, was so little and faint as fearcely to be differented. Comparing the coloured Rings made by reflexion, with these made by transtranfmiffion of the Light; I found that white was oppofite to black, red to blue, yellow to violet, and green to a compound of red and violet. That is, thofe parts of the Glafs were black when looked through, which when looked upon appeared white, and on the contrary. And fo thofe which in one cafe exhibited blue, did in the other cafe exhibit red. And the like of the other Colours. The manner you have reprefented in the third Figure, where AB, CD, are the furfaces of the Glaffes contiguous at E, and the black lines between them are their diffances in arithmetical progreffion, and the Colours written above are feen by reflected Light, and thofe below by Light tranfmitted.

#### OBS. X.

Wetting the Object-Glasses a sittle at their edges, the water crept in flowly between them, and the Circles thereby became lefs and the Colours more faint : Infomuch that as the water crept along one half of them at which it first arrived would appear broken off from the other half, and contracted into a lefs room. By measuring them I found the proportions of their Diameters to the Diameters of the like Circles made by Air to be about feven to eight, and confequently the intervals of the Glaffes at like Circles, caufed by those two mediums Water and Air, are as about three to four. Perhaps it may be a general Rule, That if any other medium more or lefs denfe than water be compreffed. between the Glaffes, their intervals at the Rings caufed thereby will be to their intervals caufed by interjacent Air.

Fig. 3.

[15]

Air, as the Sines are which measure the refraction made out of that medium into Air.

## OBS. XI.

When the water was between the Glaffes, if I preffed the upper Glafs varioufly at its edges to make the Rings move nimbly from one place to another, a little white Spot would immediately follow the center of them, which upon creeping in of the ambient water into that place would prefently vanifh. Its appearance was fuch as interjacent Air would have caufed, and it exhibited the fame Colours. But it was not Air, for where any bubbles of Air were in the water they would not vanifh. The reflexion must have rather been caufed by a fubtiler medium, which could recede through the. Glaffes at the creeping in of the water.

# OBS. XII.

These Observations were made in the open Air. But further to examin the effects of coloured Light falling on the Glasses, I darkened the Room, and viewed them by reflexion of the Colours of a Prism cast on a Sheet of white Paper, my Eye being so placed that I could fee the coloured Paper by reflexion in the Glasses, as in a Looking-glass. And by this means the Rings became distincter and visible to a far greater number than in the open Air. I have sometimes seen more than twenty of them, whereas in the open Air I could not. discern above eight or nine.

O'B.S.

[16]

#### OBS. XIII.

Appointing an affiftant to move the Prism to and fro about its Axis, that all the Colours might fucceffively fall on that part of the Paper which I faw by reflexion from that part of the Glaffes, where the Circles appeared, fo that all the Colours might be fucceffively reflected from the Circles to my Eye whilft I held it immovable, I found the Circles which the red Light made to be manifeftly bigger than those which were made by the blue and violet. And it was very pleafant to fee them gradually fwell or contract according as the Colour of the Light was changed. The interval of the Glasses at any of the Rings when they were made by the utmost red Light, was to their interval at the fame Ring when made by the utmost violet, greater than as 3 to 2, and lefs than as 13 to 8, by the most of my Observations it was as 14 to 9. And this proportion feemed very nearly the fame in all obliquities of my Eye; unleis when two Prisms were made use of instead of the Object-Glasses. For then at a certain great obliquity of my Eye, the Rings made by the feveral Colours feemed equal, and at a greater obliquity those made by the violet would be greater than the fame Rings made by the red. The refraction of the Prism in this cafe caufing the most refrangible rays to fall more obliquely on that plate of the Air than the least refrangible ones. Thus the Experiment fucceeded in the coloured Light, which was fufficiently ftrong and copious to make the Rings fenfible. And thence it may be gathered, that if the most refrangible and least refran[17] .

refrangible rays had been copious enough to make the Rings fenfible without the mixture of other rays, the proportion which here was 14 to 9 would have been a little greater, fuppofe  $14\frac{1}{4}$  or  $14\frac{1}{5}$  to 9.

#### OBS. XIV.

Whilft the Prism was turn'd about its Axis with an uniform motion, to make all the feveral Colours fall fucceffively upon the Object-Glaffes, and thereby to make the Rings contract and dilate : The contraction or dilation of each Ring thus made by the variation of its Colour was fwifteft in the red, and floweft in the violet, and in the intermediate Colours it had intermediate degrees of celerity. Comparing the quantity of contraction and dilation made by all the degrees of each Colour, I found that it was greateft in the red ; lefs in the yellow, ftill lefs in the blue, and leaft in the violet. And to make as just an estimation as I could of the proportions of their contractions or dilations, I observed that the whole contraction or dilation of the Diameter of any Ring made by all the degrees of red, was to that of the Diameter of the fame Ring made by all the degrees of violet, as about four to three, or five to four, and that when the Light was of the middle Colour between yellow and green, the Diameter of the Ring was very nearly an arithmetical mean between the greatest Diameter of the fame Ring made by the outmost red, and the least Diameter thereof made by the outmost violet : Contrary to what happens in the Colours of the oblong Spectrum made by the refraction of a Prism, where the red is most contracted, the violet most expanded, and Dd in

in the midst of all the Colours is the confine of green and blue. And hence I feem to collect that the thickneffes of the Air between the Glaffes there, where the Ring is fucceffively made by the limits of the five principal Colours (red, yellow, green, blue, violet) in order (that is, by the extreme red, by the limit of red and vellow in the middle of the orange, by the limit of yellow and green, by the limit of green and blue, by the limit of blue and violet in the middle of the indigo, and by the extreme violet ) are to one another very nearly as the fix lengths of a Chord which found the notes in a fixth Major, sol, la, mi, fa, sol, la. But it agrees fomething better with the Observation to fay, that the thickneffes of the Air between the Glaffes there, where the Rings are fucceffively made by the limits of the feven Colours, red, orange, yellow, green, blue, indigo, violet in order, are to one another as the Cuberoots of the Squares of the eight lengths of a Chord, which found the notes in an eighth, sol, la, fa, fol, la, mi, fa, fol; that is, as the Cube-roots of the Squares 

#### OBS. XV.

Thefe Rings were not of various Colours like thofe made in the open Air, but appeared all over of that prifmatique Colour only with which they were illuminated. And by projecting the prifmatique Colours immediately upon the Glaffes, I found that the Light which fell on the dark Spaces which were between the coloured Rings, was transmitted through the Glaffes without any variation of Colour. For on a white

white Paper placed behind, it would paint Rings of the fame Colour with those which were reflected, and of the bignefs of their immediate Spaces. And from thence the origin of these Rings is manifest; namely, That the Air between the Glasses, according to its various thicknefs, is difposed in some places to reflect, and in others to transmit the Light of any one Co-lour (as you may see represented in the sourch Figure) Fig. 4. and in the fame place to reflect that of one Colour where it transmits that of another.

OBS. XVI. The Squares of the Diameters of these Rings made by any prismatique Colour were in arithmetical progreffion as in the fifth Observation. And the Diameter of the fixth Circle, when made by the citrine yellow, and viewed almost perpendicularly, was about 100 parts of an Inch, or a little lefs, agreeable to the fixth Ob-

fervation. The precedent Obfervations were made with a rarer thin medium, terminated by a denfer, fuch as was Air or Water compressed between two Glasses In those that follow are fet down the appearances of a denfer medium thin'd within a rarer, fuch as are plates of Muscovy-glass, Bubbles of Water, and some other thin fubstances terminated on all fides with Air.

D'd' 2 lo tu) evend to observe

[20]

#### OBS. XVII.

If a Bubble be blown with Water first made tenacious by diffolving a little Soap in it, 'tis a common Obfervation, that after a while it will appear tinged with a great variety of Colours. To defend these Bubbles from being agitated by the external Air (whereby their Colours are irregularly moved one among another, fo that no accurate Obfervation can be made of them,) as foon as I had blown any of them I covered it with a clear Glass, and by that means its Colours emerged in a very regular order, like fo many concentrick Rings incompaffing the top of the Bubble. And as the Bubble grew thinner by the continual fubfiding of the Water, thefe Rings dilated flowly and over-fpread the whole Bubble, descending in order to the bottom of it, where they vanished fucceffively. In the mean while, after all the Colours were emerged at the top, there grew in the Center of the Rings a fmall round black Spot, like that in the first Observation, which continually dilated it felf till it became fometimes more than  $\frac{1}{2}$  or  $\frac{3}{4}$  of an Inch in breadth before the Bubble broke. At first I thought there had been no Light reflected from the Water in that place, but observing it more curioufly, I faw within it feveral smaller round Spots, which appeared much blacker and darker than the reft, whereby I knew that there was fome reflexion at the other places which were not fo dark as those Spots. And by further tryal I found that I could fee the Images of fome things (as of a Candle or the Sun ) very faintly reflected, not only from the great black Spot, but alfo

also from the little darker Spots which were within it.

Befides the aforefaid coloured Rings there would often appear finall Spots of Colours, afcending and defcending up and down the fides of the Bubble, by reafon of fome inequalities in the fubfiding of the Water. And fometimes finall black Spots generated at the fides would afcend up to the larger black Spot at the top of the Bubble, and unite with it.

#### O B S. XVIII.

Becaufe the Colours of thefe Bubbles were more extended and lively than thofe of the Air thin'd between two Glaffes, and fo more eafy to to diffinguifhed, I fhall here give you a further defcription of their order, as they were observed in viewing them by reflexion of the Skies when of a white Colour, whilft a black Subftance was placed behind the Bubble. And they were thefe, red, blue; red, blue; red, blue; red, green; red, yellow, green, blue, purple; red, yellow, green, blue, violet; red, yellow, white, blue, black.

The three first Successions of red and blue were very dilute and dirty, especially the first, where the red. feemed in a manner to be white. Among these there was fearce any other Colour sensible besides red and blue, only the blues (and principally the second blue) inclined a little to green.

The fourth red was alfo dilute and dirty, but not fo much as the former three; after that fucceeded little or no yellow, but a copious green, which at first inclined a little to yellow, and then became a pretty brifque and and good willow green, and afterwards changed to a bluish Colour; but there succeeded neither blue nor violet.

The fifth red at first inclined very much to purple, and afterwards became more bright and brisque, but yet not very pure. This was fucceeded with a very bright and intense yellow, which was but little in quantity, and foon changed to green : But that green was copious and fomething more pure, deep and lively, than the former green. After that followed an excellent blue of a bright sky-colour, and then a purple, which was les in quantity than the blue, and much inclined to red.

The fixth Red was at first of a very fair and lively Scarlet, and soon after of a brighter Colour, being very pure and brisque, and the best of all the reds. Then after a lively orange followed an intense bright and copious yellow, which was also the best of all the yellows, and this changed first to a greenish yellow, and then to a greenish blue; but the green between the yellow and the blue, was very little and dilute, seeming rather a greenish white than a green. The blue which sky-colour, but yet something inferior to the former blue; and the violet was intense and deep with little or no redness in it. And less in quantity than the blue.

In the laft red appeared a tincture of fcarlet next to violet, which foon changed to a brighter Colour, inclining to an orange; and the yellow which followed was at first pretty good and lively, but afterwards it grew more dilute, until by degrees it ended in perfect white-

whitenefs. And this whitenefs, if the Water was very tenacious and well-tempered, would flowly fpread and dilate it felf over the greater part of the Bubble; continually growing paler at the top, where at length it would crack in many places, and those cracks, as they dilated, would appear of a pretty good, but yet obscure and dark sky-colour; the white between the blue Spots diminishing, until it resembled the threds of an irregular Net-work, and foon after vanished and left all the upper part of the Bubble of the faid dark blue Colour. And this Colour, after the aforefaid manner, dilated it felf downwards, until sometimes it hath overspread the whole Bubble. In the mean while at the top, which was of a darker blue than the bottom, and appeared alfo full of many round blue Spots, fomething darker than the reft, there would emerge one or more very black Spots, and within those other Spots of an intenfer blacknefs, which I mentioned in the former Observation; and these continually dilated themfelves until the Bubble broke.

If the Water was not very tenacious the black Spots would break forth in the white, without any fenfible intervention of the blue. And fometimes they would break forth within the precedent yellow, or red, or perhaps within the blue of the fecond order, before the intermediate Colours had time to difplay themfelves.

By this defcription you may perceive how great an affinity these Colours have with those of Air defcribed in the fourth Observation, although set down in a contrary order, by reason that they begin to appear when the Bubble is thickess, and are most conveniently niently reckoned from the loweft and thickeft part of the Bubble upwards.

#### OBS. XIX.

Viewing in feveral oblique pofitions of my Eye the Rings of Colours emerging on the top of the Bubble, I found that they were fenfibly dilated by increasing the obliquity, but yet not fo much by far as those made by thin'd Air in the feventh Observation. For there they were dilated fo much as, when viewed most obliquely, to arrive at a part of the plate more than twelve times thicker than that where they appeared when viewed perpendicularly; whereas in this cafe the thickness of the Water, at which they arrived when viewed most obliquely, was to that thickness which exhibited them by perpendicular rays, fomething less than as 8 to 5. By the best of my Observations it was between 15 and  $15\frac{1}{2}$  to 10, an increase about 24 times less than in the other cafe.

Sometimes the Bubble would become of an uniform thicknels all over, except at the top of it near the black Spot, as I knew, becaufe it would exhibit the fame appearance of Colours in all pofitions of the Eye. And then the Colours which were feen at its apparent circumference by the obliqueft rays, would be different from those that were feen in other places, by rays lefs oblique to it. And divers Spectators might fee the fame part of it of differing Colours, by viewing it at very differing obliquities. Now observing how much the Colours at the fame places of the Bubble, or at divers places of equal thicknes, were varied by the feveral feveral obliquities of the rays; by the affiftance of the 4th, 14th, 16th and 18th Obfervations, as they are hereafter explained, I collect the thickness of the Water requisite to exhibit any one and the same Colour, at several obliquities, to be very nearly in the proportion expressed in this Table.

In t	iciden he W	ce on ater.	Refract to the V	ion in- Vater.	Thickness of the Water.		
-	deg.	min.	deg.	min.			
	00	00	00	00	IO		
	15	00	II	II			
	30	00	22	I	10 <sup>4</sup>		
	45	00	32	2	I I 4		
	60	00	40	30	13.		
	75	00	46	25	141		
	90	00	:48	35	155		

In the two first Columns are expressed the obliquities of the rays to the superficies of the Water, that is, their Angles of incidence and refraction. Where I suppose that the Sines which measure them are in round numbers as 3 to 4, though probably the diffolution of Soap in the Water, may a little alter its refractive Vertue. In the third Column the thickness of the Bubble, at which any one Colour is exhibited in those several obliquities, is express in parts, of which ten conflitute that thickness when the rays are perpendicular.  $\times$ 

I have sometimes observed, that the Colours which arise on polished Steel by heating it, or on Bell-metal, and some other metalline substances, when melted and "Wilk these measures the Rule found out by 7<sup>th</sup> poured on if rightly applied exactly agrees. That is, that the Thickness of a glate of Waler, by which one and colour in all changes of observations of gene appears, is proportional to g sean of that angle whose fine is first out of 106 arithmetical mean proportions between y fine of meidence and Assaction; begining from the least of the sines the from g sine of Refraction, when g Res. (metion is out of air into water, but jother was from y dine of me poured on the ground, where they may cool in the open Air, have, like the Colours of Water-bubbles, been a little changed by viewing them at divers obliquities, and particularly that a deep blue, or violet, when viewed very obliquely, hath been changed to a deep red. But the changes of these Colours are not fo great and fensible as of those made by Water. For the Scoria or vitrified part of the Metal, which most Metals when heated or melted do continually protrude, and fend out to their furface, and which by covering the Metals in form of a thin glass skin, causes these Colours, is much denser than Water; and I find that the change made by the obliquation of the Eye is least in Colours of the denser thin fubstances.

#### OBS. XX.

As in the ninth Obfervation, fo here, the Bubble, by tranfmitted Light, appeared of a contrary Colour to that which it exhibited by reflexion. Thus when the Bubble being looked on by the Light of the Clouds reflected from it, feemed red at its apparent circumference, if the Clouds at the fame time, or immediately after, were viewed through it, the Colour at its circumference would be blue. And, on the contrary, when by reflected Light it appeared blue, it would appear red by transmitted Light.

#### OBS. XXI.

By wetting very thin plates of Muscovy-glass, whose thinness made the like Colours appear, the Colours became became more faint and languid ; efpecially by wetting the plates on that fide oppofite to the Eye : But I could not perceive any variation of their fpecies. So then the thicknefs of a plate requifite to produce any Colour, depends only on the denfity of the plate, and not on that of the ambient medium: And hence, by the 10th and 16th Obfervations, may be known the thicknefs which Bubbles of Water, or Plates of Mufcovyglafs, or other fubftances, have at any Colour produced by them.

#### OBS. XXII.

A thin transparent Body, which is denser than its ambient medium, exhibits more brisque and vivid Colours than that which is fo much rarer; as I have particularly observed in the Air and Glass. For blowing Glass very thin at a Lamp-furnace, those plates Sychno-fornace incompassed with Air did exhibit Colours much more vivid than those of Air made thin between two Glasses.

#### OBS. XXIII.

Comparing the quantity of Light reflected from the feveral Rings, I found that it was most copious from the first or inmost, and in the exterior Rings became gradually lefs and lefs. Also the whiteness of the first Ring was stronger than that reflected from those parts of the thinner medium which were without the Rings; as I could manifestly perceive by viewing at a distance the Rings made by the two Object-E e 2 Glasses, Glaffes; or by comparing two Bubbles of Water blown at diftant times, in the first of which the whiteness appeared, which fucceeded all the Colours, and in the other, the whiteness which preceded them all.

#### OBS. XXIV.

When the two Object-Glaffes were lay'd upon one another, fo as to make the Rings of the Colours appear, though with my naked Eye I could not difcern above 8 or 9 of those Rings, yet by viewing them through a Prism I have seen a far greater multitude, infomuch that I could number more than forty, befides many others, that were so very small and close together, that I could not keep my Eye steddy on them feverally so as to number them, but by their extent I have so fometimes estimated them to be more than a hundred. And I believe the Experiment may be improved to the discovery of far greater numbers. For they seem to be really unlimited, though visible only fo far as they can be separated by the refraction, as I shall hereafter explain.

But it was but one fide of these Rings, namely, that towards which the refraction was made, which by that refraction was rendered distinct, and the other fide became more confused than when viewed by the naked Eye, infomuch that there I could not discern above one or two, and sometimes none of those Rings, of which I could discern eight or nine with my naked Eye. And their Segments or Arcs, which on the other fide appeared so numerous, for the most part exceeded exceeded not the third part of a Circle. If the Refraction was very great, or the Prifm very diftant from the Object-Glaffes, the middle part of those Arcs became alfo confused, so as to difappear and conftitute an even whiteness, whilst on either fide their ends, as also the whole Arcs furthest from the center, became diftincter than before, appearing in the form as you see. them designed in the fifth Figure. Fig. 5,

The Arcs, where they feemed diffincteft, were only white and black fucceffively, without any other Colours intermixed. But in other places there appeared. Colours, whofe order was inverted by the refraction in fuch manner, that if I firft held the Prifin very near the Object-Glaffes, and then gradually removed it further off towards my Eye, the Colours of the 2d, 3d, 4th, and following Rings fhrunk towards the white that emerged between them, until they wholly vanifhed into it at the middle of the Arcs, and afterwards emerged again in a contrary order. But at the ends of the Arcs they retained their order unchanged.

I have fometimes fo lay'd one Object-Glafs upon the other, that to the naked Eye they have all over feemed uniformly white, without the leaft appearance of any of the coloured Rings; and yet by viewing them through a Prifm, great multitudes of thofe Rings have difcovered themfelves. And in like manner plates of Mufcovy-glafs, and Bubbles of Glafs blown at a Lamp-furnace, which were not fo thin as to exhibit any Colours to the naked Eye, have through the Prifm exhibited a great variety of them ranged irregularly up and down in the form of waves. And fo Bubbles Bubbles of Water, before they began to exhibit their Colours to the naked Eye of a By-ftander, have appeared through a Prifm, girded about with many parallel and horizontal Rings; to produce which effect, it was neceffary to hold the Prifm parallel, or very nearly parallel to the Horizon, and to difpofe it fo that the rays might be refracted upwards.

THE

# [31] THE SECOND BOOK OF OPTICKS.

### PART II.

#### Remarks upon the foregoing Observations.

Aving given my Obfervations of these Colours, before I make use of them to unfold the Causes of the Colours of natural Bodies, it is convenient that by the simplest of them, such as are the 2d, 3d, 4th, 9th, 12th, 18th, 20th, and 24th, I first explain the more expounded. And first to shew how the Colours in the fourth and eighteenth Observations are produeed, let there be taken in any right line from the point Y, the lengths YA, YB, YC, YD, YE, YF, YG, Fig. 5. YH, in proportion to one another, as the Cube-roots of the Squares of the numbers,  $\frac{1}{2}, \frac{2}{10}, \frac{2}{3}, \frac{2}{3}, \frac{2}{5}, \frac{5}{9}, 1$ , whereby the lengths of a musical Chord to found all the Notes in an Eighth are represented; that is, in the proportion of the numbers 6300, 6814, 7114, 7631, 8255, 8855, 9243, 10000. And at the points A, B, C, D, E, F, E, F, G, H, let perpendiculars  $A \alpha$ ,  $B \beta$ ,  $\Im c$ . be erected, by whole intervals the extent of the feveral Colours fet underneath against them, is to be represented. Then divide the line  $A \alpha$  in fuch proportion as the numbers 1, 2, 3, 5, 6, 7, 9, 10, 11,  $\Im c$ . fet at the points of divifion denote. And through those divisions from Y draw lines 1 I, 2 K, 3 L, 5 M, 6 N, 7 O,  $\Im c$ .

Now if A 2 be supposed to represent the thickness of any thin transparent Body, at which the outmost violet is most copiously reflected in the first Ring, or Series of Colours, then by the 13th Observation HK, will reprefent its thickness, at which the utmost red is most copiously reflected in the fame Series. Alfo by the 5th and 16th Observations, A6 and HN will denote the thickneffes at which those extreme Colours are most copiously reflected in the fecond Series, and A 10 and HQ the thickneffes, at which they are most copiously reflected in the third Series, and so on. And the thickness at which any of the intermediate Colours are reflected most copiously, will, according to the 14th Observation, be defined by the distance of the line A H from the intermediate parts of the lines 2 K, 6 N, 10 Q, &c. against which the names of those Colours are written below.

But further, to define the latitude of these Colours in each Ring or Series, let A I defign the least thickness, and A 3 the greatest thickness, at which the extreme violet in the first Series is reflected, and let H I, and H L, defign the like limits for the extreme red, and let the intermediate Colours be limited by the intermediate parts of the lines I I, and 3 L, against which the names of those Colours are written, and so on : But

yet

yet with this caution, that the reflections be supposed Atrongest at the intermediate Spaces, 2K, 6N, 10Q, &c. and from thence to decrease gradually towards these limits, 1 I, 3 L, 5 M, 7 O, &c. on either fide; where you must not conceive them to be precisely limited, but to decay indefinitely. And whereas I have affigned the fame latitude to every Series, I did it, becaufe although the Colours in the first Series seem to be a little broader than the reft, by reason of a stronger reflexion there, yet that inequality is fo infenfible as fcarcely to be determined by Observation.

Now according to this description, conceiving that the rays originally of feveral Colours are by turns reflected at the Spaces 1 I L 3, 5 M O 7, 9 P R 11, &c. and transmitted at the Spaces A H I 1, 3 L M 5, 7 O P 9, bc. it is easy to know what Colour must in the open Air be exhibited at any thickness of a transparent thin body. For if a Ruler be applied parallel to AH, at that diftance from it by which the thickness of the body is represented, the alternate Spaces 1 IL 3, 5 MO 7, 5c. which it croffeth will denote the reflected original Colours, of which the Colour exhibited in the open Air is compounded. Thus if the conftitution of the green in the third Series of Colours be defired, apply the Ruler as you see at meon, and by its paffing through fome of the blue at  $\pi$  and yellow at  $\sigma$ , as well as through the green at e, you may conclude that the green exhibited at that thickness of the body is principally constituted of original green, but not without a mixture of fome blue and yellow.

By this means you may know how the Colours from the center of the Rings outward ought to fucceed in order as they were defcribed in the 4th and 18th Obfervations. For if you move the Ruler gradually from AH through all distances, having past over the first fpace which denotes little or no reflexion to be made by thinnest substances, it will first arrive at I the violet, and then very quickly at the blue and green, which together with that violet compound blue, and then at the yellow and red, by whole further addition that blue is converted into whitenels, which whitenels continues during the transit of the edge of the Ruler from I to 3, and after that by the fucceffive deficience of its component Colours, turns first to compound yellow, and then to red, and last of all the red ceaseth at L. Then begin the Colours of the fecond Series, which fucceed in order during the transit of the edge of the Ruler from 5 to O, and are more lively than before, becaufe more expanded and fevered. And for the fame reason, instead of the former white there intercedes between the blue and yellow a mixture of orange, yellow, green, blue and indico, all which together ought to exhibit a dilute and imperfect green. So the Co-lours of the third Series all fucceed in order; first, the violet, which a little interferes with the red of the fecond order, and is thereby inclined to a reddifh purple; then the blue and green, which are lefs mixed with other Colours, and confequently more lively than before, especially the green: Then follows the yellow, fome of which towards the green is diffinct and good, but that part of it towards the fucceeding red, as alfo that red is mixed with the violet and blue of the fourth Series,

ries, whereby various degrees of red very much inclining to purple are compounded. This violet and blue, which should succeed this red, being mixed with, and hidden in it, there fucceeds a green. And this at first is much inclined to blue, but foon becomes a good green, the only unmixed and lively Colour in this fourth Series. For as it verges towards the yellow, it begins to interfere with the Colours of the fifth Series, by whofe mixture the fucceeding yellow and red are very much diluted and made dirty, especially the yellow, which being the weaker Colour is fcarce able to fhew it felf. After this the feveral Series interfere more and more, and their Colours become more and more intermixed, till after three or four more revolutions (in which the red and blue predominate by turns) all forts of Colours are in all places pretty equally bended, and compound an even whitenefs.

And fince by the 15th Observation the rays indued with one Colour are transmitted, where those of another Colour are reflected, the reason of the Colours made by the transmitted Light in the 9th and 20th Obfervations is from hence evident.

If not only the order and species of these Colours, but also the precise thickness of the plate, or thin body at which they are exhibited, be defired in parts of an Inch, that may be also obtained by affistance of the 6th or 16th Observations. For according to those Observations the thickness of the thinned Air, which between two Glasses exhibited the most luminous parts of the first fix Rings were  $\frac{1}{178000}$ ,  $\frac{3}{178000}$ ,  $\frac{7}{178000}$ ,  $\frac{9}{178000}$ ,  $\frac{11}{178000}$  parts of an Inch. Suppose the Light reflected most copiously at these thickness be the bright citrine yellow, or con-F f 2 fine of yellow and orange, and these thicknesses will be  $G\mu$ ,  $G\nu$ ,  $G\xi$ , Go, G7. And this being known, it is easy to determine what thickness of Air is represented by  $G\rho$ , or by any other distance of the ruler from A H.

But further, fince by the 10th Observation the thicknefs of Air was to the thicknefs of Water, which between the fame Glaffes exhibited the fame Colour, as 4 to 3, and by the 21th Observation the Colours of thin bodies are not varied by varying the ambient medium; the thickness of a Bubble of Water, exhibiting any Colour, will be  $\frac{3}{4}$  of the thickness of Air producing the fame Colour. And fo according to the fame 10th and 21th Observations the thickness of a plate of Glafs, whofe refraction of the mean refrangible ray, is measured by the proportion of the Sines 31 to 20, may be  $\frac{20}{31}$  of the thickness of Air producing the same Colours; and the like of other mediums. I do not affirm, that this proportion of 20 to 31, holds in all the rays; for the Sines of other forts of rays have other proportions. But the differences of those proportions are fo little that I do not here confider them. On thefe Grounds I have composed the following Table, wherein the thickness of Air, Water, and Glass, at which each Colour is most intense and specifick, is expreffed in parts of an Inch divided. into. Ten hundred thousand equal parts.

many of the second

The The

# [37]

The thickness of coloured	Plates and	Particles	of
---------------------------	------------	-----------	----

		Ain	MZ at all	CL
	allow DI- 1	<u></u>	r aler.	Grajs.
	Plack	2	8	31
	DIACK	I	4	31
	Beginning of	2	14	$\mathbf{I}_{7}^{\frac{2}{7}}$
Their Colours of the	E Black.			13
first Order.	Blue	25	I.5	120
	, white	54	38	35
	Yellow	75	53	45
	Orange	8	6	56
:	Red	9	$6_{4}^{3}$	55
	Violet	115	83	75
	Indico	125	98	811
	Blue	14	$IO_2^{\frac{1}{2}}$	9
Of the fecond Order	Green	1518	113	95
or the record order,	Yellow	167	125	101
	Orange	173	13	II
	Bright Red	18:	134	II
	Scarlet	19	144	12
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	Purple	21	1152	1311
	Indico	22-1	164	14
	Blue	23	1711	1515
Of the third Order.	Green	25!	182	16:
	Vellow	271	201	171
	Red	20	212	182
	'Bluifh Red'	32	24	201
The state of the s	Dhill Com	124	OF!	00
	Bluin Green	34	2)2 061	22
Of the fourth Order,	JGreen	377	202	224 03 <sup>2</sup>
	Dellowith Green	50	201	239.
	Ked	403	304	20
Of the fifth Order	Greenish Blue	46	34=	293
or the men ordery	Red	521	398	34
Of the fixth Order	Greenish Blue	581	44	38
or the fixer order,	Red	65	482	42.
001 0 1 0 0	Greenith Pluse	71	52	1 45%
Of the feventh Order,	Ruddy White	77	573	491
	Circuma. At mee-		3/4	Nou

Now if this Table be compared with the 6th Scheme, you will there fee the conftitution of each Colour, as to its Ingredients, or the original Colours of which it is compounded, and thence be enabled to judge of its intenseness or imperfection; which may fuffice in ex-plication of the 4th and 18th Observations, unless it be further defired to delineate the manner how the Colours appear, when the two Object-Glaffes are lay'd upon one another. To do which, let there be defcribed a large Arc of a Circle, and a streight Line which may touch that Arc, and parallel to that Tangent feveral occult Lines, at fuch diftances from it, as the numbers fet against the several Colours in the Table denote. For the Arc, and its Tangent, will represent the fuperficies of the Glaffes terminating the interjacent Air; and the places where the occult Lines cut the Arc will flow at what diftances from the Center, or Point of contact, each Colour is reflected.

There are also other uses of this Table : For by its affistance the thickness of the Bubble in the 19th Obfervation was determined by the Colours which it ex-And fo the bigness of the parts of natural hibited. Bodies may be conjectured by their Colours, as shall be hereafter shewn. Also, if two or more very thin plates be lay'd one upon another, so as to compose one plate equalling them all in thickness, the refulting Colour may be hereby determined. For inftance, Mr. Hook in his Miscrographia observes, that a faint yellow plate of Lapidis specularis Muscovy-glass lay'd upon a blue one, constituted a very deep purple. The yellow of the first Order is a faint one, and the thickness of the plate exhibiting it, according to the Table is 43, to which add 9, the thick-V.S. 124 nefs

nefs exhibiting blue of the fecond Order, and the fum will be  $13\frac{3}{5}$ , which is the thicknefs exhibiting the purple of the third Order.

To explain, in the next place, the Circumstances of the 2d and 3d Observations; that is, how the Rings of the Colours may (by turning the Prisms about their common Axis the contrary way to that expressed in those Observations) be converted into white and black Rings, and afterwards into Rings of Colours again, the Colours of each Ring lying now in an inverted order; it must be remembred, that those Rings of Colours are dilated by the obliquation of the rays to the Air which. intercedes the Glasses, and that according to the Table in the 7th Observation, their dilatation or increase of their Diameter is most manifest and speedy when they are obliquest. Now the rays of yellow being more refracted by the first superficies of the faid Air than those of red, are thereby made more oblique to the fecond fuperficies, at which they are reflected to produce the coloured Rings, and confequently the yellow Circle in each Ring will be more dilated than the red; and the excess of its dilatation will be fo much the greater, by how much the greater is the obliquity of the rays, until at last it become of equal extent with the red of the fame Ring. And for the fame reafon the green, blue and violet, will be alfo fo much dilated by the still greater obliquity of their rays, as to become all very nearly of equal extent with the red, that is, equally diftant from the center of the Rings. And then all the Colours of the fame Ring must be coincident, and by their mixture exhibit a white Ring. And these white Rings must have black and dark Rings between them, because they do not fpread

fpread and interfere with one another as before. And for that reafon alfo they muft become diffincter and vifible to far greater Numbers. But yet the violet being obliqueft will be fomething more dilated in proportion to its extent then the other Colours, and fo very apt to appear at the exterior verges of the white.

Afterwards, by a greater obliquity of the rays, the violet and blue become more fenfibly dilated than the red and yellow, and fo being further removed from the center of the Rings, the Colours muft emerge out of the white in an order contrary to that which they had before, the violet and blue at the exterior limbs of each Ring, and the red and yellow at the interior. And the violet, by reafon of the greateft obliquity of its rays, being in proportion moft of all expanded, will fooneft appear at the exterior limb of each white Ring, and become more confpicuous than the reft. And the feveral Series of Colours belonging to the feveral Rings, will, by their unfolding and fpreading, begin again to interfere, and thereby render the Rings lefs diftinct, and not vifible to fo great numbers.

If inftead of the Prifms the Object-glaffes be made ufe of, the Rings which they exhibit become not white and diffinct by the obliquity of the Eye, by reafon that the rays in their paffage through that Air which intercedes the Glaffes are very nearly parallel to those Lines in which they were first incident on the Glaffes, and confequently the rays indued with feveral Colours are not inclined one more than another to that Air, as it happens in the Prifms.

There is yet another circumstance of these Experiments to be confidered, and that is why the black and white Rings

Rings which when viewed at a diftance appear diftinct, fhould not only become confused by viewing them near at hand, but also yield a violet Colour at both the edges of every white Ring. And the reafon is, that the rays which enter the Eye at feveral parts of the Pupil, have feveral obliquities to the Glaffes, and those which are most oblique, if confidered apart, would represent the Rings bigger than those which are the least oblique. Whence the breadth of the perimeter of every white Ring is expanded outwards by the obliquest rays, and inwards by the leaft oblique. And this expansion is fo much the greater by how much the greater is the difference of the obliquity ; that is, by how much the Pupil is wider, or the Eye nearer to the Glaffes. And the breadth of the violet must be most expanded, because the rays apt to excite a sensation of that Colour are most oblique to a second, or further superficies of the thin'd Air at which they are reflected, and have also the greatest variation of obliquity, which makes that Colour foonest emerge out of the edges of the white. And as the breadth of every Ring is thus augmented, the dark intervals must be diminished, until the neighbouring Rings become continuous, and are blended, the exterior first, and then those nearer the Center, fo that they can no longer be diftinguish'd apart, but seem to constitute an even and uniform whitenels.

Among all the Observations there is none accompanied with so odd circumstances as the 24th. Of those the principal are, that in thin plates, which to the naked Eye feem of an even and uniform transparent Gg white-

#### [42]

whitenefs, without any terminations of shadows, the refraction of a Prism should make Rings of Colours appear, whereas it ufually makes Objects appear coloured only there where they are terminated with shadows, or have parts unequally luminous; and that it should make those Rings exceedingly distinct and white, although it ufually renders Objects confused and coloured. The caufe of these things you will understand by confidering, that all the Rings of Colours are really in the plate, when viewed with the naked Eye, although by reafon of the great breadth of their circumferences they fo much interfere and are blended together, that they feem to conftitute an even whitenefs. But when the rays pass through the Prism to the Eye, the orbits of the ieveral Colours in every Ring are refracted, some more than others, according to their degrees of refrangibility : By which means the Colours on one fide of the Ring (that is on one fide of its Center) become more unfolded and dilated, and those on the other fide more complicated and contracted. And where by a due refraction they are so much contracted, that the fevral Rings become narrower than to interfere with one another, they must appear distinct, and also white, if the constituent Colours be fo much contracted as to be wholly coincident. But, on the other fide, where the orbit of every Ring is made broader by the further unfolding of its Colours, it must interfere more with other Rings than before, and fo become lefs diffinct.

Fig. 7.

To explain this a little further, fuppofe the concentrick Circles A V, and BX, reprefent the red and violet of any order, which, together with the intermediate Colours,
Colours, conftitute any one of these Rings. Now these being viewed through a Prifm, the violet Circle BX, will by a greater refraction be further translated from its place than the red AV, and fo approach nearer to it on that fide, towards which the refractions are made. For instance, if the red be translated to av, the violet may be translated to  $b \propto$ , fo as to approach nearer to it at x than before, and if the red be further translated to av, the violet may be fo much further translated to b x as to convene with it at x, and if the red be yet further translated to  $\alpha r$ , the violet may be still fo much further translated to  $\beta \xi$  as to pass beyond it at  $\xi$ , and convene with it at e and f. And this being underftood not only of the red and violet, but of all the other intermediate Colours, and alfo of every revolution of those Colours, you will easily perceive how those of the fame revolution or order, by their nearnefs at xv and  $r \xi$ , and their coincidence at x v, e and f, ought to constitute pretty distinct Arcs of Circles, especially at xv, or at e and f, and that they will appear feverally at xv, and at x v exhibit whiteness by their coincidence. and again appear feveral at  $r \xi$ , but yet in a contrary order to that which they had before, and still retain beyond e and f. But, on the other fide, at ab, a b, or « B, these Colours must become much more confufed by being dilated and fpread fo, as to interfere with those of other Orders. And the fame confusion will happen at  $T \xi$  between e and f, if the refraction be very great, or the Prism very distant from the Object-Glasses: In which cafe no parts of the Rings will be feen, fave only two little Arcs at e and f, whose distance from one Gg 2 another,

another will be augmented by removing the Prifin ftill further from the Object-Glaffes: And thefe little Arcs muft be diffincteft and whiteft at their middle, and at their ends, where they begin to grow confused they muft be coloured. And the Colours at one end of every Arc muft be in a contrary order to those at the other end, by reason that they cross in the intermediate white; namely their ends, which verge towards  $\Upsilon \xi$ , will be red and yellow on that fide next the Center, and blue and violet on the other fide. But their other ends which verge from  $\Upsilon \xi$  will on the contrary be blue and violet on that fide towards the Center, and on the other fide red and yellow.

Now as all these things follow from the Properties of Light by a mathematical way of reasoning, so the truth of them may be manifested by Experiments. For in a dark room, by viewing these Rings through a Prism, by reflexion of the several prismatique Colours, which an affistant causes to move to and fro upon a Wall or Paper from whence they are reflected, whilst the Spectator's Eye, the Prism and the Object-Glasses (as in the 13th Observation) are placed steddy : the position of the Circles made successively by the several Colours, will be found such, in respect of one another, as I have described in the Figures  $ab \times v$ , or  $ab \times v_{T}$ or  $a\beta \xi T$ . And by the same method the truth of the Explications of other Observations may be examined.

By what hath been faid the like Phænomina of Water, and thin plates of Glafs may be underftood. But in fmall fragments of those plates, there is this further

further observable, that where they lye flat upon a Table and are turned about their Centers whilft they are viewed through a Prifin, they will in fome poftures-exhibit waves of various Colours, and fome of them exhibit these waves in one or two positions only, but the most of them do in all positions exhibit them, and make them for the most part appear almost all over the plates. The reason is, that the superficies of such plates are not even, but have many cavities and fwellings, which how shallow soever do a little vary the thickness of the plate. For at the feveral fides of those cavities, forthe reafons newly defcribed, there ought to be produced waves in feveral poftures of the Prilm. Now thoughit be but fome very finall, and narrower parts of the Glafs, by which these waves for the most part are cauied, yet they may feem to extend themselves over the whole Glafs, becaufe from the narroweft of those parts there are Colours of feveral Orders that is of feveral Rings, confuledly reflected, which by refraction of the-Prism are unfolded, separated, and according to their degrees of refraction, difperfed to feveral places, fo as to constitute so many several waves, as there were divers orders of Colours promiscuously reflected from that part of the Glass.

These are the principal Phænomena of thin Plates or Bubbles, whole explications depend on the properties of Light, which I have heretofore delivered. And these you see do necessarily follow from them, and agree with them, even to their very least circumstances; and not only so, but do very much tend to their proof. Thus, by the 24th Observation, it appears, that the rays. rays of feveral Colours made as well by thin Plates or Bubbles, as by refractions of a Prifm, have feveral degrees of refrangibility, whereby those of each order, which at their reflexion from the Plate or Bubble are intermixed with those of other orders, are separated from them by refraction, and affociated together so to become visible by themselves like Arcs of Circles. For if the rays were all alike refrangible, 'tis impossible that the whitenes, which to the naked sence appears uniform, should by refraction have its parts transposed and ranged into those black and white Arcs.

It appears also that the unequal refractions of difform rays proceed not from any contingent irregularities; fuch as are veins, an uneven polifh, or fortuitous position of the pores of Glafs; unequal and cafual motions in the Air or Æther; the fpreading, breaking, or dividing the fame ray into many diverging parts, or the like. For, admitting any fuch irregularities, it would be impossible for refractions to render those Rings fo very diftinct, and well defined, as they do in the 24th Observation. It is neceffary therefore that every ray have its proper and constant degree of refrangibility connate with it, according to which its refraction is ever justly and regularly performed, and that feveral rays have feveral of those degrees.

And what is faid of their refrangibility may be alfo underftood of their reflexibility, that is of their difpofitions to be reflected fome at a greater, and others at a lefs thicknefs, of thin Plates or Bubbles, namely, that those difpositions are alfo connate with the rays, and immutable; as may appear by the 13th, 14th, and 15th 15th Observations compared with the fourth and eighth.

By the precedent Obfervations it appears alfo, that whitenefs is a diffimilar mixture of all Colours, and that Light is a mixture of rays indued with all those Colours. For confidering the multitude of the Rings of Colours, in the 3d, 12th and 24th Observations, it is manifest that although in the 4th and 18th Observations there appear no more than eight or nine of those Rings, yet there are really a far greater number, which so much interfere and mingle with one another, as after those eight or nine revolutions to dilute one another wholly, and constitute an even and fensibly uniform whiteness. And consequently that whiteness must be allowed a mixture of all Colours, and the Light which conveys it to the Eye must be a mixture of rays indued. with all those Colours.

But further, by the 24th Obfervation, it appears, that there is a conftant relation between Colours and Refrangibility, the moft refrangible rays being violet, the leaft refrangible red, and those of intermediate Colours having proportionably intermediate degrees of refrangibility. And by the 13th, 14th and 15th Obfervations, compared with the 4th or 18th, there appears to be the fame conftant relation between Colour and Reflexibility, the violet being in like circumftances reflected at leaft thickneffes of any thin Plate or Bubble, the red at greatest thickneffes, and the intermediate Colours at intermediate thickneffes. Whence it follows, that the colorifique dispositions of rays are also connate with them and immutable, and by confequence that.

## [48]

that all the productions and appearances of Colours in the World are derived not from any phyfical change caufed in Light by refraction or reflexion, but only from the various mixtures or feparations of rays, by virtue of their different Refrangibility or Reflexibility. And in this refpect the Science of Colours becomes a Speculation as truly mathematical as any other part of Optiques. I mean fo far as they depend on the nature of Light, and are not produced or altered by the power of imagination, or by ftriking or preffing the Eyes.

THE









# [49]

#### THE

# SECOND BOOK of OPTICKS.

### PART III.

Of the permanent Colours of natural Bodies, and the Analogy between them and the Colours of thin transparent Plates.

Am now come to another part of this Defign, which is to confider how the Phænomena of thin transparent Plates stand related to those of all other natural Bodies. Of these Bodies I have already told you that they appear of divers Colours, accordingly as they are disposed to reflect most copiously the rays originally indued with those Colours. But their Constitutions, whereby they reflect some rays more copiously than others, remains to be discovered, and these I shall endeavour to manifest in the following Propositions.

Hh

PROP.

## [50]

#### PROP. I.

Those superficies of transparent Bodies reflect the greatest quantity of Light, which have the greatest refracting power; that is, which intercede mediums that differ most in their refractive densities. And in the confines of equally refracting mediums there is no reflexion.

The Analogy between reflexion and refraction will appear by confidering, that when Light paffeth obliquely out of one medium into another which refracts from the perpendicular, the greater is difference of their refractive denfity, the lefs obliquity is requisite to cause a total reflexion. For as the Sines are which measure the refraction, so is the Sine of incidence at which the total reflexion begins, to the radius of the Circle, and confequently that incidence is least where there is the greatest difference of the Sines: Thus in the paffing of Light out of Water into Air, where the refraction is measured by the Ratio of the Sines 3 to 4, the total reflexion begins when the Angle of incidence is about 48 degrees 35 minutes. In paffing out of Glafs into Air, where the refraction is measured by the Ratio of the Sines 20 to 31, the total reflexion begins when the Angle of incidence is 40 deg. 10 min. and fo in paffing out of crystal, or more strongly refracting mediums into Air, there is still a less obliquity requisite to cause a total reflexion. Superficies therefore which refract most do soonest reflect all the Light which is incident on them, and fo must be allowed most strongly reflexive. 1 6 3 3

But the truth of this Proposition will further appear. by observing, that in the superficies interceding two transparent mediums, such as are (Air, Water, Oyl, Common-Glass, Crystal, Metalline-Glasses, Island-Glasses, white transparent Arsnick, Diamonds, Bc.) the reflexion is ftronger or weaker accordingly, as the fuperficies hath a greater or less refracting power. For in the confine of Air and Sal-gemm 'tis stronger than in the confine of Air and Water, and still stronger in the confine of Air and Common-Glassor Crystal, and stronger in the confine of Air and a Diamond. If any of these and fuch like transparent Solids, be immerged in Water, its reflexion becomes much weaker than before, and still weaker if they be immerged in the more ftrongly refracting Liquors of well-rectified oyl of Vitriol or spirit of Turpentine. If Water be diftinguished into two parts, by any imaginary furface, the reflexion in the confine of those two parts is none at all. In the confine of Water and Ice'tis very little, in that of Water and Oyl 'tis fomething greater, in that of Water and Sal-gemm still greater, and in that of Water and Glafs, or Crystal, or other denser substances still greater, accordingly as those mediums differ more or less in their refracting powers. Hence in the confine of Common-Glafs and Crystal, there ought to be a weak reflexion, and a ftronger reflexion in the confine of Common and Metalline-Glafs, though I have not yet tried this. But, in the confine of two Glaffes of equal denfity, there is not any fenfible reflexion, as was shewn in the first Observation. And the fame may be understood of the fuperficies interceding two Crystals, or two Liquors, or any other Subftances in which no refraction is caufed. So then the Hh 2 reason

reafon why uniform pellucid mediums, (fuch as Water, Glafs, or Cryftal) have no fenfible reflexion but in their external fuperficies, where they are adjacent to other mediums of a different denfity, is becaufe all their contiguous parts have one and the fame degree of denfity.

#### PROP. II.

The least parts of almost all natural Bodies are in some measure transparent: And the opacity of those Bodies ariseth from the multitude of reflexions caused in their internal Parts.

That this is fo has been obferved by others, and will eafily be granted by them that have been converfant with Mifcrofcopes. And it may be alfo tryed by applying any fubftance to a Hole through which fome Light is immitted into a dark room. For how opake foever that fubftance may feem in the open Air, it will by that means appear very manifeftly transparent, if it be of a fufficient thinnefs. Only white metalline Bodies must be excepted, which by reason of their exceffive density feem to reflect almost all the Light incident on their first fuperficies, unless by folution in menstruums they be reduced into very fmall particles, and then they become transparent.

#### PROP. III.

Between the parts of opake and coloured Bodies are many spaces, either empty or replenished, with mediums of other densities; as Water between the tinging corpuscles where with any Liquor is impregnated, Air between the aqueous aqueous globules that constitute Clouds or Mists; and for the most part spaces void of both Air and Water, but yet perhaps not wholly void of all substance, between the parts of hard Bodies. \*

The truth of this is evinced by the two precedent Propositions : For by the fecond Proposition there are many reflexions made by the internal parts of Bodies, which, by the first Proposition, would not happen if the parts of those Bodies were continued without any fuch interstices between them, because reflexions are caused only in superficies, which intercede mediums of a differing density by Prop. 1.

But further, that this discontinuity of parts is the principal cause of the opacity of Bodies, will appear by confidering, that opake fubstances become transparent by filling their pores with any fubstance of equal or al-most equal density with their parts. Thus Paper dipped in Water or Oyl, the Oculus mundi Stone steep'd in Water, Linnen-cloth oyled or varnished, and many other fubstances soaked in such Liquors as will intimately pervade their little pores, become by that means more transparent than otherwise; so, on the contrary, the most transparent substances may by evacuating their pores, or separating their parts, be rendred sufficiently opake, as Salts or wet Paper, or the Oculus mundi Stone by being dried, Horn by being fcraped, Glafs by being reduced to powder, or otherwife flawed, Turpen-tine by being flirred about with Water till they mix imperfectly, and Water by being formed into many fmall Bubbles, either alone in the form of froth, or by shaking it together with Oyl of Turpentine, or with fome other convenient Liquor, with which it will not

not perfectly incorporate. And to the increase of the opacity of these Bodies it conduces something, that by the 23th Observation the reflexions of very thin transparent substances are considerably stronger than those made by the same substances of a greater thickness.

#### PROP. IV.

The parts of Bodies and their Interstices must not be less than of some definite bigness, to render them opake and coloured.

For the opakeft Bodies, if their parts be fubtily divided, (as Metals by being diffolved in acid menftruums,  $\Im c$ .) become perfectly transparent. And you may also remember, that in the eighth Observation there was no sensible reflexion at the superficies of the Object-Glasses where they were very near one another, though they did not absolutely touch. And in the 17th Observation the reflexion of the Water-bubble where it became thinness was almost infensible, so as to cause very black Spots to appear on the top of the Bubble by the want of reflected Light.

On these grounds I perceive it is that Water, Salt, Glafs, Stones, and fuch like fubftances, are transparent. For, upon divers confiderations, they seem to be as full of pores or interstices between their parts as other Bodies are, but yet their parts and interstices to be too small to cause reflexions in their common furfaces.

jog

PROP.

## [55]

#### PROP. V.

The transparent parts of Bodies according to their several sizes must reflect rays of one Colour, and transmit those of another, on the same grounds that thin Plates or Bubbles do reflect or transmit those rays. And this I take to be the ground of all their Colours.

For if a thin'd or plated Body, which being of an even thicknefs, appears all over of one uniform Colour, fhould be flit into threds, or broken into fragments, of the fame thicknefs with the plate; I fee no reafon why every thred or fragment fhould not keep its Colour, and by confequence why a heap of those thredsor fragments fhould not conftitute a mass or powder of the same Colour, which the plate exhibited before it was broken. And the parts of all natural Bodies being like fo many fragments of a Plate, must on the same grounds exhibit the same Colours.

Now that they do fo, will appear by the affinity of their properties. The finely coloured Feathers of fome Birds, and particularly those of Peacocks Tails, do in the very fame part of the Feather appear of feveral Colours in feveral positions of the Eye, after the very fame manner that thin Plates were found to do in the 7th and 19th Observations, and therefore arise from the thinnels of the transparent parts of the Feathers ; that is, from the flendernels of the very fine Hairs, or *Capilla*menta, which grow out of the fides of the groffer lateral branches or fibres of those Feathers. And to the fame purpose it is, that the Webs of fome Spiders by being being fpun very fine have appeared coloured, as fome have observed, and that the coloured fibres of some filks by varying the polition of the Eye do vary their Co-Also the Colours of filks, cloths, and other fublour. stances, which Water or Oyl can intimately penetrate, become more faint and obscure by being immerged in those liquors, and recover their vigor again by being dried, much after the manner declared of thin Bodies in the 10th and 21th Observations. Leaf-gold, some forts of painted Glafs, the infusion of Lignum Nephriticum, and some other substances reflect one Colour, and transmit another, like thin Bodies in the 9th and 20th Observations. And some of those coloured powders which Painters use, may have their Colours a little changed, by being very elaborately and finely ground. Where I fee not what can be justly pretended for those changes, befides the breaking of their parts into lefs parts by that contrition after the fame manner that the Colour of a thin Plate is changed by varying its thicknefs. For which reafon alfo it is that the coloured flowers of Plants and Vegitables by being bruifed ufually become more transparent than before, or at least in some degree or other change their Colours. Nor is it much lefs to my purpofe, that by mixing divers liquors very odd and remarquable productions and changes of Co-lours may be effected, of which no caufe can be more obvious and rational than that the faline corpufcles of one liquor do varioufly act upon or unite with the tinging corpufcles of another, fo as to make them fwell, or fhrink (whereby not only their bulk but their denfity alfo may be changed) or to divide them into fmaller corpufcles, (whereby a coloured liquor may become

come transparent) or to make many of them affociate into one cluster, whereby two transparent liquors may compose a coloured one. For we see how apt those faline menstruums are to penetrate and diffolve subftances to which they are applied, and fome of them to precipitate what others diffolve. In like manner, if we confider the various Phænomena of the Atmosphære, we may obferve, that when Vapors are first raifed, they hinder not the transparency of the Air, being divided into parts too fmall to caufe any reflexion in their fuperficies. But when in order to compose drops of rain they begin to coalesce and constitute globules of all intermediate fizes, those globules when they become of a convenient fize to reflect some Colours and transmit others, may conftitute Clouds of various Colours according to their fizes. And I fee not what can be rationally conceived in so transparent a substance as Water for the production of these Colours, besides the various fizes of its fluid and globuler parcels.

#### PROP. VI.

' The parts of Bodies on which their Colours depend, are denser than the medium, which pervades their interstices.

This will appear by confidering, that the Colour of a Body depends not only on the rays which are incident perpendicularly on its parts, but on those alfo which are incident at all other Angles. And that according to the 7th Observation, a very little variation of obliquity will change the reflected Colour where the thin body or small particle is rarer than the ambient. I i medium, medium, infomuch that fuch a finall particle will at diverfly oblique incidences reflect all forts of Colours, in fo great a variety that the Colour refulting from them all, confufedly reflected from a heap of fuch particles, muft rather be a white or grey than any other Colour, or at beft it muft be but a very imperfect and dirty Colour. Whereas if the thin body or finall particle be much denfer than the ambient medium, the Colours according to the 19th Obfervation are fo little changed by the variation of obliquity, that the rays which are reflected leaft obliquely may predominate over the reft fo much as to caufe a heap of fuch particles to appear very intenfly of their Colour.

It conduces also fomething to the confirmation of this Proposition, that, according to the 22th Observation, the Colours exhibited by the denser thin body within the rarer, are more brisque than those exhibited by the rarer within the denser.

#### PROP. VII.

The bigness of the component parts of natural Bodies may be conjectured by their Colours.

For fince the parts of thefe Bodies by Prop. 5. do most probably exhibit the same Colours with a Plate of equal thickness, provided they have the same refractive density; and fince their parts seem for the most part to have much the same density with Water or Glass, as by many circumstances is obvious to collect; to determine the fizes of those parts you need only have recours to the precedent Tables, in which the thickness of Water or Glass exhibiting any Colour is expressed. Thus if if it be defired to know the Diameter of a corpufcle, which being of equal denfity with Glafs fhall reflect green of the third order; the number  $16\frac{1}{4}$  flows it to be  $\frac{16\frac{1}{4}}{1000000}$  parts of an Inch.

The greateft difficulty is here to know of what order the Colour of any Body is. And for this end we muft have recourse to the 4th and 18th Observations, from whence may be collected these particulars.

Scarlets, and other reds, oranges and yellows, if they be pure and intenfe are most probably of the fecond order. Those of the first and third order also may be pretty good, only the yellow of the first order is faint, and the orange and red of the third order have a great mixture of violet and blue.

There may be good greens of the fourth order, but the pureft are of the third. And of this order the green of all vegitables feem to be, partly by reafon of the intenfenefs of their Colours, and partly becaufe when they wither fome of them turn to a greenifh yellow, and others to a more perfect yellow or orange, or perhaps to red, paffing first through all the aforefaid intermediate Colours. Which changes feem to be effected by the exhaling of the moisture which may leave the tinging corpufices more denfe, and fomething augmented by the accretion of the oyly and earthy part of that moisture. Now the green without doubt is of the fame order with those Colours into which it changeth, because the changes are gradual, and those Colours, though usually not very full, yet are often too full and lively to be of the fourth order.

1

Blues and purples may be either of the fecond or third order, but the beft are of the third. Thus the Colour of violets feems to be of that order, becaufe their Syrup by acid Liquors turns red, and by urinous and alcalizale turns green. For fince it is of the nature of Acids to diffolve or attenuate, and of Alcalies to precipitate or incraffate, if the purple Colour of the Syrup was of the fecond order, an acid Liquor by attenuating its tinging corpufcles would change it to a red of the firft order, and an Alcaly by incraffating them would change it to a green of the fecond order ; which red and green, efpecially the green, feem too imperfect to be the Colours produced by thefe changes. But if the faid purple be fuppofed of the third order, its change to red of the fecond, and green of the third, may without any inconvenience be allowed.

If there be found any Body of a deeper and lefs reddifh purple than that of the violets, its Colour moft probably is of the fecond order. But yet their being no Body commonly known whofe Colour is conftantly more deep than theirs, I have made ufe of their name to denote the deepeft and leaft reddifh purples, fuch as manifeftly transcend their Colour in purity.

The *blue* of the firft order, though very faint and little, may poffibly be the Colour of fome fubftances; and particularly the azure Colour of the Skys feems to be of this order. For all vapours when they begin to condenfe and coalefce into finall parcels, become firft of that bignefs whereby fuch an Azure muft be reflected before they can conftitute Clouds of other Colours. And fo this being the firft Colour which vapors begin to reflect, it ought to be the Colour of the hneft and moft tranftransparent Skys in which vapors are not arrived to that grosnels requisite to reflect other Colours, as we find it is by experience.

Whiteness, if most intense and luminous, is that of the first order, if less strong and luminous a mixture of the Colours of several orders. Of this last kind is the whitenefs of Froth, Paper, Linnen, and most white fubstances; of the former I reckon that of white metals to be. For whilst the densest of metals, Gold, if foliated is transparent, and all metals become transparent if diffolved in menstruums or vitrified, the opacity of white metals arifeth not from their denfity alone. They being lefs denfe than Gold would be more transparent than it, did not fome other caufe concur with their denfity to make them opake. And this caufe I take to be fuch a bigness of their particles as fits them to reflect the white of the first order. For if they be of other thickneffes they may reflect other Colours, as is manifest by the Colours which appear upon hot Steel in tempering it, and fometimes upon the furface of melted. metals in the Skin or Scoria which arifes upon them in. their cooling. And as the white of the first order is the strongest which can be made by Plates of transparent fubstances, so it ought to be stronger in the denser subftances of metals than in the rarer of Air, Water and Glafs. Nor do I fee but that metallic fubftances of fuch. a thickness as may fit them to reflect the white of the first order, may, by reason of their great density (according to the tenour of the first of these Propositions) reflect all the Light incident upon them, and fo be as opake and fplendent as its poffible for any Body to be. Gold, or Copper mixed with lefs than half their weight of

of Silver, or Tin, or Regulus of Antimony, in fusion or amalgamed with a very little Mercury become white; which shews both that the particles of white metals have much more fuperficies, and fo are fmaller, than those of Gold and Copper, and also that they are so opake as not to fuffer the particles of Gold or Copper to fhine through them. Now it is fcarce to be doubted, but that the Colours of Gold and Copper are of the fecond or third order, and therefore the particles of white metals cannot be much bigger than is requisite to make them reflect the white of the first order. The volatility of Mercury argues that they are not much bigger, nor may they be much lefs, leaft they lofe their opacity, and become either transparent as they do when attenuated by vitrification, or by folution in menstruums, or black as they do when ground fmaller, by rubbing Silver, or Tin, or Lead, upon other fubstances to draw black The first and only Colour which white metals Lines. take by grinding their particles smaller is black, and therefore their white ought to be that which borders upon the black Spot in the center of the Rings of Colours, that is, the white of the first order. But if you would hence gather the bigness of metallic particles, you must allow for their density. For were Mercury transparent, its density is such that the Sine of incidence upon it (by my computation) would be to the fine of its refraction, as 71 to 20, or 7 to 2. And therefore the thickness of its particles, that they may exhibit the fame Colours with those of Bubbles of Water, ought to be lefs than the thicknefs of the Skin of those Bubbles in the proportion of 2 to 7. Whence its poffible that the particles of Mercury may be as little

38

as the particles of fome transparent and volatile fluids, and yet reflect the white of the first order.

Laftly, for the production of black, the corpufcles must be less than any of those which exhibit Colours. For at all greater fizes there is too much Light refle-Eted to conftitute this Colour. But if they be suppofed a little lefs than is requifite to reflect the white and very faint blue of the first order, they will, according to the 4th, 8th, 17th and 18th Observations, reflect fo very little as to appear intenfly black, and yet may perhaps varioully refract it to and fro within themfelves to long, until it happen to be ftifled and loft, by which means they will appear black in all pofitions of the Eye without any transparency. And from hence may be understood why Fire, and the more fubtile diffolver Putrefaction, by dividing the particles of fubftances, turn them to black, why fmall quantities of black fubstances impart their Colour very freely and intenfly to other fubstances to which they are applied; the minute particles of these, by reason of their very great number, eafily overfpreading the grofs particles of others; why Glafs ground very elaborately with Sand on a copper Plate, 'till it be well polished, makes the Sand, together with what is worn off from the Glafs and Copper, become very black : why black fubftances do foonest of all others become hot in the Sun's Light and burn, (which effect may proceed partly from the multitude of refractions in a little room, and partly from the eafy commotion of fo very fmall corpufcles;) and why blacks are ufually a little inclined to a bluifh Colour. For that they are fo may be feen by illuminating white Paper by Light reflected from black fubftances.

ftances. For the Paper will ufually appear of a bluith white; and the reafon is, that black borders on the obfcure blue of the first order described in the 18th Observation, and therefore reflects more rays of that Colour than of any other.

In these Descriptions I have been the more particular, because it is not impossible but that Miscroscopes may at length be improved to the discovery of the particles of Bodies on which their Colours depend, if they are not already in some measure arrived to that degree of perfection. For if those Instruments are or can be so far improved as with sufficient distinctness to represent Objects five or fix hundred times bigger, than at a Foot diffance they appear to our naked Eyes, I should hope that we might be able to discover some of the greatest of those corpuscles. And by one that would magnify three or four thousand times perhaps they might all be discovered, but those which produce blacknefs. In the mean while I fee nothing material in this Discourse that may rationally be doubted of excepting this Polition, That transparent corpuscles of the fame thickness and density with a Plate, do exhibit the fame Colour. And this I would have underftood not without some latitude, as well because those corpuscles may be of irregular Figures, and many rays must be obliquely incident on them, and fo have a shorter way through them than the length of their Diameters, as because the straitness of the medium pent in on all fides within such corpuscles may a little alter its motions or other qualities on which the reflexion depends. But yet I cannot much fuspect the last, because I have observed of some small Plates of Muscovy-Glass which were of an. even

. fr.

even thicknefs, that through a Mifcrofcope they have appeared of the fame Colour at their edges and corners where the included medium was terminated, which they appeared of in other places. However it will add much to our fatisfaction, if those corpuscles could be difcovered with Mifcrofcopes; which if we shall at length attain to, I fear it will be the utmost improvement of this sense. For it seems impossible to see the more secret and noble works of nature within the corpuscles by reason of their transparency.

#### PROP. VIII.

The cause of Reflexion is not the impinging of Light on the solid or impervious parts of Bodies, as is commonly believed.

This will appear by the following Confiderations. First, That in the passage of Light out of Glass into Air there is a reflexion as ftrong as in its paffage out of Air into Glass, or rather a little stronger, and by many degrees stronger than in its passage out of Glass into Water. And it feems not probable that Air should have more reflecting parts than Water or Glafs. But if that fhould poffibly be fuppofed, yet it will avail nothing; for the reflexion is as ftrong or ftronger when the Air is drawn away from the Glafs, (suppose in the Air-pump invented by Mr. Boyle ) as when it is adjacent to it. Secondly, If Light in its paffage out of Glafs into Air be incident more obliquely than at an Angle of 40 or 41 degrees it is wholly reflected, if lefs obliquely it is in great measure transmitted. Now it is not to be imagined that Light at one degree of obliquity should meet Kk with

with pores enough in the Air to transmit the greater part of it, and at another degree of obliquity fhould meet with nothing but parts to reflect it wholly, espe-cially confidering that in its passage out of Air into Glass, how oblique soever be its incidence, it finds pores enough in the Glafs to transmit the greatest part of it. If any Man suppose that it is not reflected by the Air, but by the outmost superficial parts of the Glass, there is still the fame difficulty : Belides, that fuch a Supposition is unintelligible, and will also appear to be false by applying Water behind some part of the Glass instead of Air. For so in a convenient obliquity of the rays fuppofe of 45 or 46 degrees, at which they are all reflected where the Air is adjacent to the Glafs, they shall be in great measure transmitted where the Water is adjacent to it; which argues, that their reflexion or transmission depends on the constitution of the Air and Water behind the Glass, and not on the striking off the rays upon the parts of the Glass. Thirdly, If the Colours made by a Prism placed at the entrance of a beam of Light into a darkened room be fucceffively cast on a second Prism placed at a greater distance from the former, in fuch manner that they are all alike incident upon it, the fecond Prism may be fo inclined to the incident rays, that those which are of a blue Colour fhall be all reflected by it, and yet those of a red Colour pretty copiously transmitted. Now if the reflexion be caufed by the parts of Air or Glafs, I would ask, why at the fame obliquity of incidence the blue should wholly impinge on those parts fo as to be all reflected, and yet the red find pores enough to be in great measure transmitted. Fourthly, where two Glasses touch one another,

another, there is no fenfible reflexion as was declared in the first Observation; and yet I fee no reason why the rays should not impinge on the parts of Glass as much when contiguous to other Glafs as when contiguous to Air. Fifthly, When the top of a Waterbubble (in the 17th Observation) by the continual subfiding and exhaling of the Water grew very thin, there was fuch a little and almost insensible quantity of Light reflected from it, that it appeared intenfly black ; whereas round about that black Spot, where the Water was thicker, the reflexion was io ftrong as to make the Water feem very white. Nor is it only at the leaft thickness of thin Plates or Bubbles, that there is no manifest reflexion, but at many other thicknesses continually greater and greater. For in the 15th Obfervation the rays of the fame Colour were by turns tranfmitted at one thickness, and reflected at another thicknefs, for an indeterminate number of fucceffions. And yet in the fuperficies of the thinned Body, where it is of any one thickness, there are as many parts for the rays to impinge on, as where it is of any other thicknefs. Sixthly, If reflexion were caufed by the parts of reflecting Bodies, it would be impoffible for thin Plates or Bubbles at the same place to reflect the rays of one Colour and transmit those of another, as they do according to the 13th and 15th Observations. For it is not to be imagined that at one place the rays which for instance exhibit a blue Colour, should have the fortune to dash upon the parts, and those which exhibit a red to hit upon the pores of the Body; and then at another place, where the Body is either a little thicker, or a little thinner, that on the contrary the blue should Kk 2 - hit

hit upon its pores, and the red upon its parts. Laftly, were the rays of Light reflected by impinging on the folid parts of Bodies, their reflexions from polithed Bodies could not be fo regular as they are. For in po-lishing Glass with Sand, Putty or Tripoly, it is not to be imagined that those substances can by grating and fretting the Glass bring all its least particles to an accurate polish; so that all their surfaces shall be truly plain or truly fpherical, and look all the fame way, fo as together to compose one even furface. The fmaller the particles of those fubstances are, the smaller will be the fcratches by which they continually fret and wear away the Glass until it be polished, but be they never so small they can wear away the Glass no otherwise than by grating and fcratching it, and breaking the proturberances, and therefore polifh it no otherwife than by bringing its roughness to a very fine Grain, so that the fcratches and frettings of the furface become too fmall to be visible. And therefore if Light were reflected by impinging upon the folid parts of the Glass, it would be scattered as much by the most polished Glass as by the roughest. So then it remains a Problem, how Glass polished by fretting substances can reflect Light fo regularly as it does. And this Problem is fcarce otherwife to be folved than by faying, that the reflexion of a ray is effected, not by a fingle point of the reflecting Body, but by fome power of the Body which is evenly diffufed all over its furface, and by which it acts upon the ray without immediate contact. For that the parts of Bodies do act upon Light at a distance shall be shewn hereafter.

Now

Now if Light be reflected not by impinging on the folid parts of Bodies, but by fome other principle; its probable that as many of its rays as impinge on the folid parts of Bodies are not reflected but ftifled and loft in the Bodies. For otherwife we muft allow two forts of reflexions. Should all the rays be reflected which impinge on the internal parts of clear Water or Cryftal, those fubftances would rather have a cloudy Colour than a clear transparency. To make Bodies look black, its neceffary that many rays be ftopt, retained and loft in them, and it feems not probable that any rays can be ftopt and ftifled in them which do not impinge on their parts.

And hence we may understand that Bodies are much more rare and porous than is commonly believed. Water is 19 times lighter, and by confequence 19 times rarer than Gold, and Gold is fo rare as very readily and without the least opposition to transmit the magnetick Effluvia, and eafily to admit Quick-filver into its pores, and to let Water pass through it. For a concave Sphere of Gold filled with Water, and fodered up, has upon preffing the Sphere with great force, let the Water squeeze through it, and stand all over its outfide in multitudes of small Drops, like dew, without burfting or cracking the Body of the Gold as I have been informed by an Eye-witnefs. From all which we may conclude, that Gold has more pores than folid parts, and by confequence that Water has above fortytimes more pores than parts. And he that shall find out an Hypothefis, by which Water may be fo rare, and yet not be capable of compression by force, may doubtles by the fame Hypothefis make Gold and Water, and all other

## [70]

other Bodies as much rarer as he pleases, so that Light may find a ready passage through transparent substances. Vid: addenda de megnoles wirhchibus st de modibus corpora

#### PROP. IX.

Bodies reflect and refract Light by one and the same power variously exercised in various circumstances.

This appears by feveral Confiderations. First, Because when Light goes out of Glass into Air, as obliquely as it can poffibly do, if its incidence be made still more oblique, it becomes totally reflected. For the power of the Glass after it has refracted the Light as obliquely as is poffible if the incidence be still made more oblique, becomes too ftrong to let any of its rays go through, and by confequence caufes total reflexions. Secondly, Because Light is alternately reflected and transmitted by thin Plates of Glass for many successions accordingly, as the thickness of the Plate increases in an arithmetical Progreffion. For here the thickness of the Glass determines whether that power by which Glass acts upon Light shall cause it to be reflected, or fuffer it to be transmitted. And, Thirdly, because those furfaces of transparent Bodies which have the greatest refracting power, reflect the greatest quantity of Light, as was shewed in the first Proposition.

#### PROP. X.

If Light be swifter in Bodies than in Vacuo in the propertion of the Sines which measure the refraction of the Bodies, the forces of the Bodies to reflect and refract Light, are are very nearly proportional to the densities of the same Bodies, excepting that uncluous and sulphureous Bodies refract more than others of this same density.

fract more than others of this same density. Let A B represent the refracting plane surface of any 4. Which: Gral. math Body, and IC a ray incident very obliquely upon the 91: 280



Body in C, fo that the Angle ACI may be infinitely little, and let CR be the refracted ray. From a given point B perpendicular to the refracting furface erect BR meeting with the refracted ray CR in R, and if CR reprefent the motion of the refracted ray, and this motion be diftinguished into two motions CB and BR, whereof CB is a parallel to the refracting plane, and BR perpendicular to it : CB shall represent the motion of the incident ray, and BR the motion generated by the refraction, as Opticians have of late explained.

Now if any body or thing in moving through any fpace of a giving breadth terminated on both fides by two parallel plains, be urged forward in all parts of that fpace by forces tending directly forwards towards the laft plain, and before its incidence on the firft plane, had no motion towards it, or but an infinitly little one; and if the forces in all parts of that fpace, between the planes be at equal diffances from the planes equal to one another, but at feveral diffances be bigger or lefs in any given proportion; the motion generated by the forces in the whole paffage of the body or thing through. through that fpace shall be in a fubduplicate proportion of the forces, as Mathematicians will easily understand. And therefore if the space of activity of the refracting superficies of the Body be confidered as such a space, the motion of the ray generated by the refracting force of the Body, during its passage through that space that is the motion BR must be in a subduplicate proportion of that refracting force : I say therefore that the square of the Line BR, and by confequence the refracting force of the Body is very nearly as the density of the same Body. For this will appear by the following Table, wherein the proportion of the Sines which measure the refraxions of several Bodies, the square of BR supposing CB an unite, the densities of the Bodies estimated by their specifick gravities, and their refractive power in respect of their densities are set down in feveral Columns.

The

		•.		.0	
The refracting Bodies.	The Proportion of the Sines of incidence and refraction of yellow Light.		The Square of BR, to which the refracting force of the Bo- dy is propor- tionate.	The denfity and speci- fic gravity of the Bo- dy.	The refra- Etive power of the Body in respect of its den- sity.
A Pfeudo-Topazius, be- ing a natural, pellucid, brittle, hairy Stone, of a vellow Colour	23 to	14	1'699	4'27	3979
Air Glafs of Antimony A Selenitis Glafs yulgar	3851 to 17 to 61 to 21 to	3850 9 41 20	0'00052 2'568 1'213 1'4025	0'00125 5'28 2'252 2'58	4160 4864 5386 5426
Cryftal of the Rock Ifland Cryftal Sal Gemmæ	25 to 5 to 17 to	16 3 11	1'445 1'778 1'388	2'65 2'72 2'143	5450 6536 - 6477
Borax Niter Dantzick Vitriol	22 to 32 to 303 to	15 21 200	1'1511 1'345 1'295	1'714 1'9 1'715	6716 7079 7551
Rain Water Gumm Arabic Spirit of Wine well recti-	10 to 529 to 31 to	7 396 21 72	1'041 0'7845 1'179 0'8765	1'7 1. 1'375	6124 7845 8574
fied Camphire Oyl Olive Lintfeed Oyl	3 to 22 to 40 to	/ 3 2 1 5 27	1 <sup>2</sup> 5 1 <sup>1</sup> 511 1 <sup>1</sup> 948	0'996 0'913 0'932	125 <b>5</b> 1 12607 12819
Spirit of Turpentine Ambar A Diamond	25 to 14 to 100 to	17 9 41	1'1626 1'42 4'949	0'874 1'04 3'4	13222 13654 14556

[73]

The refraction of the Air in this Table is determined by that of the Atmosphere observed by Astronomers. For if Light pass through many refracting substances or mediums gradually denser and denser, and terminated L l with with parallel furfaces, the fumm of all the refractions will be equal to the fingle refraction which it would have fuffered in paffing immediately out of the firft medium into the laft. And this holds true, though the number of the refracting fubftances be increased to infinity, and the distances from one another as much decreased, fo that the Light may be refracted in every point of its paffage, and by continual refractions bent into a curve Line. And therefore the whole refraction of Light in passing through the Atmosphere from the highest and rarest part thereof down to the lowest and denself part, must be equal to the refraction which it would suffer in passing at like obliquity out of a Vacuum immediately into Air of equal density with that in the lowest part of the Atmosphere.

Now, by this Table, the refractions of a Pfeudo-Topaz, a Selenitis, Rock Cryftal, Ifland Cryftal, Vulgar Glafs (that is, Sand melted together) and Glafs of Antimony, which are terreftrial ftony alcalizate concretes, and Air which probably arifes from fuch fubftances by fermentation, though these be fubftances very differing from one another in density, yet they have their refractive powers almost in the same proportion to one another as their densities are, excepting that the refraction of that ftrange fubftance Island-Cryftal is a little bigger than the reft. And particularly Air, which is 3400 times rarer than the Pseudo-Topaz, and 4200 times rarer than Glafs of Antimony, has notwithstanding its rarity the fame refractive power in respect of its density which those two very dense fubftances have in respect of theirs, excepting fo far as those two differ from one another.

Again,
Again, the refraction of Camphire, Oyl-Olive, Lintfeed Oyl, Spirit of Turpentine and Amber, which are fat fulphureous unctuous Bodies, and a Diamond, which probably is an unctuous fubftance coagulated, have their refractive powers in proportion to one another as their denfities without any confiderable variation. But the refractive power of these unctuous fubftances is two or three times greater in respect of their denfities than the refractive powers of the former fubftances in respect of theirs.

Water has a refractive power in a middle degree between those two forts of substances, and probably is of a middle nature. For out of it grow all vegetable and animal substances, which consist as well of substances fat and inflamable parts, as of earthy lean and alcalizate ones.

Salts and Vitriols have refractive powers in a middle degree between those of earthy substances and Water, and accordingly are composed of those two forts of subftances. For by distillation and rectification of their Spirits a great part of them goes into Water, and a great part remains behind in the form of a dry fixt earth capable of vitrification.

Spirit of Wine has a refractive power in a middle degree between those of Water and oyly substances, and accordingly seems to be composed of both, united by fermentation; the Water, by means of some faline Spirits with which 'tis impregnated, diffolving the Oyl, and volatizing it by the action. For Spirit of Wine is inflamable by means of its oyly parts, and being diftilled often from Salt of Tartar, grows by every diftillation more and more aqueous and flegmatick. And L1 2 Chymists Chymifts obferve, that Vegitables (as Lavender, Rue, Marjoram,  $\Im c.$ ) diftilled *per fe*, before fermentation yield Oyls without any burning Spirits, but after fermentation yield ardent Spirits without Oyls : Which fhews, that their Oyl is by fermentation converted into Spirit. They find alfo, that if Oyls be poured in fmall quantity upon fermentating Vegetables, they diftil over after fermentation in the form of Spirits.

So then, by the foregoing Table, all Bodies feem to have their refractive powers proportional to their denfities, (or very nearly;) excepting fo far as they partake more or lefs of fulphurous oyly particles, and thereby have their refractive power made greater or lefs. Whence it feems rational to attribute the refra-Etive power of all Bodies chiefly, if not wholly, to the fulphurous parts with which they abound. For it's probable that all Bodies abound more or lefs with Sulphurs. And as Light congregated by a Burning-glafs acts most upon fulphurous Bodies, to turn them into fire and flame; fo, fince all action is mutual, Sulphurs ought to act most upon Light. For that the action between Light and Bodies is mutual, may appear from this Confideration, That the denfeft Bodies which refract and reflect Light most strongly grow hottest in the Summer-Sun, by the action of the refracted or reflected Light.

I have hitherto explained the power of Bodies to reflect and refract, and shewed, that thin transparent plates, fibres and particles do, according to their several thicknesses and densities, reflect several sorts of rays, and thereby appear of several Colours, and by consequence that nothing more is requisite for producing all the the Colours of natural Bodies than the feveral fizes and denfities of their transparent particles. But whence it is that these plates, fibres and particles do, according to their feveral thickness and denfities, reflect several forts of rays, I have not yet explained. To give some infight into this matter, and make way for understanding the next Part of this Book, I shall conclude this Part with a few more Propositions. Those which preceded respect the nature of Bodies, these the nature of Light : For both must be understood before the reason of their actions upon one another can be known. And because the last Proposition depended upon the velocity of Light, I will begin with a Proposition of that kind.

# PROP. XI.

" Light is propagated from luminous Bodies in time, and Spends about seven or eight minutes of an hour in passing from the Sun to the Earth.

This was observed first by Romer, and then by others, by means of the Eclipses of the Satellites of Jupiter. For these Eclipses, when the Earth is between the Sun and Jupiter, happen about seven or eight minutes sooner than they ought to do by the Tables, and when the Earth is beyond the Sun they happen about seven or eight minutes later than they ought to do; the reason being, that the Light of the Satellites has farther to go in the latter case than in the former by the Diameter of the Earth's Orbit. Some inequalities of time may arise from the excentricities of the Orbs of the Satellites; but those cannot answer in all the Satellites, and at all times to. to the position and distance of the Earth from the Sun. The mean motions of *Jupiter*'s Satellites is also fwister in his descent from his Aphelium to his Perihelium, than in his ascent in the other half of his Orb : But this inequality has no respect to the position of the Earth, and in the three interior Satellites is insemble, as I find by computation from the Theory of their gravity.

#### PROP. XII.

\* Every ray of Light in its paffage through any refrasting surface is put into a certain transient constitution or state, which in the progress of the ray returns at equal intervals, and disposes the ray at every return to be easily transmitted through the next refracting surface, and between the returns to be easily reflected by it. M

This is manifeft by the 5th, 9th, 12th and 15th Obfervations. For by those Observations it appears, that one and the same fort of rays at equal Angles of incidence on any thin transparent plate, is alternately reflected and transmitted for many fuccess in arithmetical progression of the numbers 0, 1, 2, 3, 4, 5, 6, 7, 8,  $\Im c$ . So that if the first reflexion (that which makes the first or innermost of the Rings of Colours there described ) be made at the thickness 1, the rays shall be transmitted at the thickness 0, 2, 4, 6, 8, 10, 12,  $\Im c$ . and thereby make the central Spot and Rings of Light, which appear by transmission, and be reflected at the thickness 1, 3, 5, 7, 9, 11,  $\Im c$ . and thereby make the Rings which appear appear by reflexion. And this alternate reflexion and transmission, as I gather by the 24th Observation, continues for above an hundred vicifitudes, and by the the Observations in the next part of this Book, for many thousands, being propagated from one surface of a Glassplate to the other, though the thickness of the plate be a quarter of an Inch or above : So that this alternation seems to be propagated from every refracting furface to all distances without end or limitation.

This alternate reflexion and refraction depends on both the furfaces of every thin plate, because it depends on their diftance. By the 21th Observation, if either furface of a thin plate of Muscovy-Glass be wetted, the Colours caused by the alternate reflexion and refraction grow faint, and therefore it depends on them both.

It is therefore performed at the fecond furface, for if it were performed at the first, before the rays arrive at the second, it would not depend on the second.

It is alfo influenced by fome action or difpolition, propagated from the firft to the fecond, becaule otherwife at the fecond it would not depend on the firft. And this action or difpolition, in its propagation, intermits and returns by equal intervals, becaule in all its progrefs it inclines the ray at one diftance from the firft furface to be reflected by the fecond, at another to be transfmitted by it, and that by equal intervals for innumerable viciffitudes. And because the ray is disposed to reflexion at the distances 1, 3, 5, 7, 9,  $\Im c$ . and to transfmiffion at the distances 0, 2, 4, 6, 8, 10,  $\Im c$ , (for its transfmiffion through the first furface, is at the diftance thance  $\circ$ , and it is transmitted through both together, if their diffance be infinitely little or much less than 1) the disposition to be transmitted at the diffances 2, 4, 6, 8, 10,  $\Im c$ . is to be accounted a return of the fame disposition which the ray first had at the diffance $\circ$ , that is at its transmission through the first refracting furface. All which is the thing I would prove.

What kind of action or disposition this is? Whether it confift in 'a circulating or a vibrating motion of the ray, or of the medium, or fomething elfe? I do not here enquire. Those that are averse from assenting to any new difcoveries, but fuch as they can explain by an Hypothefis, may for the present suppose, that as Stones by falling upon Water put the Water into an undulating motion, and all Bodies by percuffion excite vibra-tions in the Air; fo the rays of Light, by impinging on any refracting or reflecting furface, excite vibrations in the refracting or reflecting medium or fubstance, and by exciting them agitate the folid parts of the refracting or reflecting Body, and by agitating them caufe the Body to grow warm or hot ; that the vibrations thus excited are propagated in the refracting or reflecting medium or fubstance, much after the manner that vibrations are propagated in the Air for caufing found, and move faster than the rays so as to overtake them; and that when any ray is in that part of the vibration which confpires with its motion, it eafily breaks through a refracting furface, but when it is in the contrary part of the vibration which impedes, its motion, it is eafily reflected; and, by confequence, that every ray is fucceffively difposed to be eafily reflected, or eafily transmitted, by every vibration which overtakes it. But whether

whether this Hypothesis be true or false I do not here confider. I content my felf with the bare discovery, that the rays of Light are by some cause or other alternately disposed to be reflected or refracted for many vicissifitudes.

# $\mathcal{D} E FINITION.$

The returns of the disposition of any ray to be reflected I will call its Fits of easy reflexion, and those of its disposition to be transmitted its Fits of easy transmission, and the space it passes between every return and the next return, the Interval of its Fits.

# PROP. XIII.

The reason why the surfaces of all thick transparent Bodies reflect part of the Light incident on them, and refract the rest, is, that some rays at their incidence are in Fits of easy reflexion, and others in Fits of easy transmission.

This may be gathered from the 24th Observation, where the Light reflected by thin plates of Air and Glass, which to the naked Eye appeared evenly white all over the plate, did through a Prism appear waved with many fucceffions of Light and Darkness made by alternate fits of easy reflexion and easy transmission, the Prism severing and distinguishing the waves of which the white reflected Light was composed, as was explained above.

And

And hence Light is in fits of eafy reflexion and eafy tranfiniffion, before its incidence on transparent Bodies. And probably it is put into fuch fits at its first emission from luminous Bodies, and continues in them during all its progress. For these fits are of a lasting Nature, as will appear by the next part of this Book.

In this Proposition I suppose the transparent Bodies to be thick, because if the thickness of the Body be much less than the interval of the fits of easy reflexion and transmission of the rays, the Body loseth its reflecting power. For if the rays, which at their entering into the Body are put into fits of easy transmission, arrive at the furthest furface of the Body before they be out of those fits they must be transmitted. And this is the reason why Bubbles of Water lose their reflecting power when they grow very thin, and why all opake Bodies when reduced into very small parts become transparent.

#### PROP. XIV.

Those surfaces of transparent Bodies, which if the ray be in a fit of refraction do refract it most strongly, if the ray be in a fit of reflexion do reflect it most easily.

For we shewed above in Prop. 8. that the cause of reflexion is not the impinging of Light on the folid impervious parts of Bodies, but some other power by which those folid parts act on Light at a distance. We shewed also in Prop. 9. that Bodies reflect and refract Light by one and the same power variously exercised in various circumstances, and in Prop. 1. that the most strongly refracting furfaces reflect the most Light : All which [83]

which compared together evince and ratify both this and the laft Proposition.

#### PROP. XV.

In any one and the same sort of rays emerging in any Angle out of any refracting surface into one and the same medium, the interval of the following fits of easy reflexion and transmission are either accurately or very nearly, as the Rectangle of the secant of the Angle of refraction, and of the secant of another Angle, whose sine is the first of 106 arithmetical mean proportionals, between the sines of incidence and refraction counted from the sine of refraction.

This is manifest by the 7th Observation.

#### PROP. XVI.

In feveral forts of rays emerging in equal Angles out of any refracting furface into the fame medium, the intervals of the following fits of eafy reflexion and eafy tranfmiffion are either accurately, or very nearly, as the Cuberoots of the Squares of the lengths of a Chord, which found the notes in an Eight, fol, la, fa, fol, la, mi, fa, fol, with all their intermediate degrees anfwering to the Colours of those rays, according to the Analogy described in the seventh Experiment of the second Book.

This is manifest by the 13th and 14th Observations.

Mm 2

PROP.

# [84]

#### PROP. XVII.

If rays of any one fort pass perpendicularly into several mediums, the intervals of the fits of easy reflexion and transmission in any one medium, is to those intervals in any other as the sine of incidence to the sine of refraction, when the rays pass out of the first of those two mediums into the second.

This is manifest by the 10th Observation.

#### PROP. XVIII.

If the rays which paint the Colour in the confine of yellow and orange pass perpendicularly out of any medium into Air, the intervals of their fits of easy reflexion are the  $\frac{1}{89000}$ th part of an Inch. And of the same length are the intervals of their fits of easy transmission.

This is manifest by the 6th Observation.

From these Propositions it is easy to collect the intervals of the fits of easy reflexion and easy transmisfion of any fort of rays refracted in any Angle into any medium, and thence to know, whether the rays shall be reflected or transmitted at their subsequent incidence upon any other pellucid medium. Which thing being useful for understanding, the next part of this Book was here to be set down. And for the same reason I add the two following Propositions.

PROP.

# [85]

# PROP. XIX.

If any fort of rays falling on the polite furface of any pellucid medium be reflected back, the fits of eafy reflexion which they have at the point of reflexion, shall still continue to return, and the returns shall be at distances from the point of reflexion in the arithmetical progression of the numbers 2, 4, 6, 8, 10, 12, &c. and between these fits the rays shall be in fits of eafy transmission.

For fince the fits of eafy reflexion and eafy tranfmiffion are of a returning nature, there is no reafon why these fits, which continued till the ray arrived at the reflecting medium, and there inclined the ray to reflexion, should there cease. And if the ray at the point of reflexion was in a fit of eafy reflexion, the progression of the distances of these fits from that point must begin from 0, and so be of the numbers 0, 2, 4, 6, 8,  $\Im c$ . And therefore the progression of the difances of the intermediate fits of easy transmission reckoned from the fame point, must be in the progression of the odd numbers 1, 3, 5, 7, 9,  $\Im c$ . contrary to what happens when the fits are propagated from points of refraction.

#### PROP. XX.

The intervals of the fits of easy reflexion and easy transmission, propagated from points of reflexion into any medium, are equal to the intervals of the like fits which the same rays would have, if refracted into the same medium

[86]

medium in Angles of refraction equal to their Angles of reflexion.

For when Light is reflected by the fecond furface of thin plates, it goes out afterwards freely at the first furface to make the Rings of Colours which appear by reflexion, and by the freedom of its egrefs, makes the Colours of these Rings more vivid and strong than those which appear on the other fide of the plates by the transmitted Light. The reflected rays are therefore in fits of eafy transmission at their egress; which would not always happen, if the intervals of the fits within the plate after reflexion were not equal both in length and number to their intervals before it. And this confirms also the proportions set down in the former Proposition. For if the rays both in going in and out at the first surface be in fits of eafy transmission, and the intervals and numbers of those fits between the first and fecond furface, before and after reflexion, be equal; the diftances of the fits of eafy transmission from either surface, must be in the fame progression after reflexion as before; that is, from the first furface which transmitted them, in the progression of the even numbers 0, 2, 4, 6, 8, 5c. and from the fecond which reflected them, in that of the odd numbers 1, 3, 5, 7, &c. But these two Propofitions will become much more evident by the Obfervations in the following part of this Book.

THE

# [87] THE SECOND BOOK OF OPTICKS. PART IV.

Observations concerning the Reflexions and Colours of thick transparent polished Plates.

Here is no Glass or Speculum how well foever polished, but, besides the Light which it refracts or reflects regularly, fcatters every way irregularly a faint Light, by means of which the polished furface, when illuminated in a dark Room by a beam of the Sun's Light, may be easily seen in all positions of the Eye. There are certain Phænomena of this scattered Light, which when I first observed them, seemed very strange and surprising to me. My Observations were as follows.

OBS.

# [88]

#### OBS. I.

The Sun shining into my darkened Chamber through a Hole i of an Inch wide, I let the intromitted beam of Light fall perpendicularly upon a Glass Speculum ground concave on one fide and convex on the other, to a Sphere of five Feet and eleven Inches Radius, and quick-filvered over on the convex fide. And holding a white opake Chart, or a Quire of Paper at the Center of the Spheres to which the Speculum was ground, that is, at the distance of about five Feet and eleven Inches from the Speculum, in fuch manner, that the beam of Light might pass through a little Hole made in the middle of the Chart to the Speculum, and thence be reflected back to the fame Hole : I observed upon the Chart four or five concentric Irifes or Rings of Colours, like Rain-bows, encompaffing the Hole much after the manner that those, which in the fourth and following Observations of the first part of this third Book appeared between the Object-Glaffes, encompassed the black Spot, but yet larger and fainter than those. These Rings as they grew larger and larger became diluter and fainter, to that the fifth was fcarce visible. Yet fometimes, when the Sun fhone very clear, there appeared faint Lineaments of a fixth and feventh. If the diftance of the Chart from the Speculum was much greater or much lefs than that of fix Feet, the Rings became dilute and vanished. And if the distance of the Speculum from the Window was much greater than that of fix Feet, the reflected beam of Light would be so broad at the diftance of fix Feet from the Speculum where the Rings appeared, the Valitude of g oeffex) ray being enlarged. appeared, as to obscure one or two of the innermost Rings.; And therefore I usually placed the Speculum at about fix Feet from the Window; fo that its Focus might there fall in with the center of its concavity at the Rings upon the Chart. And this posture is always to be understood in the following Observations where no other is exprest.

## OBS. II.

The Colours of these Rain-bows fucceeded one another from the center outwards, in the fame form and order with those which were made in the ninth Observation of the first Part of this Book by Light not reflected, but transmitted through the two Object-Glasses. For, first, there was in their common center a white round Spot of faint Light, something broader than the reflected beam of Light; which beam sometimes fell upon the middle of the Spot, and sometimes by a little inclination of the Speculum receded from the middle, and left the Spot white to the center.

This white Spot was immediately encompafied with a dark grey or ruffet, and that darknefs with the Colours of the first Iris, which were on the infide next the <u>darknels</u> a little violet and indico, and next to that a blue, which on the outfide grew pale, and then fucceeded a little greenish yellow, and after that a brighter yellow, and then on the outward edge of the Iris a red which on the outfide inclined to purple.

This Iris was immediately encompassed with a fecond, whose Colours were in order from the infide outwards, purple, blue, green, yellow, light red, a red mixed with purple.

Then immediately followed the Colours of the third Iris, which were in order outwards a green inclining to purple, a good green, and a red more bright than that of the former Iris.

The fourth and fifth Iris feemed of a bluifh green within, and red without, but fo faintly that it was difficult to difcern the Colours.

#### O B S. III.

Meafuring the Diameters of these Rings upon the Chart as accurately as I could, I found them also in the fame proportion to one another with the Rings made by Light transmitted through the two Object-Glasses. For the Diameters of the four first of the bright Rings meafured between the brightest parts of their orbits, at the distance of fix Feet from the Speculum were 1<sup>11</sup><sub>16</sub>, 2<sup>3</sup>/<sub>8</sub>, 2<sup>11</sup>/<sub>12</sub>, 3<sup>3</sup>/<sub>8</sub> Inches, whose squares are in arithmetical progression of the numbers 1, 2, 3, 4. If the white circular Spot in the middle be reckoned amongst the Rings, and its central Light, where it feems to be most luminous, be put equipollent to an infinitely little Ring; the squares of the Diameters of the Rings will be in the progression 0, 1, 2, 3, 4, 5c. I measured also the Diameters of the dark Circles between these luminous ones, and found their squares in the progression of the numbers  $\frac{1}{2}$ ,  $1\frac{1}{2}$ ,  $2\frac{1}{2}$ ,  $3\frac{1}{2}$ , 5c. the Diameters of the first four at the distance of fix Feet from the Speculum, being  $1_{16}^3$ ,  $2_{16}^1$ ,  $2_{3}^2$ ,  $3_{20}^2$  Inches. If the diftance of the Chart from the Speculum was increafed

creafed or diminished, the Diameters of the Circles were increased or diminished proportionally.

## OBS. IV.

By the analogy between these Rings and those defcribed in the Observations of the first Part of this Book, I support of the there were many more of them which spread into one another, and by interfering mixed their Colours, and diluted one another so that they could not be seen apart. I viewed them therefore through a Prism, as I did those in the 24th Observation of the first Part of this Book. And when the Prism was so placed as by refracting the Light of their mixed Colours to separate them, and distinguish the Rings from one another, as it did those in that Observation, I could then see them distincter than before, and easily number eight or nine of them, and sometimes twelve or thirteen. And had not their Light been so very faint, I question not but that I might have seen many more.

# . O B S. V.

Placing a Prifm at the Window to refract the intromitted beam of Light, and caft the oblong Spectrum of Colours on the Speculum : I covered the Speculum with a black Paper which had in the middle of it a Hole to let any one of the Colours pass through to the Speculum, whils the rest were intercepted by the Paper. And now I found Rings of that Colour only which fell upon the Speculum. If the Speculum was illuminated with red the Rings were totally red with dark inter-N n 2 vals, vals, if with blue they were totally blue, and io of the other Colours. And when they were illuminated with any one Colour, the Squares of their Diameters meafured between their most luminous parts, were in the arithmetical progression of the numbers 0, 1, 2, 3, 4, and the Squares of the Diameters of their dark intervals in the progression of the intermediate numbers  $\frac{1}{2}$ ,  $I_{\frac{1}{2}}^{1}$ ,  $2\frac{1}{2}^{1}$ ,  $3\frac{1}{2}^{1}$ : But if the Colour was varied they varied their magnitude. In the red they were largest, in the indico and violet least, and in the intermediate Colours yellow, green and blue; they were of feveral intermediate bigneffes answering to the Colour, that is, greater in yellow than in green, and greater in green than in blue. And hence I knew that when the Speculum was illuminated with white Light, the red and yellow on the outfide of the Rings were produced by the least refrangible rays, and the blue and violet by the most refrangible, and that the Colours of each Ring spread into the Colours of the neighbouring Rings on either fide, after the manner explained in the first and second Part of this Book, and by mixing diluted one another fo that they could not be diftinguished, unless near the center where they were leaft mixed. For in this Observation I could fee the Rings more diffinctly, and to a greater number than before, being able in the yellow Light to number eight or nine of them, befides a faint shadow of a tenth. To fatisfy my felf how much the Colours of the feveral Rings fpread into one another, I meafured the Diameters of the fecond and third Rings, and found them when made by the confine of the red and orange to be the fame Diameters when made by the confine of blue and indico, as 9 to 8, or thereabouts. For it was hard to

to determine this proportion accurately. Alfo the Circles made fucceffively by the red, yellow and green, differed more from one another than those made fucces fively by the green, blue and indico. For the Circle made by the violet was too dark. to be feen. To carry on the computation, Let us therefore fuppose that the differences of the Diameters of the Circles made by the outmost red, the confine of red and orange, the confine of orange and yellow, the confine of yellow and green, the confine of green and blue, the confine of blue and indico, the confine of indico and violet, and outmost violet, are in proportion as the differences of the lengths of a Monochord which found the tones in an Eight; fol, la, fa, fol, la, mi, fa, fol, that is, as the numbers  $\frac{1}{99}$  $\frac{1}{18}$ ,  $\frac{1}{12}$ ,  $\frac{1}{27}$ ,  $\frac{1}{27}$ ,  $\frac{1}{27}$ ,  $\frac{1}{18}$ . And if the Diameter of the Circle made by the confine of red and orange be 9 A, and that of the Circle made by the confine of blue and indico be 8 A as above, their difference 9 A ---- 8 A will be to the difference of the Diameters of the Circles made by the outmost red, and by the confine of red and orange, as  $\frac{1}{18} + \frac{1}{12} + \frac{1}{2} + \frac{2}{27}$  to  $\frac{1}{9}$ , that is as  $\frac{8}{27}$  to  $\frac{1}{9}$  or 8 to 3, and to the difference of the Circles made by the outmost violet, and by the confine of blue and indico, as  $\frac{1}{18} + \frac{1}{12} + \frac{1}{12} + \frac{2}{37}$ to  $\frac{1}{27} + \frac{1}{18}$ , that is, as  $\frac{5}{27}$  to  $\frac{5}{54}$ , or as 16 to 5. And therefore these differences will be & A and & A. Add the first to 9 A and subduct the last from 8 A, and your will have the Diameters of the Circles made by the leaft and most refrangible rays  $\frac{2}{8}$  A and  $\frac{61}{8}$  A. These Diameters are therefore to one another as 75 to  $61\frac{1}{2}$  or 50 to 41, and their Squares as 2500 to 1681, that is, as 3 to 2 very nearly. Which proportion differs not. much from the proportion of the Diameters of the Circles

Circles made by the outmost red and outmost violet in the 13th Observation of the first part of this Book.

#### OBS. VI.

Placing my Eye where these Rings appeared plainest, I faw the Speculum tinged all over with waves of Colours ( red, yellow, green, blue ; ) like those which in the Observations of the first Part of this Book appeared. between the Object-Glaffes and upon Bubbles of Water, but much larger. And after the manner of those, they were of various magnitudes in various politions of the Eye, fwelling and fhrinking as I moved my Eye this way and that way. They were formed like Arcs of concentrick Circles as those were, and when my Eye was over against the center of the concavity of the Speculum (that is, 5 Feet and 10 Inches distance from the Speculum) their common center was in a right Line with that center of concavity, and with the Hole in the Window. But in other postures of my Eye their center had other politions. They appeared by the Light of the Clouds propagated to the Speculum through the Hole in the Window, and when the Sun shone through that Hole upon the Speculum, his Light upon it was of the Colour of the Ring whereon it fell, but by its fplendor obfcured the Rings made by the Light of the Clouds, unlefs when the Speculum was removed to a great distance from the Window, so that his Light upon it might be broad and faint. By varying the polition of my Eye, and moving it nearer to or farther from the direct beam of the Sun's Light, the Colour of the Sun's reflected Light constantly varied upon the Speculum, as

as it did upon my Eye, the fame Colour always appearing to a By-ftander upon my Eye which to me appeared upon the Speculum. And thence I knew that the Rings of Colours upon the Chart were made by thefe reflected Colours propagated thither from the Speculum in feveral Angles, and that their production depended not upon the termination of Light and Shaddow.

## OBS. VII.

By the Analogy of all these Phænomena with those of the like Rings of Colours described in the first Part of this Book, it feemed to me that these Colours were produced by this thick plate of Glass, much after the manner that those were produced by very thin For, upon tryal, I found that if the Quickplates. filver were rubbed off from the back-fide of the Speculum, the Glais alone would caufe the fame Rings of Colours, but much more faint than before ; and therefore the Phænomenon depends not upon the Quickfilver, unlefs fo far as the Quick-filver by the increasing, the reflexion of the back-fide of the Glais increases the Light of the Rings of Colours. I found also that a Speculum of metal without Glass made some years fince for optical uses, and very well wrought, produced none of those Rings; and thence I understood that these Rings arife not from one specular surface alone, but depend upon the two furfaces of the plate of Glass whereof the Speculum was made, and upon the thickness of the Glass between them. For as in the 7th and 19th Observations of the first Part of this Book a thin plate of of Air, Water, or Glass of an even thickness appeared of one Colour when the rays were perpendicular to it, of another when they were a little oblique, of another when more oblique, of another when still more oblique, and fo on ; fo here, in the fixth Obfervation, the Light which emerged out of the Glass in several obliquities, made the Glass appear of several Colours, and being propagated in those obliquities to the Chart, there painted Rings of those Colours. And as the reason why a thin plate appeared of feveral Colours in feveral obliquities of the rays, was, that the rays of one and the fame fort are reflected by the thin plate at one obliquity and transmitted at another, and those of other forts transmitted where these are reflected, and reflected where thefe are transmitted : So the reason why the thick plate of Glass whereof the Speculum was made did appear of various Colours in various obliquities, and in those obliquities propagated those Colours to the Chart, was, that the rays of one and the fame fort did at one obliquity emerge out of the Glass, at another did not emerge but were reflected back towards the Quick-filver by the hither furface of the Glafs, and accordingly as the obliquity became greater and greater emerged and were reflected alternately for many fucceffions, and that in one and the same obliquity the rays of one fort were reflected, and those of another transmitted. This is manifest by the first Observation of this Book : For in that Obfervation, when the Speculum was illuminated by any one of the prismatick Colours, that Light made many Rings of the same Colour upon the Chart with dark intervals, and therefore at its emergence out of the Speculum was alternately transmitted, and not tranf-

transmitted from the Speculum to the Chart for many fucceffions, according to the various obliquities of its emergence. And when the Colour caft on the Speculum by the Prifm was varied, the Rings became of the Colour caft on it, and varied their bignefs with their Colour, and therefore the Light was now alternately transmitted and not transmitted from the Speculum to the Lens at other obliquities than before. It feemed to me therefore that these Rings were of one and the fame original with those of thin plates, but yet with this difference that those of thin plates are made by the alternate reflexions and transmissions of the rays at the fecond furface of the plate after one paffage through it : But here the rays go twice through the plate before they are alternately reflected and transmitted; first, they go through it from the first surface to the Quickfilver, and then return through it from the Quick-filver to the first furface, and there are either transmitted to the Chart or reflected back to the Quick-filver, accordingly as they are in their fits of easie reflexion or transmission when they arrive at that furface. For the intervals of the fits of the rays which fall perpendicularly on the Speculum, and are reflected back in the fame perpendicular Lines, by reason of the equality of these Angles and Lines, are of the same length and number within the Glass after reflexion as before by the 19th Proposition of the third Part of this Book. And therefore fince all the rays that enter through the first furface are in their fits of eafy transmission at their entrance, and as many of these as are reflected by the fecond are in their fits of eafy reflexion there, all thefe must be again in their fits of easy transmission at their O oreturn

10.3

return to the first, and by consequence there go out of the Glafs to the Chart, and form upon it the white Spot of Light in the center of the Rings. For the reafon holds good in all forts of rays, and therefore all forts must go out promiscuously to that Spot, and by their mixture caule it to be white. But the intervals of the fits of those rays which are reflected more obliquely than they enter, must be greater after reflexion than before by the 15th and 20th Prop. And thence it may happen that the rays at their return to the first furface, may in certain obliquities be in fits of eafy reflexion, and return back to the Quick-filver, and in other intermediate obliquities be again in fits of eafy transmission, and so go out to the Chart, and paint on it the Rings of Colours about the white Spot. And because the intervals of the fits at equal obliquities are greater and fewer in the lefs refrangible rays, and lefs and more numerous in the more refrangible, therefore the less refrangible at equal obliquities shall make fewer Rings than the more refrangible, and the Rings made by those shall be larger than the like number of Rings made by thefe; that is, the red Rings shall be larger than the yellow, the yellow than the green, the green than the blue, and the blue than the violet, as they were really found to be in the 5th Observation. And therefore the first Ring of all Colours incompassing the white Spot of Light shall be red without and violet within, and yellow, and green, and blue in the middle, as it was found in the fecond Obfervation; and thefe Colours in the fecond Ring, and those that follow shall be more expanded till they fpread into one another, and blend one another by interfering.

Thefe

These seems to be the reasons of these Rings in general, and this put me upon observing the thickness of the Glass, and confidering whether the dimensions and proportions of the Rings may be truly derived from it by computation.

## OBS. VIII.

I measured therefore the thickness of this concavoconvex plate of Glass, and found it every-where ‡ of an Inch precifely. Now, by the 6th Obfervation of the first Part of this Book, a thin plate of Air transmits the brighteft Light of the first Ring, that is the bright yellow, when its thickness is the spood th part of an Inch, and by the 10th Observation of the same part, a thin plate of Glass transmits the same Light of the same Ring when its thickness is less in proportion of the fine of refraction to the fine of incidence, that is, when its thickness is the TI the or TITE the part of an Inch, fuppofing the fines are as 11 to 17. And if this thickness be doubled it transmits the same bright Light of the fecond Ring, if tripled it transmits that of the third, and so on, the bright yellow Light in all these cases being in its fits of transmission. And therefore if its thicknefs be multiplied 34386 times fo as to become  $\frac{1}{4}$  of an Inch it transmits the same bright Light of the 34386th Ring. Suppose this be the bright yellow Light transmitted perpendicularly from the reflecting convex fide of the Glais through the concave fide to the white Spot in the center of the Rings of Colours on the Chart : And by a rule in the feventh Observation in the first Part of the first Book, and by the 15th and 20th Propositions  $0_{0_2}$ of

3

# [ 100 ]

of the third Part of this Book, if the rays be made ob-lique to the Glass, the thickness of the Glass requifite to transmit the same bright Light of the same Ring in any obliquity is to this thickness of  $\frac{1}{4}$  of an Inch, as the secant of an Angle whole line is the first of an hundred and fix arithmetical means between the fines of incidence and refraction, counted from the fine of incidence when the refraction is made out of any plated Body into any medium incompaffing it, that is, in this cafe, out of Glafs into Air. Now if the thicknefs of the Glafs be increased by degrees, so as to bear to its first thickness, (viz. that of a quarter of an Inch ) the proportions which 34386 (the number of fits of the perpendicular rays in going through the Glass towards the white Spot in the center of the Rings,) hath to 34385, 34384, 34383 and 34382 (the numbers of thefits of the oblique rays in going through the Glafs towards the first, fe-cond, third and fourth Rings of Colours,) and if the first thickness be divided into 100000000 equal parts, the increased thickness will be 100002908, 100005816, the increated thicknehes will be record, the property increased the increased the increased will be 26' 13", 37' 5", 45' 6" and 52' 26", the Radius being 100000000; and the fines of these Angles are 762, 1079, 1321 and 1525, and the proportional fines of refraction 1172, 1659, 2031 and 2345, the Radius being 100000. For fince the fines of incidence out of Glass into Air are to the fines of refraction as 11 to 17, and to the above-mentioned fecants as 11 to the first of 106 arithmetical means between 11 and 17, that is as 11 to  $11\frac{6}{106}$ , those fe-cants will be to the fines of refraction as  $11\frac{6}{106}$  to 17, and by this Analogy will give these fines. So then if

# [ 101 ]

if the obliquities of the rays to the concave furface of. the Glass be such that the fines of their refraction inpassing out of the Glass through that furface into the. Air be 1172, 1659, 2031, 2345, the bright Light of the 34386th Ring shall emerge at the thicknesses of the Glafs which are to ½ of an Inch as 34386 to 34385, 34384, 34383, 34382, respectively. And therefore if the thickness in all these cases be  $\frac{1}{4}$  of an Inch. (as it is in the Glass of which the Speculum was made) the bright Light of the 34385th Ring shall emerge where the fine. of refraction is 1172, and that of the 34384th, 384383th and 34382th Ring where the fine is 1659, 2031, and 2345 respectively. And in these Angles of refraction. the Light of these Rings shall be propagated from the. Speculum to the Chart, and there paint Rings about the. white central round Spot of Light which we faid was the Light of the 34386th Ring. And the Semidiameters of these Rings shall subtend the Angles of refraction made at the concave furface of the Speculum, and by confequence their Diameters shall be to the distance of the Chart from the Speculum as those fines of refraction. doubled are to the Radius that is as 1172, 1659, 2031, and 2245, doubled are to 100000. And therefore if. the diftance of the Chart from the concave furface of the Speculum be fix Feet (as it was in the third of thefe. Observations) the Diameters of the Rings of this bright yellow Light upon the Chart shall be 1'688, 2'389, 2'925, 3'375 Inches: For these Diameters are to 6 Feet as the above-mentioned fines doubled are to the Radius. Now these Diameters of the bright yellow Rings, thus found by computation are the very fame with those found in the third of these Observations by measuring them,

them, (viz. with 111, 23, 211, and 33 Inches, and therefore the Theory of deriving these Rings from the thicknefs of the plate of Glass of which the Speculum was made, and from the obliquity of the emerging rays agrees with the Observation. In this computation I have equalled the Diameters of the bright Rings made by Light of all Colours, to the Diameters of the Rings made by the bright yellow. For this yellow makes the brightest part of the Rings of all Colours. If you defire the Diameters of the Rings made by the Light of any other unmixed Colour, you may find them readily by putting them to the Diameters of the bright yellow ones in a fubduplicate proportion of the intervals of the fits of the rays of those Colours when equally inclined to the refracting or reflecting furface which caused those fits, that is, by putting the Diameters of the Rings made by the rays in the extremities and limits of the feven Colours, red, orange, yellow, green, blue, indico, violet, proportional the Cube-roots of the numbers,  $1, \frac{8}{9}, \frac{5}{6}, \frac{3}{4}$ pretty nearly in the fame proportion to one another, which they ought to have by the fifth of these Observations.

And thus I fatisfied my felf that these Rings were of the fame kind and original with those of thin plates, and by consequence that the fits or alternate dispositions of the rays to be reflected and transmitted are propagated to great distances from every reflecting and refracting furface. But yet to put the matter out of doubt 4 added the following Observation.

OBS.

# [103]

# OBS. IX.

If these Rings thus depend on the thickness of the plate of Glass their Diameters at equal distances from several Speculums made of fuch concavo-convex plates of Glafs as are ground on the fame Sphere, ought to be reciprocally in a subduplicate proportion of the thicknesses of the plates of Glass. And if this proportion be found true by experience it will amount to a demonstration that these Rings (like those formed in thin plates) do depend on the thickness of the Glass. I procured therefore another concavo-convex plate of Glass ground on both fides to the fame Sphere with the former plate : Its thickness was  $\frac{5}{62}$  parts of an Inch; and the Diameters of the three first bright Rings measured between the brightest parts of their orbits at the distance of 6 Feet from the Glass were 3. 4<sup>t</sup>/<sub>6</sub>. 5<sup>t</sup>/<sub>8</sub>. Inches. Now the thicknefs of the other Glafs being  $\frac{1}{4}$  of an Inch was to thicknels of this Glass as  $\frac{1}{4}$  to  $\frac{1}{62}$ , that is as 31 to 10, or 310000000 to 100000000, and the roots of these numbers. are 17607 and 10000, & in the proportion of the first of these roots to the second are the Diameters of the bright Rings made in this Observation by the thinner Glass, 3.  $4\frac{1}{6}$ .  $5\frac{1}{8}$  to the Diameters of the fame Rings made in the third of these Observations by the thicker Glass. 116. 23 21, that is, the Diameters of the Rings are reciprocally in a fubduplicate proportion of thickneffes of the plates of Glafs.

So then in plates of Glass which are alike concave on one fide, and alike convex on the other fide, and alike quick-filvered on the convex fides, and differ in nothing but

but their thickness, the Diameters of the Rings are reciprocally in a fubduplicate proportion of the thickneffes of the plates. And this fhews fufficiently that the Rings depend on both the surfaces of the Glass. They depend on the convex furface because they are more luminous when that furface is quick-filvered over than when it is without Quick-filver. They depend alfo upon the concave furface, becaufe without that furface a Speculum makes them not. They depend on both furfaces and on the distances between them, because their bigness is varied by varying only that distance. And this dependance is of the fame kind with that which the Colours of thin plates have on the diffance of the furfaces of those plates, because the bigness of the Rings and their proportion to one another, and the variation of their bigness arising from the variation of the thickness of the Glass, and the orders of their Colours, is fuch as ought to refult from the Propo-fitions in the end of the third Part of this Book, derived from the the Phænomena of the Colours of thin plates set down in the first Part.

There are yet other Phænomena of these Rings of Colours but such as follow from the same Propositions, and therefore confirm both the truth of those Propositions, and the Analogy between these Rings and the Rings of Colours made by very thin plates. I shall subjoyn some of them.

OBS:

# [104]

# [105]

## OBS. X.

When the beam of the Sun's Light was reflected back from the Speculum not directly to the Hole in the Window, but to a place a little diftant from it, the common center of that Spot, and of all the Rings of Colours fell in the middle way between the beam of the incident Light, and the beam of the reflected Light, and by consequence in the center of the spherical concavity of the Speculum, whenever the Chart on which the Rings of Colours fell was placed at that center. And as the beam of reflected Light by inclining the Speculum receded more and more from the beam of incident Light and from the common center of the coloured Rings between them, those Rings grew bigger and bigger, and fo alfo did the white round Spot, and new Rings of Colours emerged fucceffively out of their common center, and the white Spot became a white Ring encompaffing them; and the incident and reflected beams of Light always fell upon the opposite parts of this Ring, illuminating its perimeter like two mock Suns in the oppolite parts of an Iris. So then the Diameter of this Ring, measured from the middle of its Light on one fide to the middle of its Light on the other fide, was always equal to the diftance between the middle of the incident beam of Light, and the middle of the reflected beam meafured at the Chart on which the Rings appeared: And the rays which formed this Ring were reflected by the Speculum in Angles equal to their Angles of incidence, and by contequence to their Angles of refraction at their entrance into the Glass, but yet their Angles of reflexion Pp

reflexion were not in the fame planes with their Angles of incidence.

#### OBS. XI.

The Colours of the new Rings were in a contrary order to those of the former, and arose after this manner. The white round Spot of Light in the middle of the Rings continued white to the center till the distance of the incident and reflected beams at the chart was about  $\frac{2}{8}$  parts of an Inch, and then it began to grow dark in the middle. And when that distance was about  $1\frac{1}{16}$  of an Inch, the white Spot was become a Ring encompassing a dark round Spot which in the middle inclined to violet and indico. And the luminous Rings incompassing it were grown equal to those dark ones which in the four first Observations encompassed them, that is to fay, the white Spot was grown a white Ring equal to the first of those dark Rings, and the first of those luminous Rings was now grown equal to the fecond of those dark ones, and the second of those luminous ones to the third of those dark ones, and fo on. For the Diameters of the luminous Rings were now  $1\frac{3}{16}$ ,  $2\frac{1}{3}$ ,  $3\frac{3}{20}$ , 5c. Inches.

When the diftance between the incident and reflected beams of Light became a little bigger, there emerged out of the middle of the dark Spot after the indico a blue, and then out of that blue a pale green, and foon after a yellow and red. And when the Colour at the center was brighteft, being between yellow and red, the bright Rings were grown equal to those Rings which in the four first Observations next encompassed them; that that is to fay, the white Spot in the middle of those Rings was now become a white Ring equal to the first of those bright Rings, and the first of those bright ones was now become equal to the fecond of those, and so on. For the Diameters of the white Rings, and of the other luminous Rings incompassing it, were now  $1\frac{1}{7}$ ,  $2\frac{1}{8}$ ,  $2\frac{1}{2}$ ,  $3\frac{1}{8}$ ,  $\Im c$ . or thereabouts.

When the diftance of the two beams of Light at the Chart was a little more increased, there emerged out of the middle in order after the red, a purple, a blue, a green, a yellow, and a red inclining much to purple, and when the Colour was brightest being between yellow and red, the former indico, blue, green, yellow and red, were become an Iris or Ring of Colours equal to the first of those luminous Rings which appeared in the four first Observations, and the white Ring which was now become the fecond of the luminous Rings was grown equal to the second of those, and the first of those which was now become the third Ring was become the third of those, and so on. For their Diameters were  $1^{\frac{1}{16}}$ ,  $2^{\frac{3}{8}}$ ,  $2^{\frac{1}{12}}$ ,  $3^{\frac{3}{8}}$  Inches, the diftance of the two beams of Light, and the Diameter of the white Ring being 23 Inches.

When thefe two beams became more diffant there emerged out of the middle of the purplifh red, firft a darker round Spot, and then out of the middle of that Spot a brighter. And now the former Colours (purple, blue, green, yellow, and purplifh red) were become a Ring equal to the firft of the bright Rings mentioned in the four firft Obfervations, and the Ring about this Ring were grown equal to the Rings about that refpectively; the diffance between the two beams of P p 2 Light Light and the Diameter of the white Ring (which was now become the third Ring) being about 3 Inches.

The Colours of the Rings in the middle began now to grow very dilute, and if the diftance between the two beams was increafed half an Inch, or an Inch more, they vanifhed whilft the white Ring, with one or two of the Rings next it on either fide, continued ftill vifible. But if the diftance of the two beams of Light was ftill more increafed thefe alfo vanifhed : For the Light which coming from feveral parts of the Hole in the Window fell upon the Speculum in feveral Angles of incidence made Rings of feveral bigneffes, which diluted and blotted out one another, as I knew by intercepting fome part of that Light. For if I intercepted that part which was neareft to the Axis of the Speculum the Rings would be lefs, if the other part which was remoteft from it they would be bigger.

#### OBS. XII.

When the Colours of the Prifm were caft fucceffively on the Speculum, that Ring which in the two laft Obfervations was white, was of the fame bignefs in all the Colours, but the Rings without it were greater in the green than in the blue, and ftill greater in the yellow, and greateft in the red. And, on the contrary, the Rings within that white Circle were lefs in the green than in the blue, and ftill lefs in the yellow, and leaft in the red. For the Angles of reflexion of those rays which made this Ring being equal to their Angles of incidence, the fits of every reflected ray within the Glafs after

after reflexion are equal in length and number to the fits of the fame ray within the Glass before its incidence on the reflecting furface; and therefore fince all the rays of all forts at their entrance into the Glass were in a fit of transmission, they were also in a fit of transmission at their returning to the fame furface after reflexion; and by confequence were transmitted and went out to the white Ring on the Chart. This is the reason why that Ring was of the fame bignefs in all the Colours, and why in a mixture of all it appears white. But in rays which are reflected in other Angles, the intervals of the fits of the least refrangible being greatest, make the Rings of their Colour in their progress from this white Ring, either outwards or inwards, increase or decrease by the greatest steps; so that the Rings of this Colour without are greatest, and within least. And this is the reafon why in the last Observation, when the Speculum was illuminated with white Light, the exterior Rings made by all Colours appeared red without and blue within, and the interior blue without and red within.

Thefe are the Phænomena of thick convexo-concave plates of Glafs, which are every where of the fame thicknefs. There are yet other Phænomena when thefe plates are a little thicker on one fide than on the other, and others when the plates are more or lefs concave than convex, or plano-convex, or double-convex. For in all thefe cafes the plates make Rings of Colours, but after various manners ; all which, fo far as I have yet obferved, follow from the Propositions in the end of the third part of this Book, and fo confpire to confirm the truth of those Propositions. But the Phænomena. mena are too various, and the Calculations whereby they follow from those Propositions too intricate to be here profecuted. I content my felf with having profecuted this kind of Phænomena so far as to discover their cause, and by discovering it to ratify the Propositions in the third Part of this Book.

#### OBS. XIII.

As Light reflected by a Lens quick-filvered on the back-fide makes the Rings of Colours above defcribed, fo it ought to make the like Rings of Colours in paffing through a drop of Water. At the first reflexion of the rays within the drop, fome Colours ought to be transmitted, as in the case of a Lens, and others to be reflected back to the Eye. For inftance, if the Diameter of a finall drop or globule of Water be about the 500th part of an Inch, fo that a red-making ray in paffing through the middle of this globule has 250 fits of eafy transmission within the globule, and that all the red-making rays which are at a certain diftance from this middle ray round about it have 249 fits within the globule, and all the like rays at a certain further diftance round about it have 248 fits, and all those at a certain further diftance 24.7 fits, and fo on ; these concentrick Circles of rays after their transmission, falling on a white Paper, will make concentrick rings of red upon the Paper, fuppoing the Light which paffes through one fingle globule strong enough to be sensible. And, in like manner, the rays of other Colours will make Rings of other Colours. Suppose now that in a fair day the Sun shines through a thin Cloud of such globules
globules of Water or Hail, and that the globules are all of the fame bigness, and the Sun feen through this Cloud shall appear incompassed with the like concentrick Rings of Colours, and the Diameter of the first Ring of red shall be 7<sup>2</sup>/<sub>4</sub> degrees, that of the fecond 10<sup>2</sup>/<sub>7</sub> degrees, that of the third 12 degrees 33 minutes. And accordingly as the globules of Water are bigger or lefs, the Rings shall be less or bigger. This is the Theory, and experience answers it. For in June 1692. I faw by reflexion in a Veffel of stagnating Water three Halos Crowns or Rings of Colours about the Sun, like three little Rainbows, concentrick to his Body. The Colours of the first or innermost Crown were blue next the Sun, red without, and white in the middle between the blue and red. Those of the second Crown were purple and blue within, and pale red without, and green in the middle. And those of the third were pale blue within, and pale red without; these Crowns inclosed one another immediately, fo that their Colours proceeded in this continual order from the Sun outward : blue, white, red; purple, blue, green, pale yellow and red; pale blue, pale red. The Diameter of the fecond Crown measured from the middle of the yellow and red on one fide of the Sun, to the middle of the fame Colour on the other fide was  $9\frac{1}{3}$  degrees, or thereabouts. The Dia-meters of the first and third I had not time to measure, but that of the first seemed to be about five or fix degrees, and that of the third about twelve. The like Crowns appear fometimes about the Moon; for in the beginning of the year 1664, Febr. 19th at night, I faw two fuch Crowns about her. The Diameter of the first or innermost was about three degrees, and that of the fecond

fecond about five degrees and an half. Next about the Moon was a Circle of white, and next about that the inner Crown which was of a bluish green within next the white, and of a yellow and red without, and next about these Colours were blue and green on the infide of the outward Crown, and red on the outfide of it. At the fame time there appeared a Halo about 22 degrees 35' distant from the center of the Moon. It was Elliptical, and its long Diameter was perpendicular to the Horizon verging below fartheft from the Moon. I am told that the Moon has fometimes three or more concentrick Crowns of Colours incompaffing one another next about her Body. The more equal the globules of Water or Ice are to one another, the more Crowns of Colours will appear, and the Colours will be the more lively. The Halo at the diftance of 22<sup>1</sup>/<sub>2</sub> degrees from the Moon is of another fort. By its being oval and remoter from the Moon below than above, I conclude, that it was made by refraction in fome fort of Hail or Snow floating in the Air in an horizontal Posture, the refracting Angle being about 58 or 60 degrees.

THE

# [113]

# THE THIRD BOOK OF OPTICKS

#### Observations concerning the Inflexions of the rays of Light, and the Colours made thereby.

**G** Rimaldo has informed us, that if a beam of the Sun's Light be let into a dark Room through a very fmall Hole, the fhadows of things in this Light will be larger than they ought to be if the rays went on by the Bodies in ftreight Lines, and that thefe fhadows have three parallel fringes, bands or ranks of coloured Light adjacent to them. But if the Hole be enlarged the fringes grow broad and run into one another, fo that they cannot be diftinguifhed. Thefe broad fhadows and fringes have been reckoned by fome to proceed from the ordinary refraction of the Air, but without due examination of the matter. For the circumftances of the Phænomenon, fo far as I have obferved them, are as follows.

OBS.

# [114]

#### OBS. I.

I made in a piece of Lead a finall Hole with a Pin, whose breadth was the 42th part of an Inch. For 21 of those Pins laid together took up the breadth of half an Inch. Through this Hole I let into my darkened Chamber a beam of the Sun's Light, and found that the shadows of Hairs, Thred, Pins, Straws, and fuch like slender fubstances placed in this beam of Light, were confiderably broader than they ought to be, if the rays of Light paffed on by these Bodies in right Lines. And particularly a Hair of a Man's Head, whole breadth was but the 280th part of an Inch, being held in this Light, at the diftance of about twelve Feet from the Hole, did caft a shadow which at the distance of four Inches from the Hair was the fixtieth part of an Inch broad, that is, above four times broader than the Hair, and at the diftance of two Feet from the Hair was about the eight and twentieth part of an Inch broad, that is, ten times broader than the Hair, and at the distance of ten Feet was the eighth part of an Inch broad, that is 35 times broader.

Nor is it material whether the Hair be incompafied with Air, or with any other pellucid fubftance. For I wetted a polifhed plate of Glafs, and laid the Hair in the Water upon the Glafs, and then laying another polifhed plate of Glafs upon it, fo that the Water might fill up the fpace between the Glaffes, I held them in the aforefaid beam of Light, fo that the Light might pafs through them perpendicularly, and the fhadow of the Hair was at the tame diftances as big as before. The The fhadows of fcratches made in polifhed plates of Glafs were alfo much broader than they ought to be, and the Veins in polifhed plates of Glafs did alfo caft the like broad fhadows. And therefore the great breadth of thefe fhadows proceeds from fome other caufe than the refraction of the Air.

Let the Circle X represent the middle of the Hair ; Fig. 1. ADG, BEH, CFI, three rays passing by one fide of the Hair at feveral distances; KNQ, LOR, MPS, three other rays passing by the other fide of the Hair at the like distances; D, E, F and N, O, P, the places where the rays are bent in their paffage by the Hair; G, H, I and Q, R, S, the places where the rays fall on a Paper GQ; IS the breadth of the shadow of the Hair caft on the Paper, and T I, V S, two rays paffing to the points I and S without bending when the Hair is taken away. And it's manifest that all the Light between thefe two rays AI and VS is bent in paffing by the Hair, and turned afide from the fhadow 1S, becaufe if any part of this Light were not bent it would fall on the Paper within the shadow, and there illuminate the Paper contrary to experience. And becaufe when the Paper is at a great distance from the Hair, the shadow is broad, and therefore the rays TI and VS are at a great diftance from one another, it follows that the Hair acts upon the rays of Light at a good diftance in their paffing by it. But the action is ftrongeft on the rays which pass by at least distances, and grows weaker and weaker accordingly as the rays pafs' by at diftances greater and greater, as is represented in the Scheme: For thence it comes to pais, that the fhadow of the Hair is much broader in proportion to the distance of Qq 2 the

the Paper from the Hair, when the Paper is nearer the Hair than when it is at a great diftance from it.

#### OBS. II.

The fhadows of all Bodies (Metals, Stones, Glafs, Wood, Horn, Ice, &c.) in this Light were bordered with three parallel fringes or bands of coloured Light, whereof that which was contiguous to the shadow was broadeft and most luminous, and that which was remotest from it was narrowest, and so faint, as not easily to be visible. It was difficult to diffinguish the Colours unlefs when the Light fell very obliquely upon a fmooth Paper, or fome other finooth vvhite Body, fo as to make them appear much broader than they vvould otherwife do. And then the Colours were plainly visible in this order : The first or innermost fringe was violet and deep blue next the shadovy, and then light blue, green and yellovv in the middle, and red vvithout. The fecond fringe vvas almost contiguous to the first, and the third to the fecond, and both vvere blue vvithin and yellovv and red vvithout, but their Colours vvere very faint efpecially those of the third. The Colours therefore proceeded in this order from the fhadovv, violet, indico, pale blue, green, yellovv, red ; blue, yellovv, red ; pale blue, pale yellovv and red. The shadows made by fcratches and bubbles in polifhed plates of Glafs vvere bordered with the like fringes of coloured Light. And if plates of Looking-glass floop'd off near the edges with a Diamond cut, be held in the fame beam of Light, the Light which paffes through the parallel planes of the Glass will be be bordered with the like fringes of Colours

# [117]

lours where those Planes meet with the Diamond cut, and by this means there will fometimes appear four or five fringes of Colours. Let A B, CD represent the Fig. 2. parallel planes of a Looking-glass, and BD the plane of the Diamond-cut, making at B a very obtuse Angle with the plane A B. And let all the Light between the rays ENI and FBM pass directly through the parallel planes of the Glass, and fall upon the Paper between I and M, and all the Light between the rays GO and HD be refracted by the oblique plane of the Diamond cut B D, and fall upon the Paper between K and L; and the Light which paffes directly through the parallel planes of the Glass, and falls upon the Paper between. I and M, will be bordered with three or more fringes. at M.

#### OBS. III.

When the Hair was twelve Feet diftant from the Hole, and its fhadow fell obliquely upon a flat vvhite fcale of Inches and parts of an Inch placed half a Foot beyond it, and alfo when the fhadow fell perpendicularly upon the fame fcale placed nine Feet beyond it; I meafured the breadth of the fhadow and fringes as accurately as I could, and found them in parts of an Inch as follows.

The

# [118]

At the distance of	Foot.	nine Feet.
The breadth of the Shadow	<u>I</u> 54	19
The breadth between the middles of the brighteft Light of the innermost fringes (me) on either fide the shadow	<sup>1</sup> / <sub>35</sub> Or <sup>1</sup> / <sub>39</sub>	<u>7</u> 50
The breadth between the middles of the brighteft Light of the middlemoft frin- ges of either fide the fhadow	$\frac{1}{23^{\frac{1}{2}}}$	4
The breadth between the middles of the brighteft Light of the outmost fringes place on either fide the shadow	$\frac{1}{18}$ or $\frac{1}{182}$	3 10
The diftance between the middles of the brighteft Light of the firft and fecond fringes	<u>I</u> 120	I . 21
The diftance between the middles of the brighteft Light of the fecond and third fringes	¥ 170	<u>1</u> 31
The breadth of the luminous part (green, white, yellow and red) of the first fringe	<u>r</u> 175	<u>I</u> 32
The breadth of the darker fpace between the first and second fringes.	I 240	1 45
The breadth of the luminous part of the fecond fringe	<u>T</u> 290	1_55
The breadth of the darker fpace between the fecond and third fringes.	1 340	

These measures I took by letting the shadow of the Hair at half a Foot distance fall so obliquely on the scale as to appear twelve times broader than vyhen it fell perpendicularly on it at the same distance, and setting down in this Table the twelfth part of the meafures I then took.

#### OBS. IV.

When the shadovv and fringes vvere cast obliquely upon a smooth vvhite Body, and that Body was removed further and further from the Hair, the first fringe began to appear and look brighter than the reft of the Light at the diftance of lefs than a quarter of an Inchfrom the Hair, and the dark line or fhadovv between that and the fecond fringe began to appear at a lefs distance from the Hair than that of the third part of an Inch. The fecond fringe began to appear at a diftancefrom the Hair of less than half an Inch, and the shadow between that and the third fringe at a diftance lefs than. an Inch, and the third fringe at a diftance lefs than three Inches. At greater diftances they became much more fenfible, but kept very nearly the fame proportion of their breadths and intervals which they had at their first appearing. For the diftance between the middle of the first and middle of the second fringe, was to the distance between the middle of the fecond and middle of the third fringe, as three to two, or ten to feven. And the last of these two distances vvas equal to the breadth. of the bright Light or luminous part of the first fringe. And this breadth vvas to the breadth of the bright Light of the fecond fringe as feven to four, and to the dark interval

interval of the first and second fringe as three to two, and to the like dark interval between the second and third as two to one. For the breadths of the fringes seemed to be in the progression of the numbers 1,  $\nu'_{\frac{1}{3}}$ ,  $\nu'_{\frac{1}{3}}$  and their intervals to be in the same progression vith them; that is, the fringes and their intervals together to be in the continual progression of the numbers  $1, \nu'_{\frac{1}{2}}, \nu'_{\frac{1}{3}}, \nu'_{\frac{1}{4}}, \nu'_{\frac{1}{3}}$ , or thereabouts. And these proportions held the same very nearly at all distances from the Hair; the dark Intervals of the fringes being as broad in proportion to the fringes at their first appearance as afterwards at great distances from the Hair, though not fo dark and distinct.

#### OBS. V.

The Sun shining into my darkened Chamber through a Hole a quarter of an Inch broad; I placed at the diftance of two or three Feet from the Hole a Sheet of Past-board, vvhich vvas black'd all over on both fides, and in the middle of it had a Hole about three quarters of an Inch square for the Light to pass through. And behind the Hole I fastened to the Past-board with Pitch the blade of a sharp Knife, to intercept some part of the Light vyhich paffed through the Hole. The planes of the Past-board and blade of the Knife vvere parallel to one another, and perpendicular to the rays. And vvhen they vvere fo placed that none of the Sun's Light fell on the Past-board, but all of it passed through the Hole to the Knife, and there part of it fell upon the blade of the Knife, and part of it-paffed by its edge: I let this part of the Light vyhich paffed by, fall on a vvhite

white Paper two or three Feet beyond the Knife, and there faw two streams of faint Light shoot out both ways from the beam of Light into the shadow like the tails of Comets. But becaufe the Sun's direct Light by its brightnefs upon the Paper obfcured these faint ftreams, fo that I could fcarce fee them, I made a little Hole in the midft of the Paper for that Light to pass through and fall on a black cloth behind it; and then I faw the two ftreams plainly. They were like one another, and pretty nearly equal in length and breadth, and quantity of Light. Their Light at that end next the Sun's direct Light was pretty ftrong for the space of about a quarter of an Inch, or half an Inch, and in all its progrefs from that direct Light decreased gradually till it became insensible. The whole length of either of these streams measured upon the Paper at the distance of three Feet from the Knife was about fix or eight Inches; fo that it fubtended an Angle at the edge of the Knife of about 10 or 12, or at most 14 degrees. Yet fometimes I thought I faw it fhoot three or four degrees further, but with a Light fo very faint that I could scarce perceive it, and suspected it might ( in fome measure at least) arise from some other cause than the two ftreams did. For placing my Eye in that Light beyond the end of that ftream which was behind the Knife, and looking towards the Knife, I could fee a line of Light upon its edge, and that not only when my Eye was in the line of the streams, but also when it was without that line either towards the point of the Knife, or towards the handle. This line of Light appeared contiguous to the edge of the Knife, and was narrower than the Light of the innermost fringe, and Rr narroweft

narrowest when my Eye was furthest from the direct Light, and therefore seemed to pass between the Light of that fringe and the edge of the Knise, and that which passed nearest the edge to be most bent, though not all of it.

#### OBS. VI.

I placed another Knife by this fo that their edges might be parallel and look towards one another, and that the beam of Light might fall upon both the Knives, and fome part of it pais between their edges. And when the diftance of their edges was about the 400th part of an Inch the ftream parted in the middle, and left a fhadow between the two parts. This fhadow was fo black and dark that all the Light which paffed between the Knives feemed to be bent, and turned afide to the one hand or to the other. And as the Knives ftill approached one another the fhadow grew broader, and the ftreams fhorter at their inward ends which were next the fhadow, until upon the contact of the Knives the whole Light vanifhed leaving its place to the fhadow.

And hence I gather that the Light which is leaft bent, and goes to the inward ends of the ftreams, paffes by the edges of the Knives at the greateft diftance, and this diftance when the fhadow begins to appear between the ftreams is about the eight-hundredth part of an Inch. And the Light which paffes by the edges of the Knives at diftances ftill lefs and lefs is more and more bent, and goes to those parts of the ftreams which are further and further from the direct Light, because when when the Knives approach one another till they touch, those parts of the ftreams vanish last which are furthest from the direct Light.

#### OBS. VII.

In the fifth Observation the fringes did not appear, but by reason of the breadth of the Hole in the Window became fo broad as to run into one another, and by joyning make one continued Light in the beginning of the ftreams. But in the fixth, as the Knives approached one another, a little before the fliadow appeared between the two streams, the fringes began to appear on the inner ends of the streams on either side of the direct Light, three on one fide made by the edge of one Knife, and three on the other fide made by the edge of the other Knife. They were diffincteft when the Knives were placed at the greatest distance from the Hole in the Window, and still became more distinct by making the Hole lefs, infomuch that I could fometimes fee a faint lineament of a fourth fringe beyond the three above-mentioned. And as the Knives continually approached one another, the fringes grew diffincter and larger until they vanished. The outmost fringe vanished first, and the middlemost next, and the innermost last. And after they were all vanished, and the line of Light which was in the middle between them was grown very broad, enlarging it felf on both fides into the streams of Light described in the fifth Observation, the above-mentioned shadow began to appear in the middle of this line, and divide it along the middle. into two lines of Light, and increased until the whole R r 2 Light Light

### [124]

Light vanished. This inlargement of the fringes was fo great that the rays which go to the innermost fringe feemed to be bent above twenty times more when this fringe was ready to vanish, than when one of the Knives was taken away.

And from this and the former Obfervation compared, I gather, that the Light of the first fringe passed by the edge of the Knife at a distance greater than the eighthundredth part of an Inch, and the Light of the fecond fringe passed by the edge of the Knife at a greater diftance than the Light of the first fringe did, and that of the third at a greater distance than that of the fecond, and that of the streams of Light described in the fifth and fixth Observations passed by the edges of the Knives at less distances than that of any of the fringes.

#### OBS. VIII.

I caufed the edges of two Knives to be ground truly ftreight, and pricking their points into a board fo that their edges might look towards one another, and meeting near their points contain a rectilinear Angle, I faftned their handles together with Pitch to make this Angle invariable. The diftance of the edges of the Knives from one another at the diftance of four Inches from the angular point, where the edges of the Knives met, was the eighth part of an Inch, and therefore the Angle contained by the edges was about 1 degr. 54'. The Knives thus fixed together I placed in a beam of the Sun's Light, let into my darkened Chamber through a Hole the 42th part of an Inch wide, at the diftance of of ten or fifteen Feet from the Hole, and let the Light which paffed between their edges fall very obliquely upon a fmooth white Ruler at the diftance of half an Inch, or an Inch from the Knives, and there faw the fringes made by the two edges of the Knives run along the edges of the fhadows of the Knives in lines parallel to those edges without growing fenfibly broader, till they met in Angles equal to the Angle contained by the edges of the Knives, and where they met and joyned they ended without croffing one another. But if the Ruler was held at a much greater diftance from the Paper, the fringes became fomething broader and broader as they approached one another, and after they met they croffed one another, and then became much broader than before.

Whence I gather that the diftances at which the fringes pass by the Knives are not increased nor altered by the approach of the Knives, but the Angles in which the rays are there bent are much increased by that approach; and that the Knise which is nearest any ray determines which way the ray shall be bent, and the other Knise increases the bent.

#### OBS. IX.

When the rays fell very obliquely upon the Ruler at the diftance of the third part of an Inch from the Knives, the dark line between the first and second fringe of the shadow of one Knife, and the dark line between the first and second fringe of the shadow of the other Knifemet with one another, at the distance of the sisted between tween the Knives at the concourse of their edges. And therefore the diftance of the edges of the Knives at the meeting of these dark lines was the 160th part of an Inch. For as four Inches to the eighth part of an Inch, fo is any length of the edges of the Knives measured from the point of their concourse to the distance of the edges of the Knives at the end of that length, and fo is the fifth part of an Inch to the 160th part. So then the dark lines above-mentioned meet in the middle of the Light which paffes between the Knives where they are diftant the 160th part of an Inch, and the one half of that Light paffes by the edge of one Knife at a diftance not greater than the 320th part of an Inch, and falling upon the Paper makes the fringes of the shadow of that Knife, and the other half paffes by the edge of the other Knife, at a distance not greater than the 320th part of an Inch, and falling upon the Paper makes the fringes of the shadow of the other Knife. But if the Paper be held at a diftance from the Knives greater than the third part of an Inch, the dark lines above-mentioned meet at a greater distance than the fifth part of an Inch from the end of the Light which paffed between the Knives at the concourse of their edges; and therefore the Light which falls upon the Paper where those dark lines meet passes between the Knives where their edges are diffant above the 160th part of an Inch.

For at another time when the two Knives were diftant eight Feet and five Inches from the little Hole in the Window, made with a small Pin as above, the Light which fell upon the Paper where the aforefaid dark lines met. paffed between the Knives, where the diftance

[126]

ftance between their edges was as in the following Table, when the diftance of the Paper from the Knives was also as follows.

Diftances of the Paper	Distances between the edges
from the Knives in	of the Knives in mille-
Inches.	fimal parts of an Inch.
$     1\frac{1}{2} \cdot 3\frac{1}{3} \cdot 8\frac{1}{5} \cdot 32 \cdot 32 \cdot 96 \cdot 131 \cdot 15 $	0'012. 0'020. 0'034. 0'057. 0'081. 0'087.

And hence I gather that the Light which makes the fringes upon the Paper is not the fame Light at all diftances of the Paper from the Knives, but when the Paper is held near the Knives, the fringes are made by Light which paffes by the edges of the Knives at a lefs diftance, and is more bent than when the Paper is held at a greater diftance from the Knives.

#### OBS. X.

When the fringes of the fhadows of the Knives fell perpendicularly upon a Paper at a great diffance from the Knives, they were in the form of Hyperbolas, and their dimensions were as follows. Let CA, CB reprefent lines drawn upon the Paper parallel to the edges of the Knives, and between which all the Light would fall, if it passed between the edges of the Knives without inflexion; DE a right line drawn through C making the

the Angles ACD, BCE, equal to one another, and terminating all the Light whith falls upon the Paper from the point where the edges of the Knives meet; eis, fkt, and glv, three hyperbolical lines reprefenting the terminus of the shadow of one of the Knives, the dark line between the first and second fringes of that shadow, and the dark line between the fecond and third fringes of the fame shadow; xip, ykq and zlr, three other Hyperbolical lines reprefenting the terminus of the shadow of the other Knife, the dark line between the first and fecond fringes of that shadow, and the dark line between the fecond and third fringes of the fame shadow. And conceive that these three Hyperbolas are like and equal to the former three, and crofs them in the points i, k and 1, and that the fhadows of the Knives are terminated and diftinguished from the first luminous fringes by the lines e is and x i p, until the meeting and croffing of the fringes, and then those lines cross the fringes in the form of dark lines, terminating the first luminous fringes within fide, and diftinguishing them from another Light which begins to appear at i, and illuminates all the triangular fpace i p D Es comprehended by these dark lines, and the right line DE. Of these Hyperbolas one Afymptote is the line DE, and their other Afymptotes are parallel to the lines CA and CB. Let r v reprefent a line drawn any where upon the Paper parallel to the Afymptote DE, and let this line crofs the right lines AC in m and BC in n, and the fix dark hyperbolical lines in p, q, r; s, t, v; and by meafuring the diftances ps, qt, rv, and thence collecting the the lengths of the ordinates np, nq, nr or ms, mt, mv, and doing this at feveral diftances of the line rv, from

[129]

from the Afymptote DE you may find as many points of thefe Hyperbolas as you pleafe, and thereby know that thefe curve lines are Hyperbolas differing little from the conical Hyperbola. And by meafuring the lines Ci, Ck, Cl, you may find other points of thefe Curves.

For inftance, when the Knives were diftant from the Hole in the Window ten Feet, and the Paper from the Knives 9 Feet, and the Angle contained by the edges of the Knives to which the Angle ACB is equal, was fubtended by a chord which was to the Radius as 1 to 32, and the diftance of the line rv from the Afymptote DE was half an Inch: I meafured the lines ps, qt, rv, and found them 0'35, 0'65, 0'98 Inches refpectively, and by adding to their halfs the line  $\frac{1}{2}$  mn (which here was the 128th part of an Inch, or 0'0078 Inches) the fums np, nq, nr, were 0'1828, 0'3328, 0'4978 Inches. I meafured alfo the diftances of the brighteft parts of the fringes which run between pq and st, qr and tv, and next beyond r and v, and found them 0'5, 0'8, and 1'17 Inches.

#### OBS. XI.

The Sun fhining into my darkened Room through a fmall round Hole made in a plate of Lead with a flender Pin as above; I placed at the Hole a Prifm to refract the Light, and form on the oppofite Wall the Spectrum of Colours, defcribed in the third Experiment of the firft Book. And then I found that the fhadows of all Bodies held in the coloured Light between the Prifm and the Wall, were bordered with fringes of the Colour Ss of that Light in which they were held. In the full red Light they were totally red without any fenfible blue or violet, and in the deep blue Light they were totally blue without any fenfible red or yellow; and fo in the green Light they were totally green, excepting a little yellow and blue, which were mixed in the green Light of the Prism. And comparing the fringes made in the feveral coloured Lights, I found that those made in the red Light were largest, those made in the violet were least, and those made in the green were of a middle bignefs. For the fringes with which the shadow of a Man's Hair were bordered, being measured cross the fhadow at the diftance of fix Inches from the Hair ; the distance between the middle and most luminous part of the first or innermost fringe on one fide of the shadow, and that of the like fringe on the other fide of the shadow, was in the full red Light 1/372 of an Inch, and in the full violet 1/46. And the like diftance between the middle and most luminous parts of the second fringes on either fide the shadow was in the full red Light 12, and in the violet  $\frac{1}{27}$  of an Inch. And these diffances of the fringes held the fame proportion at all diftances from the Hair without any fenfible variation.

So then the rays which made thefe fringes in the red Light paffed by the Hair at a greater diffance than thofe did which made the like fringes in the violet; and therefore the Hair in caufing thefe fringes acted alike upon the red Light or leaft refrangible rays at a greater diftance, and upon the violet or most refrangible rays at a lefs diffance, and by those actions disposed the red Light into larger fringes, and the violet into finaller, and the Lights of intermediate Colours into fringes of inter-

### [131]

intermediate bigneffes without changing the Colour of of any fort of Light.

When therefore the Hair in the first and second of these Observations was held in the white beam of the Sun's Light, and caft a fhadow which was bordered with three fringes of coloured Light, those Colours arose not from any new modifications imprest upon the rays of Light by the Hair, but only from the various inflections whereby the feveral forts of rays were feparated from one another, which before feparation by the mixture of all their Colours, composed the white beam of the Sun's Light, but whenever fepasated compose Lights of the feveral Colours which they are originally difpofed to exhibit. In this 13th Observation, where the Colours are feparated before the Light paffes by the Hair, the least refrangible rays, which when separated from the rest make red, were inflected at a greater distance from the Hair, so as to make three red fringes at a greater diftance from the middle of the fhadow of the Hair; and the most refrangible rays which when feparated make violet, were inflected at a lefs diftance from the Hair, fo as to make three violet fringes at a lefs diftance from the middle of the fhadow of the Hair. And other rays of intermediate degrees of refrangibility were inflected at intermediate diftances from the Hair, fo as to make fringes of intermediate Colours at intermediate diftances from the middle of the shadow of the Hair. And in the second Observation, where all the Colours are mixed in the white Light which passes by the Hair, these Colours are separated by the various inflexions of the rays, and the fringes which they make appear all together, and the innermoft Ss 2 fringes fringes

# [132]

fringes being contiguous make one broad fringe compofed of all the Colours in due order, the violet lying on the infide of the fringe next the fhadow, the red on the outfide furtheft from the fhadow, and the blue, green and yellow, in the middle. And, in like manner, the middlemost fringes of all the Colours lying in order, and being contiguous, make another broad fringe composed of all the Colours; and the outmost fringes of all the Colours lying in order, and being contiguous, make a third broad fringe composed of all the Colours. These are the three fringes of coloured Light with which the fhadows of all Bodies are bordered in the second Observation.

When I made the foregoing Obfervations, I defigned to repeat most of them with more care and exactnels, and to make some new ones for determining the manner how the rays of Light are bent in their passage by Bodies for making the fringes of Colours with the dark lines between them. But I was then interrupted, and cannot now think of taking these things intofurther confideration. And fince I have not finished this part of my Defign, I shall conclude, with propofing only some Queries in order to a further fearch to be made by others.

Query 1. Do not Bodies act upon Light at a diftance, and by their action bend its rays, and is not this action (cateris paribus) strongest at the least distance?

Qu. 2. Do not the rays which differ in refrangibility. differ also in flexibility, and are they not by their different inflexions separated from one another, so as after separation to make the Colours in the three fringes above [133]

above defcribed ? And after what manner are they inflected to make those fringes?

Qu. 3. Are not the rays of Light in paffing by the edges and fides of Bodies, bent feveral times backwards and forwards, with a motion like that of an Eel? And do not the three fringes of coloured Light above-mentioned, arife from three fuch bendings?

Qu. 4. Do not the rays of Light which fall upon Bodies, and are reflected or refracted, begin to bend before they arrive at the Bodies; and are they not reflected, refracted and inflected by one and the fame Principle, acting varioufly in various circumftances?

Qu. 5. Do not Bodies and Light act mutually upon one another, that is to fay, Bodies upon Light in emitting, reflecting, refracting and inflecting it, and Light upon Bodies for heating them, and putting their parts into a vibrating motion wherein heat confifts ?

Qu. 6. Do not black Bodies conceive heat more eafily. from Light than those of other Colours do, by reason that the Light falling on them is not reflected outwards, but enters the Bodies, and is often reflected and refracted within them, until it be stilled and lost?

Qu. 7. Is not the ftrength and vigor of the action between Light and fulphureous Bodies observed above, one reason why fulphureous Bodies take fire more readily, and burn more vehemently, then other Bodies do?

Qu. 8. Do not all fixt Bodies when heated beyond a certain degree, emit Light and shine, and is not this emission performed by the vibrating motions of their parts? wie. allenda.

Qu. 9.

### [134]

Qu. 9. Is not fire a Body heated fo hot as to emit Light copioufly? For what elfe is a red hot Iron than fire? And what elfe is a burning Coal than red hot Wood?

Qu. 10. Is not flame a vapour, fume or exhalation heated red hot, that is, fo hot as to fhine? For Bodies do not flame without emitting a copious fume, and this fume burns in the flame. The Ignis Fatuus is a vapour fhining without heat, and is there not the fame difference between this vapour and flame, as between rotten Wood shining without heat and burning Coals of fire? In diftilling hot Spirits, if the head of the ftill be taken off, the vapour which afcends out of the Still will take fire at the flame of a Candle, and turn into flame, and the flame will run along the vapour from the Candle to the Still. Some Bodies heated by motion or fermentation, if the heat grow intense fume copiously, and if the heat be great enough the fumes will fhine and become flame. Metals in fusion do not flame for want of a copious fume, except Spelter which fumes copioufly, and thereby flames. All flaming Bodies, as Oyl, Tallow, Wax, Wood, foffil Coals, Pitch, Sulphur, by flaming wafte and vanish into burning smoke, which fmoke, if the flame be put out, is very thick and visible, and fometimes finells ftrongly, but in the flame lofes its fmell by burning, and according to the nature of the finoke the flame is of feveral Colours, as that of Sulphur blue, that of Copper opened with Sublimate green, that of Tallow yellow. Smoke paffing through flame cannot but grow red hot, and red hot imoke can have no other appearance than that of flame.

Qu. 11.

Qu. 11. Do not great Bodies conferve their heat the longest, their parts heating one another, and may not great dense and fix'd Bodies, when heated beyond a certain degree, emit Light fo copioufly, as by the emiffion and reaction of its Light, and the reflexions and refractions of its rays within its pores to grow still hotter, till it comes to a certain period of heat, fuch as is that of the Sun? And are not the Sun and fix'd Stars great Earths vehemently hot, whole heat is conferved by the greatness of the Bodies, and the mutual action and reaction between them, and the Light which they emit, and whole parts are kept from fuming away, not only by their fixity, but also by the vast weight and denfity of the Atmospheres incumbent upon them, and very ftrongly compreffing them, and condenfing the vapours and exhalations which arife from them? Wide addenda.

Qu. 12. Do not the rays of Light in falling upon the bottom of the Eye excite vibrations in the Tunica retina? Which vibrations, being propagated along the folid fibres of the optick Nerves into the Brain, caufe the sense of seeing. For because dense Bodies conferve their heat a long time, and the denfest Bodies conferve their heat the longest, the vibrations of their parts are of a lafting nature, and therefore may be propagated along folid fibres of uniform dense matter to a great distance, for conveying into the Brain the impressionsmade upon all the Organs of sense. For that motions which can continue long in one and the fame part of a Body, can be propagated a long way from one part to. another, fuppofing the Body homogeneal, fo that the motion may not be reflected, refracted, interrupted or difordered by any unevennefs of the Body.

Qu. 13 ...

Qu. 13. Do not feveral fort of rays make vibrations of feveral bignefics, which according to their bignefics excite fentations of feveral Colours, much after the manner that the vibrations of the Air, according to their feveral bignefics excite fenfations of feveral founds? And particularly do not the most refrangible rays excite the shortest vibrations for making a fensation of deep violet, the least refrangible the largest for making a fensation of deep red, and the feveral intermediate forts of rays, vibrations of the several intermediate bigneffes to make fensations of the feveral intermediate Colours?

Qu. 14. May not the harmony and difcord of Colours arife from the proportions of the vibrations propagated through the fibres of the optick Nerves into the Brain, as the harmony and difcord of founds arifes from the proportions of the vibrations of the Air? For fome Colours are agreeable, as those of Gold and Indico, and others difagree.

Qu. 15. Are not the Species of Objects feen with both Eyes united where the optick Nerves meet before they come into the Brain, the fibres on the right fide of both Nerves uniting there, and after union going thence into the Brain in the Nerve which is on the right fide of the Head, and the fibres on the left fide of both Nerves uniting in the fame place, and after union going into the Brain in the Nerve which is on the left fide of the Head, and thefe two Nerves meeting in the Brain in fuch a manner that their fibres make but one entire Species or Picture, half of which on the right fide of the Senforium comes from the right fide of both Eyes through the right fide of both



both optick Nerves to the place where the Nerves meet, and from thence on the right fide of the Head into the Brain, and the other half on the left fide of the Senforium comes in like manner from the left fide of both Eyes. For the optick Nerves of fuch Animals as look the fame way with both Eyes (as of Men, Dogs, Sheep, Oxen,  $\Im c$ .) meet before they come into the Brain, but the optick Nerves of fuch Animals as do not look the fame way with both Eyes (as of Fifhes and of the Chameleon) do not meet, if I am rightly informed.

Qu. 16. When a Man in the dark preffes either corner of his Eye with his Finger, and turns his Eye away from his Finger, he will fee a Circle of Colours like those in the Feather of a Peacock's Tail? Do not these Colours arise from such motions excited in the bottom of the Eye by the preffure of the Finger, as at other times are excited there by Light for causing Vision? And when a Man by a stroke upon his Eye sees a Flash of Light, are not the like Motions excited in the *Retina* by the stroke?

Tt



#### BOOK III, Plate I,



# [138]

## ENUMERATIO

# LINEARUM TERTII ORDINIS.

# 

. L BEI ?

ENUMERATIO LINEARUM TERTII ORDINIS.

[139]

Inez Geometricz fecundum numerum dimenfionum æquationis qua relatio inter Ordinatas Line avium Or-& Absciffas definitur, vel (quod perinde est) secun-dum numerum punctorum in quibus a linea recta secari possunt, optime distinguuntur in Ordines. Qua ratione linea primi Ordinis erit Recta fola, eæ fecundi five quadratici ordinis erunt fectiones Conicæ & Circulus, & ex tertii five cubici Ordinis Parabola Cubica, Parabola Neiliana, Ciffois veterum & reliquæ quas hic enumerare susceptimus. Curva autem primi generis, (fiquidem recta inter Curvas non est numeranda) eadem est cum Linea secundi Ordinis, & Curva sécundi generis eadem cum Linea Ordinis tertii. Et Linea Ordinis infinitefimi ea est quam recta in punctis infinitis secare potest, qualis est Spiralis, Cyclois, Quadratrix & linea omnis quæ per radii vel rotæ revolutiones infinitas generatur.

Γ.

Tt 2

Sectionum

11. Proprietates Se-Stionem Conicarum competunt curvis superiorum generum.

III. cundi generis Ordinate, Diametri, Vertices, Centra, Axes.

Sectionum Conicarum proprietates præcipuæ a Geometris passim traduntur. Et confimiles sunt proprietates Curvarum secundi generis & reliquarum, ut ex sequenti proprietatum præcipuarum enumeratione constabit.

Nam fi rectæ plures parallelæ & ad conicam fe-Curvarum se- Aionem utrinq; terminatæ ducantur, recta duas earum bisecans bisecabit alias omnes, ideoq; dicitur Diameter figuræ & rectæ bisectæ dicuntur Ördinatim applicatæ ad Diametrum, & concursus omnium Dia-metrorum est Centrum figuræ, & intersectio Curvæ & diametri Vertex nominatur, & diameter illa Axis est cui ordinatim applicate infistunt ad angulos rectos. Et ad eundem modum in Curvis secundi generis, fi rectæ duæ quævis parallelæ ducantur occurrentes Curvæ in tribus punctis : recta quæ ita secat has parallelas ut fumma duarum partium ex uno fecantis latere ad curvam terminatarum æquetur parti tertiæ ex altero latere ad curvam terminatæ, eodem modo fecabit omnes alias his parallelas curvæq; in tribus punctis occurrentes rectas, hoc est, ita ut summa partium duarum ex uno ipfius latere semper æquetur parti tertiæ ex altero latere. Has itaq; tres partes qua hinc inde æquantur, Ordinatim applicatas & rectam secantem cui ordinatim applicantur. Diametrum & intersectionem diametri & curvæ Verticem & concursum duarum diametrorum Centrum. nominare licet. Diameter autem ad Ordinatas re-Etangula fi modo aliqua fit, etiam Axis dici potest, & ubi omnes diametri in eodem puncto concurrunt. istud erit Centrum generale ...

Hyper-

Hyperbola primi generis duas Asymptotos, ea fe-cundi tres, ea tertii quatuor & non plures habere, po-earum proprieta-test, & fic in reliquis. Et quemadmodum partes tes lineæ cujusvis rectæ inter Hyperbolam Conicam & duas ejus Afymptotos sunt hinc inde æquales : fic in Hyperbolis secundi generis si ducatur recta quævis fecans tam Curvam quàm tres ejus Afymptotos in tribus punctis, summa duarum partium istius rectæ quæ a duobus quibusvis Afymptotis in eandem plagam ad duo puncta Curvæ extenduntur æqualis erit parti tertiæ quæ a tertia Afymptoto in plagam contrariam ad tertium Curvæ punctum extenditur.

Et quemadmodum in Conicis sectionibus non Parabolicis quadratum Ordinatim applicatæ, hoc est, Latera resta cor rectangulum Ordinatarum quæ ad contrarias partes Diametri ducuntur, est ad rectangulum partium Diametri quæ ad Vertices Ellipseos vel Hyperbolæ terminantur, ut data quædam linea quæ dicitur Latus rectum, ad partem diametri quæ inter Vertices jacet & dicitur Latus transversum : fic in Curvis non Parabolicis fecundi generis Parallelepipedum sub tribus. Ordinatim applicatis est ad Parallelepipedum sub partibus Diametriad Ordinatas & tres Vertices figuræabfciffis, in ratione quadam data : in qua ratione fi fumantur tres rectæ ad tres partes diametri inter vertices figuræ fitas fingulæad fingulas, tunc illæ tresrecte dici poffunt Latera recla figure, & ille partes. Diametri inter Vertices Latera transversa. Et ficut: in Parabola Conica quæ ad unam & eandem diametrum unicum tantum habet Verticem, rectangulum: sub Ordinatis æquatur rectangulo sub parte Diametri quæ ad Ordinatas & Verticem abscinditur & recta: quadam

quadam data que Latus rectum dicitur, fic in Curvis secundi generis quæ non nifi duos habent Vertices ad eandem Diametrum, Parallelepipedum fubOrdinatis tribus æquatur Parallelepipedo iub duabus partibus Diametri ad Ordinatas & Vertices illos duos abfciffis, & recta quadam data quæ proinde *Latus rectum* dici potest.

VI. Ratio contentorum sub Paralle-

larum segmentis.

J.a.

Deniq; ficut in Conicis sectionibus ubi duæ parallelæ ad Čurvam utrinq; terminatæ fecantur a duabus parallelis ad Curvam utrinq; terminatis, prima a tertia & fecunda a quarta, rectangulum partium primæ est ad rectangulum partium tertiæ ut rectangulum partium fecundæ ad rectangulum partium quartæ: fic ubi quatuor tales rectæ occurrunt Curvæ fecundi generis fingulæ in tribus punctis, parallele-pipedum partium primæ rectæ erit ad parallelepidedum partium tertiæ, ut parallelepipedum partium fecundæ ad parallelepipedum partium quartæ.

Curvarum secundi & superiorum generum æque VII. bolica & Parabo- atq; primi crura omnia in infinitum progredientia Crura Hyperlica & eorum pla- vel Hyperbolici funt generis vel Parabolici. Crus Hyperbolicum voco quod ad Afymptoton aliquam in infinitum appropinquat, Parabolicum quod Afymptoto destituitur. Hæc crura ex tangentibus optime dignoscuntur. Nam si punctum contactus in infinitum abeat tangens cruris Hyperbolici cum Afymptoto coincidet & tangens cruris Parabolici in infinitum recedet, evanescet & nullibi reperietur. Invenitur igitur Afymptotos cruris cujuívis quærendo tangen-tem cruris illius ad punctum infinite diftans. Plaga autem cruris infiniti invenitur quærendo pofitionem rectæ cujuívis quæ tangenti parallela est ubi punctum
[143]

Etum contactus in infinitum abit. Nam hæc recta in eandem plagam cum crure infinito dirigitur.

Linex omnes Ordinis primi, tertii, quinti, sep-Linex omnes Ordinis primi, tertii, quinti, fep- VIII. timi & imparis cujufq; duo habent ad minimum varum omnium ctura in infinitum versus plagas oppositas progre-generis secundi ad dientia. Et lineæ omnes tertii Ordinis duo habent aquationum casus quatuor. Casus ejusmodi crura in plagas oppositas progredientia in primus. quas nulla alia earum crura infinita (præterquam in Parabola Cartefiana) tendunt. Si crura illa fint Hyperbolici generis, fit GAS eorum Afymptotos & huic parallela agatur recta quævis CBc ad Curvam utrinque (fi fieri potest ) terminata eademq; bisectur in puncto X, & locus puncti il- Fig. 1. lius X erit Hyperbola Conica (puta X •) cujus una Afymptotos eft AS. Sit ejus altera Áfymptotos AB, & aquatio qua relatio inter Ordinatam BC & Absciffam AB definitur, fi AB dicatur x & BC y, femper induct hanc formam  $x y y + e y = a x^3$ -+bxx+cx+d. Ubi termini e, a, b, c, d, defignant quantitates datas cum fignis suis + & - affe-Aas, quarum quælibet deeffe poffunt modo ex earum defectu figura in sectionem conicam non vertatur. Poteft autem Hyberbola illa Conica cum afymptotis suis coincidere, id est punctum X in recta AB locari : & tunc terminus + e y deeft.

At fi recta illa CBc non poteft utrinq; ad Curvam terminari fed Curvæ in unico tantum puncto occurrit: age quamvis pofitione datam rectam A B afymptoto AS occurrentem in A, ut & aliam quamvis BC afymptoto illi parallelam Curvæque occurrentem in puncto C, & æquatio qua relatio inter Ordinatam BC

IX. Čafus fecundus,

### [144]

BC & Absciffam AB definitur, semper induct hanc formam  $xy = ax^3 + bxx + cx + d$ .

X. Casus tertius. Quod fi crura illa oppofita Parabolici fint generis, recta CB c ad Curvam utrinque, fi fieri poteft, terminata in plagam crurum ducatur & bifecetur in B, & locus puncti B erit linea recta. Sit ifta AB, terminata ad datum quodvis punctum A, & æquatio qua relatio inter Ordinatam BC & Abfciffam AB definitur, femper induet hanc formam,  $yy=ax^3$ +bxx+cx+d.

At vero fi recta illa CB c in unico tantum puncto occurrat Curvæ, ideoq; ad Curvam utrinq; terminari non poffit : fit punctum illud C, & incidat recta illa ad punctum B in rectam quamvis aliam pofitione datam & ad datum quodvis punctum A terminatam AB: & æquatio qua relatio inter Ordinatam BC & Abfciffam AC definitur femper induet hanc formam,  $y=ax^3+bxx+cx+d$ .

Enumerando curvas horum cafuum, Hyperbolam vocabimus *infcriptam* quæ tota jacet in Afymptoton angulo ad inftar Hyperbolæ conicæ, *circumfcriptam* quæ Afymptotos fecat & partes abfciffas in finu fuo amplectitur, *ambigenam* quæ uno crure infinito infcribitur & altero circumfcribitur, *convergentem* cujus crura concavitate fua feinvicem refpiciunt & in plagam eandem diriguntur, *divergentem* cujus crura convexitate fua feinvicem recipiunt & in plagas contrarias diriguntur, *cruribus contrariis præditam* cujus crura in partes contrarias convexa funt & in plagas contrarias infinita, *Conchoidalem* quæ vertice concavo & cruribus divergentibus ad afymptoton applicatur, *anguineam* quæ flexibus contrariis afymptoton fecat

XI. Caſus quartus.

XII. Nomina forma-

rum.

& utrinq; in crura contraria producitur, cruciformem quæ conjugatam decuffat, nodatam quæ feipfam decuffat in orbem redeundo, cuspidatam cujus partes duæ in angulo contactus concurrunt & ibi terminantur, punctatam quæ conjugatam habet Ovalem infinite parvam id eft punctum, & puram quæ per impoffibilitatem duarum radicum Ovali, Nodo, Cufpide & Puncto conjugato privatur. Eodem fenfu Parabolam quoq; convergentem, divergentem, cruribus contrariis præditam, cruciformem, nodatam, cufpidatam, punctatam & puram nominabimus.

In cafu primo fi terminus a x<sup>3</sup> affirmativus eft Figura erit Hyperbola triplex cum fex cruribus Hy-redundante & perbolicis quæ juxta tres Afymptotos quarum nullæ eius tribus Afunt parallelæ in infinitum progrediuntur, binæ juxta unamquamq; in plagas contrarias. Et hæ Afymptoti fi terminus bxx non deeft fe mutuo fecabunt in tribus punctis triangulum (Dd $^{\circ}$ ) inter fe continentes, fin terminus bxx deeft convergent omnes ad idem punctum. In priori cafu cape AD =  $\frac{1}{24}$ , & Ad = A $^{\circ} = \frac{b}{27d}$ , ac junge Dd, D $^{\circ}$ , & erunt AD, Dd, D $_{\circ}$  tres Afymptoti. In pofteriori duc ordinatam quamvis BC, & in ea utrinq; producta cape hine inde BF & Bf fibi mutuo æquales & in ea ratione ad A B quam habet  $\sqrt{d}$  ad a, jungeq; AF, Af, & erunt AB, AF, Af tres Afympoti. Hanc autem Hyperbolam vocamus redundantem quia numero crurum Hyperbolicorum Sectiones Conicas fuperat.

In Hyperbola omni redundante fi neq; terminus XIV. De hujus Hye y defit neq; fit bb-4ac æquale ± a e v a curva nul-perbola diametris lam habebit diametrum, fin eorum alterutrum ac-U u cidat

cidat curva habebit unicam diametrum, & tres fi utrumque. Diameter autem femper transit per intersectionem duarum Asymptoton & bisecat rectas omnes quæ ad Afymptotos illas utrinq; terminantur & parallelæ funt & Afymptoto tertiæ. Eftq; absciffa AB diameter Figuræ quoties terminus ey deeft. Diametrum vero absolute dictam hic & in sequentibus in vulgari fignificatu ufurpo, nempe pro absciffa que passim habet ordinatas binas equales ad idem punctum hinc inde infiftentes.

XV. habent Asymptotos triangulum capientes.

Fig. 1,2.

Si Hyperbola redundans nullam habet diametrum vem redundantes quærantur Æquationis hujus ax4+bx3+cxx+dx qua diametro de-  $+\frac{1}{4}=0$  radices quatuor seu valores ipsius x. Ex stituuntur & tres sunto AP, Aw, Aw, Ap. Erigantur ordinate PT,  $\pi\tau$ ,  $\pi$ , pt, & hæ tangent Curvam in punctis totidem T,  $\tau$ ,  $\eta$ , t, & tangendo dabunt limites Curvæ per quos species ejus innotescet.

Nam fi radices omnes AP, A =, A =, A p funt reales, ejusdem figni & inæquales, Curva constat ex tribus Hyperbolis, (infcripta circumfcripta & ambigena ) cum Ovali. Hyperbolarum una jacet verfus D, altera versus d, tertia versus o, & Ovalis femper jacet intra triangulam Dd , atq; etiam inter medios limites  $7 \& \tau$ , in quibus utiq; tangitur ab ordinatis #7 & wr. Et hæc eft species prima.

Fig. 3.4.

Si e radicibus duæ maximæ A $\pi$ , A p, vel duæ minimæ AP, A = æquantur inter se, & ejusdem sunt figni cum alteris duobus, Ovalis & Hyperbola circumscripta fibi inxicem junguntur coeuntibus earum punctis contactus 7 & t vel T & 7, & crura Hyperbolæ sefe decussando in Ovalem continuantur, figuram nodatam efficientia. Quæ species est secunda.

Si

Si e radicibus tres maximæ Ap,  $A\pi$ ,  $A\pi$ ,  $A\pi$ , vel tres Fig. 5, 6. minimæ  $A\pi$ ,  $A\pi$ ,

Si e radicibus dux medix  $A^{\varpi} & A^{\pi}$  xquentur in-Fig. 7. ter fe, puncta contactus  $\tau & \gamma$  coincidunt, & propterea Ovalis interjecta in punctum evanuit, & conftat figura ex tribus Hyperbolis, inferipta, circumferipta & ambigena cum *puncto* conjugato. Qux eft fpecies quarta.

Si duæ ex radicibus funt impoffibiles & reliquæ Fig. 7,8,13,14. duæ inæquales & ejufdem figni (nam figna contraria habere nequeunt,) puræ habebuntur Hyperbolæ tres fine Ovali vel Nodo vel cufpide vel puncto conjugato, & hæ Hyperbolæ vel ad latera trianguli ab Afymptotis comprehenfi vel ad angulos ejus jacebunt & perinde fpeciem vel quintam vel fextam conftituent.

Si e radicibus duæ funt æquales & alteræ duæ Fig. 9,10,15,16 vel impoffibiles funt vel reales cum fignis quæ a fignis æqualium radicum diversa funt, figura cruciformis habebitur, nempe duæ ex Hyperbolis seinvicem decussabunt idq; vel ad verticem trianguli ab Asymptotis comprehensi, vel ad ejus basem. Quæ duæ species sunt septima & octava.

Si deniq; radices omnes funt impoffibiles vel fi Fig. 11, 12 omnes funt reales & inæquales & earum duæ funt affirmativæ & alteræ duæ negativæ, tunc duæ habebuntur Hyperbolæ ad angulos oppofitos duarum U u 2 Afymp-

### [148]

Afymptoton cum Hyperbola anguinea circa Afymptoton tertiam. Quæ species est nona.

Et hi funt omnes radicum cafus poffibiles. Nam fi duæ radices funt æquales inter fe, & aliæ duæ funt etiam inter fe æquales, Figura evadet Sectio Conica cum linea recta.

XVI. Si Hyperbola redundans habet unicam tantum Hyperbola duodecim redundantes cum unicatan- tionis hujus  $ax^3 + bxx + cx + d = 0$  quære tres ratum Diametro. dices feu valores x.

Fig. 17.

Si radices illæ funt omnes reales & ejufdem figni, Figura conftabit ex *Ovali* intra triangulum  $Dd \circ ja-$ cente & tribus Hyperbolis ad angulos ejus, nempe circumfcripta ad angulum D & infcriptis duabus ad angulos d &  $\circ$ . Et hæc eft fpecies decima.

Si radices duæ majores funt æquales & tertia ejufdem figni, crura Hyperbolæ jacentis verfus D fefe decuffabunt in forma *Nodi* propter contactum Ovalis. Quæ fpecies eft undecima.

Si tres radices funt æquales, Hyperbola ista fit cuspidata fine Ovali. Quæ species est duodecima.

Si radices duæ minores funt æquales & tertia ejufdem figni, Ovalis in *punctum* evanuit. Quæ fpecies eft decima tertia. In fpeciebus quatuor noviffimis Hyperbola quæ jacet verfus D Afymptotos in finu fuo amplectitur, reliquæ duæ in finu Afymptotôn jacent.

Si duæ ex radicibus funt impoffibiles habebuntur tres Hyperbolæ *puræ* fine Ovali decuffatione vel cufpide. Et hujus cafus fpecies funt quatuor, nempe decima quarta fi Hyperbola circumfcripta jacet verfus D & decima

Fig. 18.

Fig. 19.

Fig. 20.

Fig. 20.

Fig. 21. Fig. 22. Fig. 23. decima quinta fi Hyperbola infcripta jacet verfus D, decima fexta fi Hyperbola circumfcripta jacet fub bafi d<sup>o</sup> trianguli Dd<sup>o</sup>, & decima feptima fi Hyperbola infcripta jacet fub eadem bafi.

Si dux radices funt xquales & tertia figni diverfi Fig. 24. figura erit cruciformis. Nempe dux ex tribus Hyperbolis feinvicem decuffabunt idq; vel ad verticem trianguli ab Afymptotis comprehenfi vel ad ejus bafem. Qux dux fpecies funt decima octava & decima nona.

Si duæ radices funt inæquales & ejufdem figni & tertia eft figni diverfi, duæ habebuntur Hyperbolæ in oppofitis angulis duarum afymptotôn cum Conchoidali intermedia. Conchoidalis autem vel jace-Fig. 27. bit ad eafdem partes afymptoti fuæ cum triangulo ab afymptotis conftituto, vel ad partes contrarias; & hi duo cafus conftituunt fpeciem vigefimam & vigefimam primam.

Hyperbola redundans quæ habet tres diametros XVII. Hyperbola duæ conftat ex tribus Hyperbolis in finubus afymptoôn redundantes cum jacentibus, idq; vel ad angulos trianguli ab afymptotribus Diametris. tis comprehenfi vel ad ejus latera. Cafus prior dat Fig. 29. fpeciem vigefimam fecundam,& posterior speciem vigefimam tertiam.

Si tres afymptoti in puncto communi fe mutuo XVIII. decuffant, vertuntur species quinta & fexta in vigefimam quartam, feptima & octava in vigefimam cum Asymptotis quintam, & nona in vigefimam fextam ubi Anguinea non transit per concursum afymptoton, & in vigefimam septimam ubi transit per concursum illum, quo casu termini b ac d defunt, & concursus afymptotôn est centrum figuræ ab omnibus ejus partibus Fig. 33oppositis

XVIII. Kyperbole novem redundantes um Afymptotis

#### [150]

oppositis æqualiter distans. Et hæ quatuor species Diametrum non habent.

Vertuntur etiam species decima quarta ac decima fexta in vigefimam octavam, decima quinta ac decima feptima in vigefimam nonam, decima octava & decima nona in tricefimam, & vigefima cum vige-fima prima in tricefimam primam. Et hæ fpecies unicam habent diametrum.

Ac deniq; species vigesima secunda & vigesima tertia vertuntur in speciem tricesimam secundam cujus tres funt Diametri per concursum asymptotôn transeuntes. Quæ omnes conversiones facillime intelliguntur faciendo ut triangulum ab asymptotis comprehensum diminuatur donec in punctum evanescat.

Si in primo æquationum casu terminus ax<sup>3</sup> negativus est, Figura erit Hyberbola defectiva unicam etrum non haben- habens afymptoton & duo tantum crura Hyperbolica juxta afymptoton illam in plagas contrarias in-finite progredientia. Et afymptotos illa est Ordinata prima & principalis AG. Si terminus e y non deest figura nullam habebit Diametrum, si deest habebit unicam. In priori cafu fpecies fic enumerantur.

> Si æquationis hujus  $ax^4 = bx^3 + cxx + dx + \frac{1}{4}ee$ , radices omnes A m, A P, A p, A m, funt reales & inæquales, Figura erit Hyperbola anguinea afymptoton flexu contrario amplexa, cum Ovali conjugata. Quæ species est tricesima tertia.

> Si radices dux medix AP & Ap xquentur inter se, Ovalis & Anguinea junguntur sese decussantes in forma Nodi. Quæ est species tricesima quarta. Si

Fig. 34. Fig. 35. Fig. 36. Fig. 37.

Fig. 38.

XIX. Hyperbola Sex defective diamer .t.eso

F10.39.

F19.40.

Si tres radices funt æquales, Nodus vertetur in Fig: 41.  $cu \int pidem$  acutiffimum in vertice anguineæ. Et hæc eft species tricesima quinta.

[ IJI ]

Si e tribus radicibus ejusdem figni dux maxim $x^{Fig. 43}$ . A p & A = fibi mutuo æquantur, Ovalis in *punclum* evanuit. Quæ species est tricesima sexta.

Si radices dux quxvis imaginarix funt, fola manebit Anguinea *pura* fine Ovali, decuffatione, cufpide vel puncto conjugato. Si Anguinea illa non Fig. 42. transit per punctum A species est tricesima septima, fin transit per punctum illud A (id quod contingit Fig. 43. ubi termini b ac d defunt,) punctum illud A erit centrum sigurx rectas omnes per ipsum ductas & ad Curvam utrinq; terminatas bisecans. Et hxc est species trices trices actions.

In altero cafu ubi terminus e y deeft & propterea XX. figura Diametrum habet, fi æquationis hujus a x<sup>3</sup> Hyperbola feptem defettiva, ditem 
Si duæ radices funt inæquales & ejufdem figni & Fig. 44. tertia eft figni contrarii, Ovalis jacebit ad concavitatem Conchoidalis. Eftq; fpecies quadragefima.

Si radices dux minores AT, At, funt xquales Fig. 46. & tertia A7 est ejusdem signi, Ovalis & Conchoidalis jungentur seste decussando in modum Nodi. Qux species est quadragesima prima.

Si tres radices funt æquales, Nodus mutabitur in Fig. 47? Cuspidem & figura erit Cissi Veterum. Et hæc eft fpecies quadragefima fecunda.

Si

Fig. 49.

Fig. 49.

Fig. 48,49.

Si radices duæ majores funt æquales, & tertia eft ejufdem figni,Conchoidalis habebit *punctum* conjugatum ad convexitatem fuam, eftq; species quadragefima tertia.

Si radices dux funt xquales & tertia est figni contrarii Conchoidalis habebit *punctum* conjugatum ad concavitatem fuam, estq; species quadragesima quarta.

Si radices duæ funt impossibiles habebitur Conchoidalis *pura* fine Ovali, Nodo, Cuspide vel puncto conjugato. Quæ species est quadragesima quinta.

XXI. Hyperbola feptem Parabolica Diametrum non habentes.

Siquando in primo æquationum cafu terminus a x<sup>3</sup> deeft & terminus b x x non deeft, Figura erit Hyperbola Parabolica duo habens crura Hyperbolica ad unam Afymptoton SAG & duo Parabolica in plagam unam & eandem convergentia. Si terminus e y non deeft figura nullam habebit diametrum, fin deeft habebit unicam. In priori cafu fpecies funt hæ.

Fig. 50.

Fig. 51.

Si tres radices AP, A=, A=, A= æquationis hujus bx<sup>3</sup>+cx +dx+ $\frac{1}{4}$  ee=o funt inæquales & ejuídem figni, figura conftabit ex *Ovali* & aliis duabus Curvis quæ partim Hyperbolicæ funt & partim Parabolicæ. Nempe crura Parabolica continuo ductu junguntur cruribus Hyperbolicis fibi proximis. Et hæc eft fpecies quadregefima fexta.

Si radices duæ minores funt æquales & tertia eft ejufdem figni, Ovalis & una Curvarum illarum Hyperbolo-Parabolicarum junguntur & fe decuffant in formam *Nodi*. Quæ fpecies eft quadragefima feptima.

Si

[153]

Si tres radices funt æquales, Nodus ille in Cuf-Fig. 52. pidem vertitur. Estq; species quadragesima octava.

Si radices dux majores funt æquales & tertia eft Fig. 53. ejufdem figni, Ovalis in *punctum* conjugatum evanuit. Qux fpecies eft quadragefima nona.

nuit. Que fpecies est quadragesima nona. Si due radices sunt impossibiles, manebunt pur e Fig. 53,54ille due curve Hyperbolo-parabolice fine Ovali, decussation e, cuspide vel puncto conjugato, & speciem quinquagesimam constituent.

Si radices dux funt xquales & tertia est figni con-<sup>Fig. 55</sup>. trarii, Curvx illx hyperbolo-parabolicx junguntur fese decussion in morem crucis. Estq; species quinquagesima prima.

Si radices duæ funt inæquales & ejufdem figni & <sup>Fig. 56.</sup> tertia eft figni contrarii, figura evadet Hyperbola anguinea circa Afymptoton AG, cum Parabola conjugata. Et hæc eft fpecies quinquagefima fecunda.

In altero cafu ubi terminus ey deeft & figura XXII. Diametrum habet, fi duæ radices æquationis hujus tuor Parabolica bxx-|-cx-|-d=0 funt impoffibiles, duæ habentur Diametrum hafiguræ hyperbolo-parabolicæ a Diametro AB hinc Fig. 57. inde æqualiter diftantes. Quæ fpecies eft quinquagefima tertia.

Si æquationis illius radices duæ funt impoffibiles, <sup>Fig. 58.</sup> Figuræ hyperbolo-parabolicæ junguntur fefe decuffantes in morem crucis, & fpeciem quinquagefimam quartam constituunt.

Si radices illæ funt inæquales & ejufdem figni, ha- Fig. 59 betur Hyperbola Conchoidalis cum Parabola ex eodem latere Afymptoti. Estq; species quinquagesima quinta.

Хх

Fig. 60.

XXIII. Quatuor Hyperbolifmi Hyperbola.

Si radices illæ funt figni contrarii, habetur Conchoidalis cum Parabola ad alteras partes Afymptoti. Quæ fpecies eft quinquagefima fexta.

[154]

Siquando in primo æquationum cafu terminus uterq; a x<sup>3</sup> & b x x deeft, figura erit Hyperbolifmus fectionis alicujus Conicæ. Hyperbolifmum figuræ voco cujus Ordinata prodit applicando contentum fub Ordinata figuræ illius & recta data ad Abfciffam communem. Hac ratione linea recta vertitur in hyperbolam Conicam, & fectio omnis Conica vertitur in aliquam figurarum quas hic Hyperbolifmos fectionum Conicarum voco. Nam æquatio ad figuras de quibus agimus, nempe x y y + e y = cx + d, feu  $y = \frac{e \pm \sqrt{ee + 4} dx + 4 cxx}{2x}$  generatur applicando contentum fub Ordinata fectionis Conicæ  $e \pm \sqrt{ee + 4} dx + 4 cxx}$  secta data m ad curvarum

Absciffam communem x. Unde liquet quod figura genita Hyperbolismus erit Hyperbolæ, Ellipseos vel Parabolæ perinde ut terminus cx affirmativus est vel negativus vel nullus.

Hyperbolifmus Hyperbolæ tres habet afymptotos quarum una eft Ordinata prima & principalis A d, alteræ duæ funt parallelæ Abfciffæ A B & ab eadem hinc inde æqualiter diftant. In Ordinata principali A d cape A d, A & hinc inde æquales quantitati V c & per puncta d ac & age d g,  $\delta \gamma$  Afymptotos Abfciffæ A B parallelas.

Ubi terminus e y non deest figura nullam habet diametrum. In hoc casu fi æquationis hujus  $cxx+dx+\frac{1}{4}ee=0$  radices duæ AP, Ap sunt reales & inæquales (nam æquales effe nequeunt nifi figura Fig. 61) fit Conica fectio) figura conftabit ex tribus Hyperbolis fibi oppofitis quarum una jacet inter afymptotos parallelas & alteræ duæ jacent extra. Et hæc eft fpecies quinquagefima feptima.

Si radices illæ duæ funt impoffibiles, habentur Hyperbolæ duæ oppofitæ extra afymptotos parallelas & Anguinea hyperbolica intra eafdem. Hæc figura duarum eft fpecierum. Nam centrum non habet Fig. 621 ubi terminus d non deeft ; fed fi terminus ille deeft <sup>Fig. 634</sup> punctum A eft ejus centrum. Prior fpecies eft quinquagefima octava, pofterior quinquagefima nona.

Quod fi terminus e y deeft, figura conftabit ex Fig. 643tribus hyperbolis oppofitis quarum una jacet inter afymptotos parallelas & alteræ duæ jacent extra ut in fpecie quinquagefima quarta, & præterea diametrum habet quæ eft abfciffa AB. Et hæc eft fpecies fexagefima.

Hyperbolifmus Ellipfeos per hanc æquationem definitur x y y + e y = cx + d, & unicam habet afympmi Ellipfeos. toton quæ eft Ordinata principalis A d. Si terminus Fig. 65. e y non deeft, figura eft Hyperbola anguinea fine diametro atq; etiam fine centro fi terminus d non deeft. Quæ fpeeies eft fexagefima prima.

At si terminus d'deest, figura habet centrum sine Fig. 66. diametro & centrum ejus est punctum A. Species vero est sexagesima secunda.

Et si terminus e y deest & terminus d non deest, Fig. 67. figura est Conchoidalis ad afymptoton AG, habetq; diametrum sine centro, & diameter ejus est Abscissa AB. Quæ species est sexagesima tertia.

X x 2

Hyper-

XXV. Duo Hyperbolismi Parabola.

Fig. 68.

Fig. 69.

XXVI. Tridens.

Fig. 76.

XXVII. que divergentes.

Fig. 70, 71.

Fig. 72. Fig. 73.

Hyperbolifmus Parabolæ per hanc æquationem definitur x y y + e y = d; & duas habet alymptotos, Absciffam AB & Ordinatam primam & principalem AG. Hyperbolæ vero in hac figura funt duæ, nonin afymptoton angulis oppofitis sed in angulis qui funt deinceps jacentes, idq; ad utrumq; latus absciffa AB, & vel fine diametro si terminus ey habetur, vel cum diametro si terminus ille deest. Quæ dux species sunt sexagesima quarta & sexagesima. quinta:

In fecundo æquationum cafu habebatur æquatio  $xy=ax^{3}+bxx+cx+d$ . Et figura in hoc cafu habet quatuor crura infinita quorum duo funt hyperbolica circa afymptoton AG in contrarias partes tendentia & duo Parabolica convergentia & cum prioribus speciem Tridentis fere efformantia. Estq; hæc Figura Parabola illa per quam Cartefius æquationes sex dimensionum construxit. Hæc est igitur species sexagesima sexta.

In tertio cafu æquatio erat  $yy = ax^3 + bxx + cx$ Parabola quin- + d, & Parabolam defignat cujus crura divergunt ab invicem & in contrarias partes infinite progrediuntur. Absciffa AB est ejus diameter & species ejus funt quinq; sequentes.

> Si æquationis a  $x^3 + bx^2 + cx + d = 0$  radices omnes A7, AT, At funt reales & inxquales, figura est Parabola divergens campaniformis cum Ovali ad verticem. Et species est sexagesima septima.

> Si radices duæ funt æquales, Parabola prodit vel nodata contingendo Ovalem, vel punctata ob Ovalem infinite parvam. Quæ duæ species sunt sexagesima octava & fexagefima nona.

Si tres radices sunt æquales Parabola erit cuspi- Fig. 73. data in vertice. Et hæc eft Parabola Neiliana quæ vulgo femicubica dicitur.

Si radices duz funt impoffibiles, habetur Parabola Fig. 73, 74. pura campaniformis speciem septuagesimam primam constituens.

In quarto cafu æquato erat y = ax + bxx + cxXXVIII. +d, & hæc æquatio Parabolam illam Wallifianam Parabola cubica. designat que crura habet contraria & cubica dici solet. Et sic species omnino sunt septuaginta dux.

Si in planum infinitum a puncto lucido illuminatum umbræ figurarum projiciantur, umbræ sectio- rum per Umbras. num Conicarum femper erunt fectiones Conica, eæ Curvarum secundi generis semper erunt Curvæ secundi generis, eæ curvarum tertii generis femper erunt Curvæ tertii generis, & fic deinceps in infinitum. Et quemadmodum Circulus umbram projiciendo generat sectiones omnes conicas, fic Parabolæ quinq; divergentes umbris suis generant & exhibent alias omnes secundi generis curvas, & sic Curvæ quædam fimpliciores aliorum generum inveniri possunt quæ alias omnes eorundem generum curvas umbris fuis a puncto lucido in planum projectis formabunt.

Diximus Curvas fecundi generis a linea recta in XXX. punctis tribus fecari posse. Horum duo nonnun- ta duplicia, quam coincidunt. Ut cum recta per Ovalem infi-Curvarum puna= nite parvam transit vel per concursum duarum partium Curvæ se mutuo secantium vel in cuspidem coeuntium ducitur. Et siquando rectæ omnes in plagam

XXIX. Genesis Curvaplagam cruris alicujus infiniti tendentes Curvam in unico tantum puncto fecant (ut fit in ordinatis Parabolæ Cartefianæ & Parabolæ cubicæ, nec non in rectis Abfciffæ Hyperbolifmorum Hyberbolæ & Parabolæ parallelis ) concipiendum eft quod rectæ illæ per alia duo Curvæ puncta ad infinitam diftantiam fita (ut ita dicam) tranfeunt. Hujufmodi interfectiones duas coincidentes five ad finitam fint diftantiam five ad infinitam, vocabimus punctum duplex. Curvæ autem quæ habent punctum duplex defcribi poffunt per fequentia Theoremata.

XXXI.

Theoremata de Curvarum defcriptione organica. Fig. 78.

1. Si anguli duo magnitudine dati PAD, PBD circa polos pofitione datos A, B rotentur, & eorum crura AP, BP concurfu fuo P percurrant lineam rectam; crura duo reliqua AD, BD concurfu fuo D defcribent fectionem Conicam per polos A, B tranfeuntem: præterquam ubi linea illa recta tranfit per polorum alterutrum A vel B, vel anguli BAD, ABD fimul evanefcunt, quibus in cafibus punctum D defcribet lineam rectam.

2. Si crura prima A P, B P concurfu fuo P percurrant fectionem Conicam per polum alterutrum A tranfeuntem, crura duo reliqua A D, BD concurfu fuo D defcribent Curvam fecundi generis per polum alterum B tranfeuntem & punctum duplex habentem in polo primo A per quem fectio Conica tranfit : præterquam ubi anguli BAD, A B D fimul evanefcunt, quo cafu punctum etum D describet aliam sectionem Conicam per polum A transeuntem.

[159]

2. At fi fectio Conica quam punctum P percurrit transeat per neutrum polorum A, B, punctum D describet curvam secundi vel tertii generis pun-Aum duplex habentem. Et punAum illud duplex in concursu crurum describentium, AD, BD invenietur ubi anguli BAP, ABP fimul evanefcunt. Curva autem descripta secundi erit generis fi anguli BAD, ABD fimul evanescunt, alias erit tertii generis & alia duo habebit puncta duplicia in polis A & B.

Jam sectio Conica determinatur ex datis ejus XXXII. punctis quinq; & per cadem fic describi potest. nicarum descrip-Sectionum Co-Dentur ejus puncta quinq; A, B, C, D, E. Jun- tio per data quin-gantur eorum tria quævis A, B, C & trianguli ABC que paneta. rotentur anguli duo quivis CAB, CBA circa vertices suos A & B, & ubi crurum AC, BC interfectio C fucceffive applicatur ad puncta duo reliqua D, E, incidat intersectio crurum reliquorum AB & BA in puncta P & Q. Agatur & infinite producatur recta PQ, & anguli mobiles ita rotentur ut interfectio crurum AB, BA percurrat rectam PQ, & crurum reliquorum intersectio C describet propositam fectionem Conicam per Theorema primum.

XXXIII.

Curvæ omnes secundi generis punctum duplex cundigeneris punhabentes determinantur ex datis earum punctis ctum duplex hafeptem, quorum unum est punctum illud duplex, tio per data sep-

Curvarum sebentium descrip-& tem punsta.

#### [160]

& per eadem puncta sic describi possunt. Dentur Curvæ describendæ puncta quælibet septem A, B, C, D, E, F, G quorum A est punctum duplex. Jungantur punctum A & alia duo quævis e punctis puta B&C; & trianguli ABC rotetur tum angulus CAB circa verticem suum A, tum angulorum reliquorum alteruter ABC circa verticem suum B. Et ubi crurum AC, BC concurfus C fucceffive applicatur ad puncta quatuor reliqua D, E, F, G incidat concursus crurum reliquorum A B & B A in puncta quatuor P, Q, R, S. Per puncta illa quatuor & quintum A describatur sectio Conica, & anguli præfati CAB, CBA ita rotentur ut crurum AB, BA concursus percurrat sectionem illam Conicam, & concursus reliquorum crurum AC, BC describet Curvam propofitam per Theorema fecundum.

Si vice puncti C datur pofitione recta BC quæ Curvam defcribendam tangit in B, lineæ AD, AP coincident, & vice anguli DAP habebitur linea recta circa polum A rotanda.

Si punctum duplex A infinite diftat debebit Recta ad plagam puncti illius perpetuo dirigi & motu parallelo ferri interea dum angulus ABC circa polum B rotatur.

Defcribi etiam poffunt hæ curvæ paulo aliter per Theorema tertium, fed defcriptionem fimpliciorem pofuiffe fufficit.

Eadem methodo Curvas tertii, quarti & fuperiorum generum defcribere licet, non omnes quidem fed quotquot ratione aliqua commoda per motum localem defcribi poffunt. Nam curvam aliquam fecundi

#### [ 161 ]

fecundi vel fuperioris generis punctum duplex non habentem commode describere Problema est inter difficiliora numerandum.

Curvarum usus in Geometria est ut per earum Proponatur quationum per deintersectiones Problemata folvantur. æquatio construenda dimensionum novem  $x^{9*} + bx^{7}$  scriptionem Cur- $-cx^{6}-dx^{5}-ex^{4}+fx^{3}-gxx+hx-k=0$ . Ubi varum. --- m

b, c, d,  $\Im c$ . fignificant quantitates quasivis datas fignis suis + & - affectas. Assume a quatio ad Parabolam cubicam  $x^3 = y$ , & æquatio prior, fcribendo y pro x<sup>3</sup>, evadet  $y^3$  + bx yy + cyy + dx xy  $+exy+my+fx^{3}-gxx+hx+k=0$ , æquatio ad Curvam aliam fecundi generis. Ubi m vel f deeffe potest vel pro lubitu assumi. Et per harum Curvarum descriptiones & intersectiones dabuntur radices æquationis construendæ. Parabolam cubicam semel describere sufficit.

Si æquatio conftruenda per defectum duorum terminorum ultimorum hx & k reducatur ad septem dimenfiones, Curva altera delendo m, habebit punctum duplex in principio absciffa, & inde facile defcribi potest ut supra.

Si æquatio construenda per defectum terminorum trium ultimorum gxx+hx+k reducatur ad fex dimensiones, Curva altera delendo f evadet fectio Conica.

Et si per defectum sex ultimorum terminorum æquatio construenda reducatur ad tres dimensiones, incidetur in constructionem Wallisianam per Parabolam cubicam & lineam rectam.

Yy

Con-

XXXIV. Constructio &-

#### [162]

Conftrui etiam poffunt æquationes per Hyperbolifinum Parabolæ cum diametro. Ut fi conftruenda fit hæc æquatio dimenfionum novem termino penultimo carens,  $a+cxx+dx^3+ex^4+fx+gx^6+hx^7$ -+m

 $+kx^{8}+1x^{9}=0$ ; affumatur æquatio ad Hyperbolifmum illum xxy=1, & fcribendo y pro  $\frac{1}{xx}$ , æquatio conftruenda vertetur in hanc  $ay^{3}+cyy+dxyy+ey$  $+fxy+mxxy+g+hx+kxx+1x^{3}=0$ , quæ curvam fecundi generis defignat cujus defcriptione Problema folvetur. Et quantitatum m ac g alterutra hic deeffe poteft, vel pro lubitu affumi.

Per Parabolam cubicam & Curvas tertii generis conftruuntur etiam æquationes omnes dimenfionum non plufquam duodecim, & per eandem Parabolam & curvas quarti generis conftruuntur omnes dimenfionum non plufquam quindecim, Et fic deinceps in infinitum. Et curvæ illæ tertii quarti & fuperiorum generum defcribi femper poffunt inveniendo eorum puncta per Geometriam planam. Ut fi conftruenda fit æquatio x<sup>11</sup>\* +a x<sup>10</sup>+bx<sup>9</sup>+cx<sup>8</sup>+dx<sup>7</sup>+ex<sup>6</sup>+fx<sup>5</sup> +gx<sup>4</sup>+hx<sup>3</sup>+ixx+kx+l=0, & defcripta habeatur Parabola Cubica; fit æquatio ad Parabolam illam cubicam x<sup>3</sup> = y, & fcribendo y pro x<sup>3</sup> æquatio conftruenda vertetur in hanc y<sub>4</sub> +axy<sup>3</sup>+cxxyy+fxxy+ixx=0, quæ eft +b +dx +gx +kx +e +h +l

æquatio ad Curvam tertii generis cujus descriptione Problema solvetur. Describi autem potest hæc Curva inveniendo ejus puncta per Geometriam planam, propterea quod indeterminata quantitas x non nisi ad duas dimensiones ascendit.













#### Curvarum Tab. IV.









and the second 
.





# TRACTATUS D E Quadratura Curvarum.



## INTRODUCTIO:

[ 165 ]

Quantitates Mathematicas non ut ex partibus quam minimis conftantes, fed ut motu continuo defcriptas hic confidero. Lineæ defcribuntur ac defcribendo generantur non per appofitionem partium fed per motum continuum punctorum, fuperficies per motum linearum, folida per motum fuperficierum, anguli per rotationem laterum, tempora per fluxum continuum, & fic in cæteris. Hæ Genefes in rerum natura locum vere habent & in motu corporum quotidie cernuntur. Et ad hunc modum Veteres ducendo rectas mobiles in longitudinem rectarum immobilium genefin docuerunt rectangulorum.

Confiderando igitur quod quantitates æqualibus temporibus crefcentes & crefcendo genitæ, pro velocitate majori vel minori qua crefcunt ac generantur, evadunt majores vel minores; methodum quærebam deterdeterminandi quantitates ex velocitalibus motuum vel incrementorum quibus generantur; & has motuum vel incrementorum velocitates nominando Fluxiones & quantitates genitas nominando Fluentes, incidi paulatim Annis 1665 & 1666 in Methodum Fluxionum qua hic ufus fum in Quadratura Curvarum.

Fluxiones sunt quam proxime ut Fluentium augmenta æqualibus temporis particulis quam minimis genita, & ut accurate loquar, funt in prima ratione augmentorum nascentium; exponi autem possunt per lineas quafcunq; quæ funt ipfis proportionales. Ut fi areæ ABC, ABDG Ordinatis BC, BD super basi A B uniformi cum motu progredientibus describantur, harum arearum fluxiones erunt inter se ut Ordinatæ defcribentes BC & BD, & per Ordinatas illas exponi poffunt, propterea quod Ordinatæ illæ funt ut arearum augmenta nascentia. Progrediatur Ordinata BC de loco fuo BC in locum quemvis novum b c. Compleatur parallelogrammum BCEb, ac ducatur recta VTH que Curvam tangat in C ipfisq; bc & BA productis occur-rat in T & V : & Absciffæ AB, Ordinatæ BC, & Lineæ Curvæ ACc augmenta modo genita erunt Bb, Ec & Cc; & in horum augmentorum nascentium ratione prima funt latera trianguli CET, ideoq; fluxiones ipsarum AB, BC & AC sunt ut trianguli illius CET latera CÉ, ET & CT & per eadem latera exponi poffunt, vel quod perinde est per latera trianguli confimilis VBC.

Eodem recidit fi fumantur fluxiones in ultima ratione partium evanefcentium. Agatur recta Cc & producatur eadem ad K. Redeat Ordinata bc in

Fig. L.
in locum fuum priorem BC, & coeuntibus punctis C & c, recta CK coincidet cum tangente CH, & triangulum evanescens CEc in ultima sua forma evadet fimile triangulo CET, & ejus latera evanescentia CE, Ec & Cc erunt ultimo inter se ut sunt trianguli alterius CET latera CE, ET & CT, & propterea in hac ratione sunt fluxiones linearum AB, BC & AC. Si puncta C & c parvo quovis intervallo ab invicem distant recta CK parvo intervallo a tangente CH distabit. Ut recta CK cum tangente CH coincidat & rationes ultimæ linearum CE, Ec & Cc inveniantur, debent puncta C & c coire & omnino coincidere. Errores quam minimi in rebus mathematicis non funt contemnendi.

Simili argumento fi circulus centro B radio BC defcriptus in longitudinem Abfciffæ AB ad angulos rectos uniformi cum motu ducatur, fluxio folidi geniti ABC erit ut circulus ille generans, & fluxio fuperficiei ejus erit ut perimeter Circuli illius & fluxio lineæ curvæ AC conjunctim. Nam quo tempore folidum ABC generatur ducendo circulum illum in longitudinem Abfciffæ AB, eodem fuperficies ejus generatur ducendo perimetrum circuli illius in longitudinem Curvæ AC.

Recta PB circa polum datum P revolvens secet aliam Fig. 2... positione datam rectam AB: quæritur proportio fluxionum rectarum illarum AB & PB. Progrediatur recta PB de loco suo PB in locum novum Pb. In Pb capiatur PC ipsi PB æqualis, & ad AB ducatur PD sic, ut angulus bPD æqualis sit angulo bBC; & ob similitudinem triangulorum bBC, bPD erit augmentum Bb ad augmentum Cb ut Pb ad Db. Redeat Redeat jam P b in locum fuum priorem P B ut augmenta illa evanefcant, & evanefcentium ratio ultima, id eft ratio ultima P b ad D b, ea erit quæ eft P B ad D B, exiftente angulo P D B recto, & propterea in hac ratione eft fluxio ipfius A B ad fluxionem ipfius P B.

Recta P B circa datum Polum P revolvens fecet alias duas positione datas rectas  $AB \oslash AE$  in  $B \oslash$ E: quæritur proportio fluxionum rectarum illarum  $AB \oslash AE$ . Progrediatur recta revolvens PB de loco suo PB in locum novum P b rectas AB, AE in punctis b & e secantem, & recta AE parallela BC ducatur ipsi P b occurrens in C, & erit Bb ad BC ut Ab ad Ae, & BC ad Ee ut PB ad PE, & conjunctis rationibus Bb ad Ee ut AbxPB ad AexPE. Redeat jam linea P b in locum sum priorem PB, & augmentum evanescens Bb erit ad augmentum evanescens Ee ut ABxPB ad AExPE, ideoq; in hac ratione est fluxio recta AB ad fluxionem recta AE.

Hinc fi recta revolvens PB lineas quafvis Curvas pofitione datas fecet in punctis B & E, & rectæ jam mobiles AB, AE Curvas illas tangant in Sectionum punctis B & E: erit fluxio Curvæ quam recta, AB tangit ad fluxionem Curvæ quam recta AE tangit ut ABxPB ad AExPE. Id quod etiam eveniet fi recta PB Curvam aliquam pofitione datam perpetuo tangat in puncto mobili P.

Fluat quantitas x uniformiter & invenienda sit fluxio quantitatis  $x^n$ . Quo tempore quantitas x fluendo evadit x+0, quantitas  $x^n$  evadet  $x+0|^n$ , id est per methodum serierum infinitarum,  $x^n-n o x^{n-1}$ 

Fig. 3.

# [ 169 ]

 $-\frac{1}{2}\frac{nn-n}{2}\cos^{n-2}+\mathcal{G}c$ . Et augmenta o &  $nox^{n-1}+\frac{nn-n}{2}\cos^{n-2}$ - $-\mathcal{G}c$ . funt ad invicem ut I &  $nx^{n-1}-\frac{nn-n}{2}\cos^{n-2}+\mathcal{G}c$ . Evanefcant jam augmenta illa, & eorum ratio ultima erit I ad  $nx^{n-1}$ : ideoq; fluxio quantitatis x eft ad fluxionem quantitatis  $x^n$  ut I ad  $nx^{n-1}$ .

Similibus argumentis per methodum rationum primarum & ultimarum colligi poffunt fluxiones linearum feu rectarum feu curvarum in cafibus quibufcunque, ut & fluxiones fuperficierum, angulorum & aliarum quantitatum. In finitis autem quantitatibus Analyfin fic inftituere, & finitarum naicentium vel evanefcentium rationes primas vel ultimas inveftigare, confonum eft Geometriæ Veterum : & volui oftendere quod in Methodo Fluxionum non opus fit figuras infinite parvas in Geometriam introducere. Peragi tamen poteft Analyfis in figuris quibufcunq; feu finitis feu infinite parvis quæ figuris evanefcentibus finguntur fimiles, ut & in figuris quæ pro infinite parvis haberi folent, modo caute procedas.

Ex Fluxionibus invenire Fluentes Problema difficilius est, & folutionis primus gradus æquipollet Quadraturæ Curvarum; de qua sequentia olim scripsi.

Żz

DE

[170]

# TRACTATUS

# Quadratura Curvarum.

Uantitates indeterminatas ut motu perpetuo crefcentes vel decrefcentes, id eft ut fluentes vel defluentes in fequentibus confidero, defignoq; literis z, y, x, v, & earum fluxiones feu celeritates crefcendi noto iifdem literis punctatis z, y, x, v. Sunt & harum fluxionum fluxiones feu mutationes magis aut minus celeres quas ipfarum z, y, x, v fluxiones fecundas nominare licet & fic dignare z, y, x, v, & harum fluxiones primas feu ipfarum z, y, x, v, & harum fluxiones primas feu ipfarum z, y, x, v, Et quemadmodum z, y, x, v funt fluxiones quantitatum z, y, x, v, & hæ funt fluxiones quantitatum z, y, x, v : fic hæ quantitates confiderari poffunt ut fluxiones aliarum quas fic defignabo,

 $Z_{2}$ 

# 

z, y, x, v, & hæ ut fluxiones aliarum z, y, x, v, & hæ ut fluxiones aliarum z, y, x, v. Defignant igitur z, z, z, z, z, z, z, z &c. feriem quantitatum quarum quælibet posterior est fluxio præcedentis & quælibet prior est fluens quantitas fluxionem habens subsequentem. Similis est series Vaz-zz, Vaz-zz, Vaz-zz, Vaz-zz, Vaz-zz, Vaz-zz, ut & feries  $\frac{az+z^2}{z^2} = \frac{az+z^2}{az+z^2} =$ 2----Z 3----7 a-z a----Z 9 az--z2 . Et notandum est quod quantitas quælibet a---7 prior in his seriebus est ut area figuræ curviliniæ cujus ordinatim applicata rectangula est quantitas posterior & abscissa est z : uti Vaz-zz area curvæ cujus ordinata est vaz-zz & abscissa z. Quo autem spectant hæc omnia patebit in Propositionibus quæ sequuntur.

Zz 2

PROP.

# [172]

#### PROP. I. PROB. I.

Data æquatione quotcunq; fluentes quantitates involvente, invenire fluxiones.

#### Solutio.

Multiplicetur omnis æquationis terminus per indicem dignitatis quantitatis cujusq; fluentis quam involvit, & in fingulis multiplicationibus mutetur dignitatis latus in fluxionem suam, & aggregatum factorum omnium sub propriis signis erit æquatio nova.

## Explicatio.

Sunto a, b, c, d  $\Im c$ . quantitates determinatæ & immutabiles, & proponatur æquatio quævis quantitates fluentes z, y, x  $\Im c$ . involvens, uti x<sup>3</sup> — x y y  $-+aaz-b^3=0$ . Multiplicentur termini primo per indices dignitatum x, & in fingulis multiplicationibus pro dignitatis laterc, feu x unius dimenfionis, fcribatur x,& fumma factorum erit  $3 \times x^2 - x y y$ . Idem fiat in y & prodibit—<sup>2</sup>x y y. Idem fiat in z & prodibit a a z. Ponatur fumma factorum æqualis nihilo, & habebitur æquatio  $3 \times x^2 - x y y - x y y$ -|-aaz=0. Dico quod hac æquatione definitur relatio fluxionum.

De-

# [173]

# Demonstratio.

Nam fit o quantitas admodum parva & funto oz, oy, ox, quantitatum z, y, x momenta id eft incrementa momentanea synchrona. Et si quantitates fluentes jam funt z, y & x, hæ post momentum temporis incrementis suis oz, oy, ox aucta, evadent z-l-oz, y-l-oy, x-l-ox, quæ in æquatione prima pro z, y & x scriptæ dant æquationem x<sup>3</sup> -- 3xxox  $-3x00xx - 0^{3}x^{3} - xyy - 0xyy - 2x0yy - 2x00yy$  $-xooyy-xo^3yy-aaz-aaoz-b_3=0$ . Subducatur æquatio prior, & refiduum divifum per o erit 2xx2  $-3xxox - x^{3}oo - xyy - 2xyy - 2xoyy - xoyy - xooyy$ lectis terminis evanescentibus restabit 3xx2-xyy -xyy+aaz=0. Q. E. D.

## Explicatio plenior.

# [174]

& (per hanc Propositionem)  $ax^{-2}yy = 2z$  feu  $ax^{-2}yy = z$ , hoc eft  $ax^{-2}yy = \sqrt{ax} - yy$ . Et  $2z^{-2}z$ 

inde 
$$3x^2x - xyy - 2xyy + \frac{a^3x - 2aayy}{\sqrt{ax - yy}} = 0$$

Et per operationem repetitam pergitur ad fluxiones fecundas, tertias & fequentes. Sit æquatio  $zy^3-z_4-a^4=0$ , & fiet per operationem primam  $zy^3+3zyy^2-4zz^3=0$ , per fecundam  $zy^3+6zyy^2$  $+3zyy^2+6zy^2y-4zz^3-12z^2z^2=0$ , per tertiam  $zy^3+9zyy^2-9zyy_2+18zy_2y+3zyy^2+18zyyy$  $-6zy^3-4zz^3-36zzz^2-24z^3z=0$ .

Ubi vero fic pergitur ad fluxiones fecundas, tertias & fequentes, convenit quantitatem aliquam ut uniformiter fluentem confiderare, & pro ejus fluxione prima unitatem fcribere, pro fecunda vero & fequentibus nihil. Sit æquatio  $zy^3 - z^4 + a^4 = 0$ , ut fupra; & fluat z uniformiter, fitq; ejus fluxio unitas, & fiet per operationem primam  $y^3 + 3zyy^2 - 4z_3 = 0$ , per fecundam  $6yy^2 + 3zyy^2 + 6zy^2y - 12z^2 = 0$ , per tertiam  $9yy^2 + 18y^2y + 3zyy^2 + 18zyyy + 6zy_3 - 24z = 0$ .

-In

# [175]

In hujus autem generis æquationibus concipiendum eft quod fluxiones in fingulis terminis fint ejufdem ordinis, id eft vel omnes primi ordinis y, z, vel omnes fecundi y, y<sup>2</sup>, yz, z<sup>2</sup>, vel omnes tertii y, yy, yz, y<sup>3</sup>, y<sup>2</sup>z, yz<sup>2</sup> z<sup>3</sup> &c. Et ubi res aliter fe habet complendus eft ordo per fubintellectas fluxiones quantitatis uniformiter fluentis. Sic æquatio noviffima complendo ordinem tertium fit  $9zyy^{2}$  $+18zy^{2}y--3zyy^{2}--18zyyy--6zy^{3}--24zz^{3}=0.$ 

#### PROP. II. PROB. II.

#### Invenire Curvas quæ quadrari possunt.

Sit ABC figura invenienda, BC Ordinatim ap-Fig. 4. plicata rectangula, & AB abfciffa. Producatur CB ad E ut fit BE=1, & compleatur parallelogrammum ABED: & arearum ABC, ABED fluxiones erunt ut BC & BE. Affumatur igitur æquatio quævis qua relatio arearum definiatur, & inde dabitur relatio ordinatarum BC & BE per Prop. I. Q. E. I.

Hujus rei exempla habentur in Propositionibus duabus sequentibus.

PROP.

# [176]

#### PROP. III. THEOR. I.

Si pro abfciffa AB & area AE feu AB×1 promifcue fcribatur z, & fi pro e  $-|-fz^n + gz^{2n} + hz^{3n} - kc$ . fcribatur R: fit autem area Curvæ  $z_{\theta}R^{\Lambda}$  erit. ordinatim applicata BC=

 ${}^{\theta}e + {}^{\theta}_{+\lambda n} fz^{n} + {}^{\theta}_{+2\lambda n} gz^{2n} + {}^{\theta}_{+3\lambda n} hz^{3n} + \&c. in z^{\theta-1} R^{\lambda-1}.$ 

## Demonstratio.

Nam fi fit  $z^{\theta}R^{\lambda} = v$ , erit per Prop. 1,  $\theta z z^{\theta-1}R_{\lambda}$   $-1 - \lambda z^{\theta}RR^{\lambda-1} = v$ . Pro  $R^{\lambda}$  in primo æquationis termino &  $z^{\theta}$  in fecundo foribe  $RR^{\lambda-1}$  &  $zz^{\theta-1}$ , & fiet  $\theta z R - \lambda z R$  in  $z^{\theta-1}R^{\lambda}_{-1} = v$ . Erat autem  $R = e - fz^{\mu}$   $(-gz^{2n} + hz^{3n})$  & c. & inde per Prop. 1. fit  $R = i^{\mu}$   $r fzz^{\mu-1} + 2^{\mu}gzz^{2\mu-1} + 3^{\mu}hzz^{3\mu-1} + \infty$ . quibus fubfitutis & foripta BE feu I pro z, fiet  $\theta e + \theta + fz^{\mu} + \theta = z^{2\mu} + \theta + 3^{2\mu} + z^{3\mu} + \infty$ . in  $z^{\theta-1}R^{\lambda-1} = v = BC$ . Q. E. D.

PROP.

# [177]

#### PROP. IV. THEOR. II.

 $\begin{array}{c} \theta e k_{-1-\lambda n}^{-1-\theta} f k \ z^{n} \stackrel{+\theta}{\xrightarrow{}} g k z^{2n} \cdot \ast \cdots \ast \\ \stackrel{-1-\theta}{\xrightarrow{}} e l \ z^{n} \stackrel{-\theta}{\xrightarrow{}} f l \ z^{2n} \stackrel{+\theta}{\xrightarrow{}} g l z^{3n} \ast \cdots \ast \\ \stackrel{-1-\lambda n}{\xrightarrow{}} e^{-1-\lambda n} \stackrel{-1-2\lambda n}{\xrightarrow{}} g l z^{3n} \ast \cdots \ast \\ \stackrel{-1-\lambda n}{\xrightarrow{}} \stackrel{-1-\lambda n}{\xrightarrow{}} \frac{e l z^{2n} -e \theta}{\xrightarrow{}} f l \ z^{2n} \stackrel{-\theta}{\xrightarrow{}} g l z^{3n} \ast \cdots \ast \\ \stackrel{-1-\lambda n}{\xrightarrow{}} \frac{e l z^{2n} -e \theta}{\xrightarrow{}} f l \ z^{2n} \stackrel{-\theta}{\xrightarrow{}} g l z^{3n} \ast \cdots \ast \\ \stackrel{-1-\lambda n}{\xrightarrow{}} \frac{e l z^{2n} -e \theta}{\xrightarrow{}} f l \ z^{2n} \stackrel{-\theta}{\xrightarrow{}} g l z^{3n} \stackrel{-\theta}{\xrightarrow{}} g l z^{2n} \xrightarrow{} \theta g l z$ 

Demonstratur ad modum Propositionis superioris.

PROP. V. THEOR. III.

Si Curvæ abfeiffa AB fit z, & pro e+fz<sup>n</sup>+gz<sup>2n</sup> +hz<sup>3n</sup> + &c. feribatur R : fit autem ordinatim applicata  $z^{\theta-1}R^{\lambda-1}$  in a +  $bz^n$  +  $cz^{2n}$  +  $dz^{3n}$  + &c. & ponatur  $\int_{n}^{\theta} = r. r + \lambda = s. s + \lambda = t. t + \lambda = v.$  &c. erit area  $z^{\theta}R^{\lambda}$  in  $\int_{re}^{\frac{1}{n}a} + \frac{\int_{r}^{\frac{1}{n}b} - sfA}{r + 1, e} + \frac{\int_{r}^{\frac{1}{n}c} - \int_{r}^{s} fB - tgA}{r + 2, e} + \frac{\int_{n}^{\frac{1}{n}d} - \int_{r}^{s} fC - \int_{r}^{t} gB - vhA}{r + 3, e}$  $+ \frac{\int_{r}^{-s} fD - \int_{r}^{t} gC - \int_{r}^{v} hB}{r + 4, e} z^{4n} +$ &c. Ubi A, B, C, D, &c. A = a denotant

# [178]

denotant totas coefficientes datas terminorum fingulorum in ferie cum fignis fuis-|-&-, nempe A primi termini coefficientem  $\frac{\frac{1}{n}a}{\frac{1}{r}e}$ , B fecundi coefficientem  $\frac{\frac{1}{n}c}{\frac{1}{r}+1,e}$ , C tertii coefficientem  $\frac{\frac{1}{n}c}{\frac{1}{r}+2,e}$ , & fic deinceps.

# Demonstratio.

# Sunto juxta Propositionem tertiam,

Curvarum Ordinatæ	& earundem areæ.
1. $\theta e A \xrightarrow{-1-\theta}_{-1-\lambda_n} f A z^n \xrightarrow{-1-\theta}_{-1-\lambda_n} g A z^{2n} \xrightarrow{-1-\theta}_{-1-\lambda_n} h A z^{3n} \& c.$	$Az^{\theta} R^{\lambda}$ .
$2 \cdots \theta \overline{f_{n}}, eBz^{n} \overline{f_{-n}} fBz^{2n} \overline{f_{-n}} gBz^{3n} \&c.$	Bzo+" R <sup>1</sup> .
$3 \qquad \qquad +\theta+2n, eCZ^{2n-\int_{-\lambda_n}} \theta CZ^{3n} \& c.$	$Cz^{\theta+2n} \mathbb{R}^{\lambda}$ .
4	$Dz^{g+3u} \mathbb{R}^{\lambda}$ .

Et fi fumma ordinatarum ponatur æqualis ordinatæ a- $|-bz^n-|-cz^{2n}-|-dz^{3n}-|-\&c.$  in  $z^{\theta-1}R^{\lambda-1}$ , fumma arearum  $z^{\theta}R^{\lambda}$  in  $A-j-Bz^n-j-Cz^{2n}-j-Dz^{3n}-j-\&c.$  æqualis erit areæ Curvæ cujus ifta eft ordinata. Æquentur igitur Ordinatarum termini correipondentes, & fiet  $a=\theta eA$ ,  $b=\frac{\theta}{-f\lambda n}fA \xrightarrow{+\theta}{-f\theta}eB$ ,  $c=\frac{\theta}{-f-2\lambda n}gA \xrightarrow{-f\theta+n}{-f\lambda n}fB$  $-f-\overline{\theta+2n}, eC$  &c. & inde  $\frac{a}{\theta e} = A$ .  $\frac{b--(\theta-f-\lambda n)fA}{\theta-f-n} = B$ .  $c-(\theta-f-2\lambda n), gA \xrightarrow{-\overline{\theta+n}+\lambda n}fB} = C$ . Et fic deinceps in infinitum

# [179]

nitum. Pone jam  $\frac{\theta}{n} = r$ .  $r + \lambda = s$ .  $s + \lambda = t$  &c. & in area  $z^{\theta}R^{\lambda} \times A + Bz^{n} + Cz^{2n} + Dz^{3n}$  &c. fcribe ipforum A, B, C, &c. valores inventos & prodibit feries propofita. Q. E. D.

Et notandum est quod Ordinata omnis duobus modis iu feriem resolvitur. Nam index " vel affirmativus est potest vel negativus. Proponatur Ordi-nata  $\frac{3 k-12 z}{zz \sqrt{kz-1z} 3-1-mz_4}$ . Hæc vel sic scribi potest Hæc vel fic fcribi poteft  $z = \frac{1}{2} \times \frac{1}{2} \times \frac{1}{2} \times \frac{1}{2} - \frac{1}{2} = \frac{1}{2}$ , vel fic  $z \times -\frac{1}{2} - \frac{3}{2} \times \frac{1}{2}$  $\times m - lz^{-1} + kz^{-3} - \frac{1}{2}$ . In cafu priore eft a = 3k.b = o.  $c=-1. e=k. f=0. g=-1. h=m. \lambda = -\frac{1}{2}. n=1.$  $\theta - I = -\frac{5}{2}, \quad \theta = -\frac{3}{2} = I, \quad S = -I, \quad t = -\frac{1}{2}, \quad V = 0.$  In posteriore est a = -1. b = o. c = 3k. e = m. f = -1.  $g=0, h=1, \lambda=-\frac{1}{2}, \mu=-1, \theta=1, \theta=2, r=-2.$  $s = -I_{\frac{1}{2}}$ . t = -I.  $v = -I_{\frac{1}{2}}$ . Tentandus eft cafus uterque. Et si serierum alterutra ob terminos tandem deficientes abrumpitur ac terminatur, habebitur area Curvæ in terminis finitis. Sic in exempli hujus priore cafu scribendo in ferie valores ipforum a, b, c, e, f, g, h, x, 0, r, s, t, v, termini omnes post primum evanescunt in infinitum & area Curvæ prodit  $-2\sqrt{\frac{k-lzz-l-mz_3}{z_3}}$ . Et hæc area ob fignum negativum adjacet absciffæ ultra ordinatam productæ. Nam area omnis affirmativa adjacet tam absciffæ quam ordinatæ, negativa vero cadit ad contrarias partes ordinatæ & adjacet abscisse productæ, manente scilicet signo Ordinate. Hoc modo series alterutra & nonnunquam utraque semper terminatur & finita evadit fi Curva geometrice quadrari poteft. At fi Curva talem quadraturam non admittit, feries utraq; continuabitur in infinitum, & ea-Aaa 2 rum

# [ 180 ]

rum altera converget & aream dabit approximando, præterquam ubi r (propter aream infinitam) vel nihil eft vel numerus integer & negativus, vel ubi  $\frac{2}{e}$ æqualis eft unitati. Si  $\frac{2}{e}$  minor eft unitate, converget feries in qua index " affirmativus eft : fin  $\frac{2}{e}$  unita te major eft, converget feries altera. In uno cafu area adjacet abfciffæ ad ufq; ordinatam ductæ, in altero adjacet abfciffæ ultra ordinatam productæ.

Nota infuper quod fi Ordinata contentum eft fub factore rationali Q & factore furdo irreducibili R<sup> $\pi$ </sup>, & factoris furdi latus R non dividit factorem rationalem Q; erit  $\lambda - I = \pi \& R^{\lambda-I} = R^{\pi}$ . Sin factoris furdi latus R dividit factorem rationalem femel, erit  $\lambda - I = \pi - I \& R^{\lambda-I} = R^{\pi+I}$ : fi dividit bis, erit  $\lambda - I = \pi - I \& R^{\lambda-I} = R^{\pi+I}$ : fi dividit bis, erit  $\lambda - I = \pi - I \& R^{\lambda-I} = R^{\pi+2}$ : fi ter, erit  $\lambda - I = \pi - I - 3$ , &  $R^{\lambda-I} = R^{\pi+3}$ : & fic deinceps.

Si Ordinata eft fractio rationalis irreducibilis cum Denominatore ex duobus vel pluribus terminis compofito : refolvendus eft denominator in divifores fuos omnes primos. Et fi divifor fit aliquis cui nullus alius eft æqualis, Curva quadrari nequit : Sin duo vel plures fint divifores æquales, rejiciendus eft eorum unus, & fi adhuc alii duo vel plures fint fibi mutuo æquales & prioribus inæquales, rejiciendus eft etiam eorum unus, & fic in aliis omnibus æqualibus fi adhuc plures fint : deinde divifor qui relinquitur vel contentum fub diviforibus omnibus qui relinquuntur, fi plures funt, ponendum eft pro R, & ejus quadrati reciprocum R<sup>-2</sup> pro R<sup>\*-1</sup>, præterquam ubi contentum illud eft quadratum vel cubus vel quadrato quadratum,&c. quo cafu ejus latus ponen-

# [ 181 ]

ponendum est pro R & potestatis index 2 vel 3 vel 4 negative sumptus pro x. & Ordinata ad denominatorem R<sup>2</sup> vel R<sup>3</sup> vel R<sup>4</sup> vel R<sup>5</sup> &c. reducenda.

Ut fi ordinata fit  $\frac{25+24-8z_3}{25+24-5z_3-22+8z-4}$ ; quoniam hac fractio irreducibilis eft & denominatoris divifores funt pares, nempe z-1, z-1, z-1 & z+2, z-1-2, rejicio magnitudinis utriufque diviforem unum & reliquorum z-1, z-1, z-2 contentum z<sup>3</sup>-3z-2 pono pro R & ejus quadrati reciprocum  $\overline{R^2}$  feu R-2 pro R<sup>A-1</sup>. Dein Ordinatam ad denominatorem R<sup>2</sup> feu R<sup>1-A</sup> reduco, & fit  $z^6-9z^4+8z^3$  $\overline{z^3-3z+2}$  quad., id eft  $z_3 \times 8-9z+z_3 \times 2-3z+z^3/2^2$ Et inde eft a=8. b=-9. c=0. d=-1, &c. e=2. f=-3. g=0. h=1. A-1=-2. A=-1.  $n=1. \theta-1=3. \theta=4=r. s=3. t=2. v=1.$  Et his in ferie fcriptis prodit area  $\frac{z^4}{z^3-3z+2}$ , terminis omnibus in tota ferie poft primum evanefcentibus.

Si deniq; Ordinata est fractio irreducibilis & ejus denominator contentum est sub factore rationali Q-& factore surdo irreducibili R<sup> $\pi$ </sup>, inveniendi sunt lateris R divisores omnes primi, & rejiciendus est divisor unus magnitudinis cujusq; & per divisores qui restant, siqui sint, multiplicandus est factor rationalis Q: & fi factum æquale est lateri R vel lateris illius potestati alicui cujus index est numerus integer, esto index ille m, & erit  $\lambda - I = -\pi - m$ , &  $R^{\lambda-1} = R^{-\pi-m}$ . Ut fi Ordinata sit  $\frac{3q^3 - q + x - pq^3 xx - qqx^3 - 6qx^4}{qq - xx \sqrt{cub} \cdot q^3 - pqx - qxx - x^3}$ quoniam quoniam factoris furdi latus R feu q3 --qqx-qxx-x3 divifores habet q-+x, q+x, q--x qui duarum funt magnitudinum, rejicio diviforem unum magnitudinis utriufq; & per diviforem q+x qui relinquitur multiplico factorem rationalem qq--xx. Et quoniam factum q3-+qqx-qxx-x3 æquale eft lateri R,pono m=1. & inde, cum  $\pi$  fit  $\frac{1}{3}$ , fit  $\lambda$ -I =  $-\frac{4}{3}$ . Ordinatam igitur reduco ad denominatorem R- $\frac{4}{3}$ & fit -Z° × 3q<sup>6</sup>+ 2q<sup>5</sup>x+8q<sup>4</sup>xx+8q<sup>3</sup>x<sup>3</sup>--7qqx'-6qx<sup>5</sup> ×q<sup>3</sup>--qqx-qxx-x<sup>3</sup>+ $\frac{4}{3}$ . Unde eft a = 3q<sup>6</sup>. b = 2q<sup>5</sup> & c. e=q3. f=qq & c.  $\theta$ -I = 0.  $\theta$ =I = ".  $\lambda$  =  $-\frac{1}{3}$ . r= I.  $s = \frac{2}{3}$ .  $t = \frac{1}{3}$ . v = 0. Et his in ferie fcriptis prodit area  $\frac{3qqx-3x^3}{\sqrt{cub.a_3+ax-axx-x^3}}$ , terminis omnibus in ferie tota poft tertium evanefcentibus.

#### PROP. VI. THEOR. IV.

Si Curvæ abſcifſa AB ſit z, & ſcribantur R pro e-[-fz<sup>n</sup> -[-gz<sup>2n</sup> -[-hz<sup>3n</sup>-]-&c. & S pro k -[-lz<sup>n</sup> +mz<sup>2n</sup> -[-nz<sup>3n</sup> &c. ſit autem ordinatim applicata  $z^{\theta-1}R^{\lambda-1}S^{\mu-1}$ in a-[-bz<sup>n</sup> -[-cz<sup>2n</sup> -[-dz<sup>3n</sup> &c. & ſi terminorum, e, f, g, h, &c. & k, l, m, n. &c. rectangula ſint.

ek	fk	g k	hk &c.	
el	fl	gl	hl &c.	
em	fm	gm	hm &c.	
en	fn	gn	hn &c.	

Et

# [183]

Et fi rectangulorum illorum coefficientes numerales fint respective

$$\vec{r} = \vec{r} \cdot \vec{r} - \vec{r} \cdot \vec{r} = \vec{s} \cdot \vec{s} - \vec{r} \cdot \vec{r} = \vec{r} \cdot \vec{t} + \vec{r} \cdot \vec{r} = \vec{v} \cdot \vec{k} \cdot \vec{c} \cdot \vec{r} + \vec{r} \cdot \vec{r} = \vec{v} \cdot \vec{k} \cdot \vec{r} + \vec{r} \cdot \vec{r} = \vec{v} \cdot \vec{k} \cdot \vec{r} + \vec{r} \cdot \vec{r} \cdot \vec{r} + \vec{r} \cdot \vec{r} \cdot \vec{r} + \vec{r} \cdot \vec{r} \cdot \vec{r} \cdot \vec{r} + \vec{r} \cdot \vec{r} \cdot \vec{r} \cdot \vec{r} + \vec{r} \cdot $

area Curvæ erit hæc



Ubi A denotat termini primi coefficientem datam <sup>1</sup>/<sub>a</sub> cum figno fuo – vel – , B coefficientem datam fecundi, C coefficientem datam tertii, & fic deinceps. Terminorum vero, a, b, c, &c. k, l, m, &c. unus vel plures deeffe poffunt. Demonstratur Propositio ad modum præcedentis, & quæ ibi notantur hic obtinent. Pergit autem feries talium Propositionum in infinitum, & Progressio feriei manifesta est.

#### PROP.

# [184]

## PROP. VII. THEOR. V.

Si pro  $e + fz^n - |-gz^{2n} + \&c.$  fcribatur R ut fupra, & in Curvæ alicujus Ordinata  $z_{\theta} \pm \pi \sigma R^{\lambda} \pm \pi$  maneant quantitates datæ, ", », e, f, g, &c. & pro o ac 7 scri-bantur fuccessive numeri quicunq; integri : & fi detur area unius ex Curvis quæ per Ordinatas innumeras fic prodeuntes defignantur fi Ordinatæ funt duorum nominum in vinculo radicis, vel fi dentur areæ duarum ex Curvis fi Ordinatæ funt trium nominum in vinculo radicis, vel arex trium ex Curvis fi Ordinatæ sunt quatuor nominum in vinculo radicis, & fic deinceps in infinitum : dico quod dabuntur areæ curvarum omnium. Pro nominibus hic habeo terminos omnes in vinculo radicis tam deficientes quam plenos quorum indices dignitatum funt in progressione arithmetica. Sic ordinata  $\sqrt{a^4 - ax^3 + x^4}$  ob terminos duos inter  $a^4 \& -ax^3$ deficientes pro quinquinomio haberi debet. At  $\sqrt{a^4} - x_4$  binomium eft &  $\sqrt{a^4} - x_4 - x_8^8$  trinonium, cum progressio jam per majores differentias procedat. Propositio vero sic demonstratur.

#### CAS. I.

Sunto Curvarum duarum Ordinatæ pz<sup>*θ*-1</sup> R<sup>*λ*-1</sup> & qz<sup>*θ*+1</sub>, <sup>*n*-1</sup>, <sup>*k*</sup> areæ pA & qB, exiftente R quantitate trium nominum e-[-fz<sup>*n*</sup>-[-gz<sup>2</sup>]. Et cum per Prop.</sup>

# [ 185 ]

**P**rop. III. fit  $z^{\theta} \mathbb{R}^{\lambda}$  area curvæ cujus Ordinata eft  $\theta \in \frac{+\theta}{+\lambda_{n}} fz^{n} \frac{+\theta}{+2\lambda_{n}} gz^{2n}$  in  $z^{\theta-1} \mathbb{R}^{\lambda-1}$ , fubduc Ordinatas & areas priores de area & Ordinata posteriori, & manebit  $\frac{\theta e}{-p} \frac{+\theta}{+\lambda_{n}} fz^{n} \frac{+\theta}{+2\lambda_{n}} gz^{2n}$  in  $z^{\lambda-1} \mathbb{R}^{\lambda-1}$  Ordinata nova Curvæ,  $--- q z^{n}$ 

 $z^{\theta}R^{*}-pA-qB$  ejuídem area. Pone e=p &  $\theta f = q \& Ordinata evadet \int_{\frac{1}{2}\lambda_{H}}^{\theta} gz^{2n} in z^{\theta-1} R^{\lambda-1}, \& area z^{\theta} R^{\lambda} = \theta e A = \theta f B = \lambda_{H} f B.$  Divide utramq; per eg-]-<sup>2</sup><sup>An</sup>g, & aream prodeuntem dic C, & affumpta utcunq; r, erit r C area Curvæ cujus Ordinata eft  $rz^{\theta+2\eta-1}R^{\lambda-1}$ . Et qua ratione ex areis pA & qB aream rC Ordinatæ rz<sup>0-|-2n-1</sup> R<sup>A-1</sup> congruentem invenimus, licebit ex areis qB & rC aream quartam puta sD, ordinatæ sz<sup>0+3\*-1</sup>R<sup>x-1</sup> congruentem invenire, & fic deinceps in infinitum. Et par est ratio progreffionis ab areis B & A in partem contrariam pergentis. Si terminorum 0, 0-1-10, & 0-1-21 aliquis deficit & feriem abrumpit, affumatur area pA in principio progressionis unius & area qB in principio alterius, & ex his duabus areis dabuntur areæ omnes in progressione utraque. Et contra, ex aliis duabus areis affumptis fit regreffus per analyfin ad areas A & B, adeo ut ex duabus datis cæteræ omnes dentur. Q. E. O. Hic eft cafus Curvarum ubi ipfius z index 1 augetur vel diminuitur perpetua additione vel subductione quantitatis ». Casus alter est Curvarum ubi index > augetur vel diminuitur unitatibus.

CAS.

# [186]

## CAS. II.

Ordinatæ pz<sup>θ-1</sup>R<sup>^</sup> & qz<sup>θ-1</sup>m-1</sup>R<sup>^</sup>, quibus areæ pA & qB jam refpondeant, fi in R feu e-|-fz<sup>n</sup> + gz<sup>2n</sup> ducantur ac deinde ad R viciffim applicentur, evadunt pe -|- pfz<sup>n</sup> -|- pgz<sup>2n</sup> × z<sup>θ-1</sup>R<sup>^-1</sup> & qez<sup>n</sup> -|- qfz<sup>2n</sup> -|-qgz<sup>3n</sup> × z<sup>θ-1</sup>R<sup>^-1</sup>. Et per Prop. III. eft az<sup>6</sup>R<sup>^</sup> area Curvæ cujus Ordinata eft θae +θ afz<sup>n</sup> +θ afz<sup>n</sup> + agz<sup>2n</sup> in z<sup>θ-1</sup>R<sup>^-1</sup>, & bz<sup>θ+n</sup>R<sup>^</sup> area Curvæ cujus ordinata eft -h<sup>0</sup>bez<sup>n</sup> +<sup>0</sup>bfz<sup>2n</sup> +<sup>0</sup>bgz<sup>3n</sup> in z<sup>θ-1</sup>R<sup>^-1</sup>. Et harum qua-+<sup>1</sup>/<sub>n</sub> +<sup>0</sup>/<sub>n</sub>bfz<sup>2n</sup> +<sup>0</sup>/<sub>n</sub>bgz<sup>3n</sup> in z<sup>θ-1</sup>R<sup>^-1</sup>. Et harum quatuor arearum fumma eft pA-|-qB-|- az<sup>θ</sup>R<sup>^</sup>-|-bz<sup>θ+n</sup>R<sup>^</sup> & fumma refpondentium ordinatarum

 $\stackrel{\theta_{ae}}{\stackrel{+}{\rightarrow}}_{n} afz^{n} \stackrel{+}{\stackrel{+}{\rightarrow}}_{1-2\lambda n} agz^{2n} \stackrel{+}{\stackrel{-}{\rightarrow}}_{-n} bgz^{3n} in z^{\theta-1} \mathbb{R}^{\lambda-1}.$   $\stackrel{+}{\stackrel{+}{\rightarrow}}_{n} be \stackrel{+}{\stackrel{+}{\rightarrow}}_{n} bf \stackrel{+}{\stackrel{+}{\rightarrow}}_{n} dg \stackrel{+}{\rightarrow}_{n} dg \stackrel{$ 

Si terminus primus tertius & quartus ponantur feorfim æquales nihilo, per primum fiet  $\theta_{ae-pe=0}$ feu  $-\theta_a = p$ , per quartum  $-\theta_{b-nb-2\lambda nb} = q$ , & per tertium (eliminando p & q)  $\frac{2ag}{f} = b$ . Unde fecundus fit  $\frac{\lambda naff-4\lambda nage}{f}$ , adeoq; fumma quatuor Ordinatarum eft  $\lambda naff-4\lambda nage}{f} z^{\theta+n-1}R^{\lambda-1}$ , & fumma totidem refpondentium arearum eft a  $z^{\theta}R^{\lambda} - \frac{2ag}{f}z^{\theta+n}R_{\lambda-\theta a}A - \frac{2\theta+2n+4\lambda n}{f}agB$ . DiviDividantur hæ fummæ per  $\frac{\lambda \mu^{aff} - 4\lambda \mu^{age}}{i}$ , & fi Quotum pofterius dicatur D, erit D area curvæ cujus ordinata eft Quotum prius  $z^{\theta+\eta-1}R^{\lambda-1}$ . Et eadem ratione ponendo omnes Ordinatæ terminos præter primum æquales nihilo poteft area Curvæ inveniri cujus Ordinata eft  $z^{\theta-1}R^{\lambda-1}$ . Dicatur area ifta C, & qua ratione ex areis A & B inventæ funt areæ C ac D, ex his areis C ac D inveniri poffunt aliæ duæ E & F ordinatis  $z^{\theta-1}R^{\lambda-2}$  &  $z^{\theta+\eta-1}R^{\lambda-2}$  congruentes, & fic deinceps in infinitum. Et per analyfin contrariam regredi elicet ab areis E & F ad areas C ac D, & inde ad areas A & B, aliafq; quæ in progreffione fequuntur. Igitur fi index  $\lambda$  perpetua unitatum additione vel fubductione augeatur vel minuatur, & ex areis quæ Ordinatis fic prodeuntibus refpondent duæ fimpliciffimæ habentur ; dantur aliæ omnes in infinitum. Q. E. O.

#### CAS. III.

Et per cafus hosce duos conjunctos, si tam index  $\theta$  perpetua additione vel subductione ipsus, quam index  $\lambda$  perpetua additione vel subductione unitatis, utcunq; augeatur vel minuatur, dabuntur areæ singulis prodeuntibus Ordinatis respondentes. Q. E. O.

Bbbz

CAS.

# [188]

# CAS. IV.

Et fimili augmento fi ordinata conftat ex quatuor nominibus in vinculo radicali & dantur tres arearum, vel fi conftat ex quinq; nominibus & dantur quatuor arearum, & fic deinceps : dabuntur areæ omnes quæ addendo vel fubducendo numerum n indici n vel unitatem indici  $\lambda$  generari poffunt. Et par eft ratio Curvarum ubi ordinatæ ex binomiis conflantur, & area una earum quæ non funt geometrice quadrabiles datur. Q. E. O.

#### PROP. VIII. THEOR. VI.

Si pro  $e_{+}fz^{n}_{+}gz^{2n}_{+}$ -&c. & k +  $|z^{n}_{+}-mz^{2n}_{+}$ -&e. fcribantur R & S ut fupra, & in Curvæ alicujus Ordinata  $z^{\theta+n\sigma} \mathbb{R}^{n+\frac{1}{2}\tau} S^{\mu+\frac{1}{2}\sigma}$  maneant quantitates datæ  $\theta$ , ",  $\lambda$ ,  $\mu$ , e, f, g, k, l, m, &c. & pro  $\sigma$ ,  $\tau$ , &  $\nu$ , fcribantur fucceffive numeri quicunq; integri : & fi dentur areæ duarum ex curvis quæ per ordinatas fic prodeuntes defignantur fi quantitates R & S funt binomia, vel fi dentur areæ trium ex curvis fi R & S conjunctim ex quinq; nominibus conftant, vel areæ quatuor ex curvis fi R & S conjunctim ex fex nominibus conftant, & fic deinceps in infinitum : dico quod dabuntur areæ curvarum omnium.

Demonstratur ad modum Propositionis superioris.

PROP.

[189]

## PROP.IX. THEOR. VII.

Æquantur Curvarum areæ inter fe quarum Ordinatæ funt reciproce ut fluxiones Abfciffarum.

Nam contenta fub Ordinatis & fluxionibus Abfciffarum erunt æqualia, & fluxiones arearum funt ut hæc contenta.

#### COROL. I.

Si affumatur relatio quævis inter Absciffas duarum Curvarum, & inde per Prop. 1. quæratur relatio fluxionum Absciffarum, & ponantur Ordinatæ reciproce proportionales fluxionibus, inveniri poffunt innumeræ Curvæ quarum areæ fibi mutuo æquales erunt.

## COROL. II.

Sic enim Curva omnis cujus hæc eft Ordinata  $z^{\theta-1}$  in  $e^{-fz^n} + gz^{2n} + \&c.|^{\lambda}$  affumendo quantitatem quamvis pro, & ponendo  $\frac{n}{p} = s \& z^s = x$ , migrat in aliam fibi æqualem cujus ordinata eft  $\frac{n}{n}x\frac{p\theta-n}{n}$  in  $e^{-fx^{\nu}} + gx^{2\nu} + kc.|^{\lambda}$ .

CO-

# [190]

## COROL. III.

Et Curva omnis cujus Ordinata eft  $z^{\theta-1}$  in  $a + bz^{n} + cz^{2n} + \&c. \times e + fz^{n} + gz^{2n} \&c.[^{h}, affumen-do quantitatem quamvis pro " & ponendo "= s & z^{s} = x, migrat in aliam fibi æqualem cujus ordinata$  $eft <math>\frac{v}{n}x^{\nu\theta-n}$  in  $a + bx^{\nu} + cx^{2\nu} + \&c. \times e + fx^{\nu} + gx^{2\nu} + \&c.[^{h}.$ 

#### COROL. IV.

Et Curva omnis cujus Ordinata eft  $z^{\theta-1}$  in  $a \rightarrow bz^{n} + cz^{2n} + \&c. \times e + fz^{n} + gz^{2n} + \&c.|^{h}$   $\times k \rightarrow 1z^{n} - mz^{2n} + \&c.|^{\mu}$ , affumendo quantitatem quamvis pro v ponendo  $\frac{n}{v} = s \& z^{s} = x$ , migrat in aliam fibi æqualem cujus ordinata eft  $\frac{v}{v} \times \frac{v\theta-u}{u}$  in  $a \rightarrow bx^{v}$  $(-cx^{2v} - \&c.)^{k} + c.|^{h} \otimes c.|^{h} \times k + 1x^{v} - mx^{2v} + \&c.|^{h}$ 

## COROL. V.

Et Curva omnis cujus Ordinata eft  $z^{\theta-1}$  in  $e -f \overline{z^n} - g\overline{z^{2n}} + \&c.|^{\wedge}$  ponendo  $\frac{1}{z} = x$  migrat in aliam fibi æqualem cujus ordinata eft  $\frac{1}{x^{\theta+1}} \times \overline{e+fx^{*n}}$   $\overline{+gx^{-2n}} + \&c.|^{\wedge}$  id eft  $\overline{x^{\theta+1+m\lambda}} \times \overline{f+ex^{*n}}^{\wedge}$  fi duo funt nomina in vinculo radicis vel  $\frac{1}{x^{\theta+1+m\lambda}} \times \overline{g+fx^{*n}+ex^{2n}}^{\wedge}$ fi tria funt nomina ; & fic deinceps. CO-

# [ 191 ]

#### COROL. VI.

Et Curva omnis cujus Ordinata eft  $z^{\theta-1}$  in  $e - fz^{n} + gz^{2n} + \&c. \land k + Iz^{n} + mz^{2n} + \&c. \land^{\mu}$ ponendo  $\frac{1}{z} = x$  migrat in aliam fibi æqualem cujus ordinata eft  $\frac{1}{x^{\theta+1}} \times e + fx^{-n} + gx^{-2n} + \&c. \land^{n}$   $\times \overline{k+1x^{-n}} + mx^{-2n} + \&c. \land^{\mu}$  id eft  $\frac{1}{x^{\theta+1+\frac{1}{2}m\lambda+\frac{1}{2}\mu}} \times \overline{f+ex^{n}}$   $\times \overline{l+kx^{n}}$  fi bina funt nomina in vinculis radicum, vel  $\overline{x^{\theta+1+\frac{1}{2}m\lambda+\frac{1}{2}\mu}} \times \overline{g+fx^{n}} + ex^{2n} \land^{\lambda} \times \overline{1+kx^{n}}$  fi tria funt nomina in vinculo radicis prioris ac duo in vinculo pofterioris : & fic in aliis. Et nota quod areæ duæ æquales in noviffimis hifce duobus Corollariis jacent ad contrarias partes ordinatarum. Si area in alterutra curva adjacet abfciffæ, area huic æqualis in altera curva adjacet abfciffæ productæ.

#### COROL. VII.

Si relatio inter Curvæ alicujus Ordinatam y & Absciffam z definiatur per æquationem quamvis fectam hujus formæ, y<sup>a</sup> in e + fy<sup>n</sup>z<sup>3</sup> + gy<sup>2n</sup>z<sup>20</sup> + hy<sup>3n</sup>z<sup>3<sup>3</sup></sup> + &c. = z<sup>β</sup> in k + ly<sup>n</sup>z<sup>3</sup> + my<sup>2n</sup>z<sup>20</sup> + &c. hæc figura affumendo  $s = \frac{n-3}{n}$ ,  $x = \frac{1}{s}Z^{s} & \lambda = \frac{n-3}{a\beta+\beta n}$ , migrat in aliam fibi æqualem cujus Absciffa x, ex data Ordinata

# [192]

Ordinata v, determinatur per æquationem non affectam  $\frac{1}{s}v^{\alpha\lambda} \times e^{-\frac{1}{2}}fv^{n} + gv^{2n} + hv^{3n} + \&c.|^{\lambda} \times k + 1v^{m}$  $+ mv^{2n} + \&c.|^{-\lambda} = x.$ 

#### COROL. VIII.

Si relatio inter Curvæ alicujus Ordinatam y & Abfciffam z definitur per æquationem quamvis affectam hujus formæ, y<sup>a</sup> in  $e - fy^n z^0 + gy^{2n} z^{2\theta} + \&c.$  $= z^{\beta}$  in  $k - 1y^n z^0 - my^{2n} z^{2\theta} + \&c. - z^{\gamma}$  in  $p - qy^n z^{\theta}$  $- ry^{2n} z^{2\theta} + \&c.$  hæc figura affumendo  $s = \frac{n-\theta}{n}, x = \frac{1}{s} z^s,$  $\mu = \frac{\alpha \delta + \beta n}{n-\theta} \& v = \frac{\alpha \delta + \gamma n}{n-\theta},$  migrat in aliam fibi æqualem cujus Abfciffa x ex data Ordinata v determinatur per æquationem minus affectam v<sup>a</sup> in  $e + fv^n + gv^{2n}$  $- \&c. = s^{\mu} x^{\mu}$  in  $k + 1v^n + mv^{2n} - \&c. + s^{\nu} x^{\nu}$  in  $p + qv^n + rv^{2n} - \&c.$ 

#### COROL. IX.

Curva omnis cujus Ordinata eft  $\pi z^{\theta \cdot 1}$  in  $e \ddagger_{n}^{\nu} f z^{n} \ddagger_{2n}^{\nu} g z^{2n} + \&c. \times e + f z^{n} + g z^{2n} \&c.|^{n-1} \times [a + b]e z^{\nu} + f z^{\nu+n} + g z^{\nu+n} + \&c.|^{\tau}]^{*}$ , fi fit  $\theta = n \& a$ affumantur  $x = e z^{\nu} + f z^{\nu+n} + g z^{\nu+2n} + \&c.|^{\pi}$ ,  $\sigma = \frac{\tau}{\pi}$   $\& & = \underbrace{h = \tau}{\pi}$ , migrat in aliam fibi æqualem cujus ordinata eft  $x^{\cdot \theta} \times a + b x^{\sigma}$ . Et nota quod ordinata prior in in hoc Corollario evadit fimplicior ponendo  $\lambda = 1$ , vel ponendo  $\tau = 1$  & efficiendo ut radix dignitatis extrahi poffit cujus index eft  $\omega$ , vel etiam ponendo  $\omega = -1 \& \lambda = 1 = \tau = \sigma = \pi$ , ut alios cafus præteream.

## COROL. X.

Pro  $ez^{\nu} + fz^{\nu+n} + gz^{\nu+2n} + \&c. \nu ez^{\nu-1} + fz^{\nu+n-1}$   $+ 2n gz^{\nu+2n-1} + \&c. k + 1z^n + mz^{2n} + \&c. \& nlz^{n-1}$   $+ 2nmz^{2n-1} + \&c.$  for ibantur R, r, S & s refpective, & Curva omnis cujus ordinata eft  $\pi Sr + eRs$  in  $R^{n-1} S^{\mu-1}$   $\times \overline{aS^{\nu} + bR^{\tau}}$ , fi fit  $\mu = \frac{\nu}{n} = \frac{\varrho}{\pi}$ ,  $\frac{\tau}{\pi} = \sigma$ ,  $\frac{n-\pi}{\pi} = s$ ,  $\& R^{\pi}S^{\varphi} = x$ , migrat in aliam fibi æqualem cujus ordinata eft  $x^{\varphi} \times \overline{a-k} = bx^{\sigma}/2^{\omega}$ . Et nota quod Ordinata prior evadit fimplicior, ponendo unitates pro  $\tau$ ,  $\nu$ ,  $\& \wedge vel \mu$ , & faciendo ut radix dignitatis extrahi poffit cujus index eft  $\omega$ , vel ponendo  $\omega = -1$  vel  $\mu = 0$ .

#### PROP. X. PROB. III.

Invenire figuras fimpliciffimas cum quibus Curva quævis geometrice compari poteft, cujus ordinatim applicata y per æquationem non affectam ex data abfciffa z determinatur.

# [194]

Sit Ordinata  $az^{\theta -1}$ , & area erit  $\frac{1}{\theta}az^{\theta}$ , ut ex Prop.V. ponendo b = o = c = d = f = g = h & e = 1, facile colligitur.

#### CAS. II.

Sit Ordinata  $az^{\theta-1} \times e^{-j-fz^{n}} + gz^{2n}|^{\lambda-1} + \&c. \& fi$ curva cum figuris rectilineis geometrice comparari poteft, quadrabitur per Prop. V. ponendo b = o = c=d. Sin minus convertetur in aliam curvam fibi æqualem cujus Ordinata eft  $\frac{a}{n}x^{\frac{\theta-n}{n}} \times e^{-j-fx-j-gx^{2}\&c.} \stackrel{\lambda-1}{}$ per Corol. 2. Prop. IX. Deinde fi de dignitatum indicibus  $\frac{\theta-n}{n} \& \lambda - 1$  per Prop. VII. rejiciantur unitates donec dignitates illæ fiant quam minimæ, devenietur ad figuras fimpliciffimas quæ hac ratione colligi poffunt. Dein harum unaquæq; per Corol.5. Prop. IX. dat aliam quæ nonnunquam fimplicior eft. Et ex his per Prop. III. & Corol. 9 & 10, Prop. IX. inter fe collatis, figuræ adhuc fimpliciores quandoq; prodeunt. Deniq; ex figuris fimpliciffimis aflumptis facto regreffu computabitur area quæfita.

CAS.

# [195]

## CAS. III.

Sit Ordinata  $z^{\theta-1} \times \overline{a + bz^n + cz^{2n} + \&c.}$   $\times \overline{e + fz^n + gz^{2n} + \&c.}^{h-1}$ , & hæc figura fi quadrari poteft, quadrabitur per Prop. V. Sin minus, diftinguenda eft ordinata in partes  $z^{\theta-1} \times a \times \overline{e + fz^n}$  $\overline{+ gz^{2n} + \&c.}^{h-1}, z^{\theta-1} \times bz^n \times \overline{e + fz^n + gz^{2n} + \&c.}^{h-1},$ &c. & per Caf. 2. inveniendæ funt figuræ fimpliciffimæ cum quibus figuræ partibus illis refpondentes comparari poffunt. Nam areæ figurarum partibus illis refpondentium fub fignis fuis + & conjunctæ component aream totam quæfitam.

## CAS. IV.

Sit Ordinata  $z^{\theta \cdot 1} \times \overline{a + b z^n + c z^{2n} + \&c.} \times \overline{e + f z^n + g z^{2n} + \&c.} \times \overline{k + 1 z^n + m z^{2n} + \&c.} \times \overline{k + 1 z^n + m z^{2n} + \&c.} = \frac{u \cdot 1}{2}$ & fi Curva quadrari poreft, quadrabitur per Prop. VI. Sin minus, convertetur in fimpliciorem per Corol.4. Prop. IX. ac deinde comparabitur cum figuris fimpliciffimis per Prop. VIII. & Corol. 6, 9 & 10. Prop. IX. ut fit in Cafu 2 & 3.

## CAS. V.

Si Ordinata ex variis partibus conftat, partes fingulæ pro ordinatis curvarum totidem habendæ funt,& curvæ illæ quotquot quadrari poffunt,figilla-Ccc 2 tim tim quadrandæ funt, earumq; ordinatæ de ordinata tota demendæ. Dein Curva quam ordinatæ pars refidua defignat feorfim ( ut in Cafu 2, 3 & 4,) cum figuris fimpliciffimis comparanda est cum quibus comparari potest. Et fumma arearum omnium pro area Curvæ propositæ habenda est.

# COROL. I.

Hinc etiam Curva omnis cujus Ordinata eft radix quadratica affecta æquationis fuæ, cum figuris fimpliciffimis feu rectilineis feu curvilineis compari poteft. Nam radix illa ex duabus partibus femper conftat quæ feorfim fpectatæ non funt æquanum radices affectæ. Proponatur æquatio aayy  $-\neg zzyy = 2a^3y - \frac{2}{2^3}y - \frac{2}{2^4}$ , & extracta radix erit  $y = \frac{a^3 - \frac{1}{2^3} - \frac{2}{2^3}y - \frac{2}{2^4}}{a^3 - \frac{2}{2^2}}$  & pars irrationalis  $\frac{a\sqrt{a^4 - 2az^3 - z^4}}{a^3 - \frac{2}{2^2}}$  funt ordinatæ curvarum quæ per hanc Propofitionem vel quadrari poffunt vel cum figuris fimpliciffimis comparari cum quibus collationem geometricam admittunt.

# COROL. II.

Et curva omnis cujus Ordinata per æquationem quamvis affectam definitur quæ per Corol. 7. Prop. IX. in æquationem non affectam migrat, vel quadratur dratur per hanc Propositionem fi quadrari potest vel comparatur cum figuris simpliciss cum quibus compari potest. Et hac ratione Curva omnis quadratur cujus æquatio est trium terminorum. Nam æquatio illa fi affecta sit transmutatur in non affectam per Corol.7. Prop.IX. ac deinde per Corol. 2 & 5. Prop. IX. in simplicissimam migrando, dat vel quadraturam figuræ si quadrari potest, vel curvam simplicissimam quacum comparatur.

#### COROL. III.

Et Curva omnis cujus Ordinata per æquationem quamvis affectam definitur quæ per Corol. 8. Prop. IX. in æquationem quadraticam affectam migrat ; vel quadratur per hanc Propositionem & hujus Corol. 1. fi quadrari potest, vel comparatur cum figuris fimpliciss cum quibus collationem geometricam-admittit.

### SCHOLIUM.

Ubi quadrandæ funt figuræ; ad Regulas hafce generales femper recurrere nimis moleftum effet : præftat Figuras quæ fimpliciores funt & magis ufui effe poffunt femel quadrare & quadraturas in Tabulam referre, deinde Tabulam confulere quoties ejufmodi Curvam aliquam quadrare oportet. Hujus autem generis funt Tabulæ duæ fequentes, in quibus z denotat Abfciffam, y Ordinatam rectangulam

# [ 198 ]

gulam & t Aream Curvæ quadrandæ, & d, e, f, g, g. h, " funt quantitates datæ cum fignis fuis-|-&--.

# TABULA

Curvarum simpliciorum quæ quadrari possunt.

Curvarum formæ. Curvarum areæ.

Forma prima.

$$dz^{n-1} = y, \qquad \qquad \overset{d}{n} z^n = t.$$

Forma fecunda.

 $\frac{\mathrm{d} z^{n-1}}{\frac{\mathrm{ce} - \left[-2\mathrm{ef} z_n - \left[-\mathrm{ff} z_2\right]\right]}{\mathrm{ff} z_2}} = y.$ 

$$\frac{\mathrm{d} z_n}{n^{\mathrm{ee-}}+n^{\mathrm{efz}_n}} = t, \text{ vel } \frac{-\mathrm{d}}{n^{\mathrm{ef-}}+n^{\mathrm{ffz}_n}} = t.$$

Forma tertia.

1. 
$$dz_{-1}^{n}\sqrt{e_{-}+fz^{n}} = y$$
.  $\frac{2d}{3n}R^{3} = t$ , exiftente  $R = \sqrt{e_{-}+fz^{n}}$   
2.  $dz_{-1}^{2n}\sqrt{e_{-}+fz^{n}} = y$ .  $\frac{-4e_{-}+6fz_{n}}{15n}dR^{3} = t$ .  
3.  $dz_{-1}^{3n}\sqrt{e_{-}+fz^{n}} = y$ .  $\frac{16ee_{-}-24efz_{n}+30ffz_{2n}}{105nf_{3}}dR^{3} = t$ .  
4.  $dz_{-1}^{4n}\sqrt{e_{-}+fz^{n}} = y$ .  $\frac{-96e_{3}-1-144eefz_{n}-180effz_{2n}-1210f_{3}z_{3}n}{945n^{f+}}dR^{3} = t$ .

Forma quarta.

I. 
$$\frac{dZ^{n-1}}{\sqrt{e_{-}-fz_{n}}} = y.$$
$$\frac{2d}{nf}R = t.$$
  
2. 
$$\frac{dZ^{2n-1}}{\sqrt{e_{-}-fz_{n}}} = y.$$
$$\frac{-4e_{-}-2fz_{n}}{3\mu ff}dR = t.$$

dz3MI



# TABULA

# Curvarum simpliciorum quæ cum Ellipsi & Hyperbola compari possunt.

Sit jam aGD vel PGD vel GDS Sectio Conica cujus area ad Quadraturam Curvæ pro-Fig. 5,6,7,8. pofitæ requiritur, fitq; ejus centrum A, Axis Ka, Vertex a, Semiaxis conjugatus AP, datum Absciffæ principium A vel a vel a, Abscissa AB vel a B vel  $_{a}B=x$ , Ordinata rectangula BD=v, & Area A B D P vel a B DG vel  $\alpha$  B DG = s, existente  $\alpha$ G Ordinata ad punctum «. Jungantur KD, AD, a D. Ducatur Tangens DT occurrens Absciffæ AB in T, & compleatur parallelogrammum ABDO. Et fiquando ad quadraturam Curvæ propofitæ requiruntur arez duarum Sectionem Conicarum, dicatur posterioris Abscissa &, Ordinata T, & Area o. Sit autem - differentia duarum quantitatum ubi incertum est utrum posterior de priori an prior de pofteriori fubduci debeat.

Curva-

						2	
						<sup>4 ces</sup> = t.	Form
		ئە <sup>.</sup>		DGa.	=t.	2	
aGDB	t.	$\frac{de}{dH}Z^{n} \frac{ee}{nH}S =$		$s = t = \frac{4}{n} A$	$2\frac{n}{2} + \frac{4es-2}{nf}$	$\frac{3}{2}$ $\frac{2dee}{n}$ <b>Z</b>	
$\frac{1}{n}S = t =$	d Zu c	d Z <sup>2</sup> h Z <sup>2</sup> h		$= V_{\circ} \frac{2XV_{\circ} \cdot 4}{n}$	-V. <sup>2de</sup>	$= V_{e} \frac{2de}{3nf} 2$	
$\frac{d}{c-ft} = V.$	$\frac{d}{dt} = v$ .	$\frac{\mathrm{d}}{\mathrm{e}_{-}-\mathrm{fx}} = \mathbf{V}_{*}$			$\frac{d}{f} - \frac{e}{f}XX$	/d _f _f XX	
Z <sup>ti</sup> ≐X.	Z'=X.	Z <sup>4</sup> = X.		$\frac{d}{dt_{2n}} = \mathbf{x}. \ \mathbf{v}$	$\frac{d}{dt_n} = x$ .	$\frac{d}{dt} = X.$	
-= y.	= y.	·= y.	ecunda.	= y. V_ <u>e_</u>	$= y$ . $\sqrt{c_{e}}$	·= y. <	
I. e-f-fzu	2. e-j-fzn	dz <sup>31-1</sup> 3. e- -fz <sup>n</sup>	Forma fo	I. $\frac{dz^{\frac{1}{2}h-1}}{e+fz^{n}}$	2. e-j-fZn	3. e+fzn	
ig. 5.				Tig. 6, 7.			
	Fig. 5. 1. $\frac{dz^{n-1}}{e-fz_n} = y$ . $z^n = x$ . $\frac{d}{e-fx} = v$ . $\frac{1}{n}s = t = \frac{aGDB}{n}$	Fig. 5. I. $\frac{dz^{n-1}}{e-fz_n} = y$ . $z^n = x$ . $\frac{d}{e-fx} = v$ . $\frac{1}{n}s = t = \frac{dDB}{n}$ 2. $\frac{dz^{2n-1}}{e-fz_n} = y$ . $z^n = x$ . $\frac{d}{e-fx} = v$ . $\frac{d}{nt}z^{n-\frac{e}{n}}s = t$ .	Fig. 5. 1. $\frac{dz^{\mu t}}{c - fz_{\mu}} = y$ . $z^{\mu} = x$ . $\frac{d}{c + fx} = v$ . $\frac{1}{n}s = t = \frac{dDB}{n}$ 2. $\frac{dz^{2\mu t}}{c - fz_{\mu}} = y$ . $z^{\mu} = x$ . $\frac{d}{c + fx} = v$ . $\frac{d}{n}z^{\mu} - \frac{c}{n}s = t$ . 3. $\frac{dz^{3\mu t}}{c - fz_{\mu}} = y$ . $z^{\mu} = x$ . $\frac{d}{c - fx} = v$ . $\frac{d}{2n}z^{2n} - \frac{dc}{n}z^{2n} + \frac{cc}{n}s = t$ .	Fig. 5. 1. $\frac{dz^{n-1}}{e-1} = y$ . $z^n = x$ . $\frac{d}{e-1} = v$ . $\frac{1}{n} s = t = \frac{d\Omega B}{n}$ 2. $\frac{dz^{2n-1}}{e-1} = y$ . $z^n = x$ . $\frac{d}{e-1} = v$ . $\frac{d}{n!} z^{n-\frac{e}{n!}} s = t$ . 3. $\frac{dz^{3n-1}}{e-1} = y$ . $z^n = x$ . $\frac{d}{e-1} = v$ . $\frac{d}{n!} z^{n-\frac{e}{n!}} s = t$ . Forma fecunda.	Fig. 5. 1. $\frac{dz^{n-1}}{e^{-1}fz_n} = y$ . $z^n = x$ . $\frac{d}{e^{-1}fx} = v$ . $\frac{1}{n}s = t = \frac{dGDB}{n}$ 2. $\frac{dz^{2n-1}}{e^{-1}fz_n} = y$ . $z^n = x$ . $\frac{d}{e^{-1}fx} = v$ . $\frac{d}{n!}z^{n-\frac{n}{n!}}s = t$ . 3. $\frac{dz^{2n-1}}{e^{-1}fz^n} = y$ . $z^n = x$ . $\frac{d}{e^{-1}fx} = v$ . $\frac{d}{2n}z^{n-\frac{n}{n!}}s^{n-\frac{n}{n!}}s = t$ . Forma fecunda. Fig. 6.7. 1. $\frac{dz^{2n-1}}{e^{-1}fz^n} = y$ . $\sqrt{\frac{d}{n}} = x$ . $\sqrt{\frac{d}{n}} - \frac{e^{-\frac{n}{n!}}s^{n-\frac{n}{n!}s^{n-\frac{n}{n!}}s^{n-\frac{n}{n!}s^{n-\frac{n}{n!}}s^{n-\frac{n}{n!}s^{n-\frac{n}{n!}}s^{n-\frac{n}{n!}s^{n-$	Fig. 5. 1. $\frac{dz^{n+1}}{e-tz_n} = y$ . $z^n = x$ . $\frac{d}{e^{t+x}} = v$ . $\frac{1}{a}s = t = \frac{dB}{a}$ 2. $\frac{dz^{2n+1}}{e-tz_n} = y$ . $z^n = x$ . $\frac{d}{e^{t+x}} = v$ . $\frac{d}{at} z^{n-\frac{n}{n}} s^s = t$ . 3. $\frac{dz^{2n+1}}{e-tz_n} = y$ . $z^n = x$ . $\frac{d}{e^{t+x}} = v$ . $\frac{d}{at} z^{n-\frac{n}{n}} s^s = t$ . Forma founda. Fig. 6,7. 1. $\frac{dz^{2n+1}}{e-tz_n} = y$ . $\sqrt{\frac{d}{t-1}} = x$ . $\sqrt{\frac{d}{t} - \frac{e}{t}xx} = v$ . $\frac{z^{n+\frac{n}{n}}}{a} s^{-\frac{n}{n}} t^s = t^{-\frac{n}{n}} A DGa$ . 2. $\frac{dz^{2n+1}}{e-tz_n} = y$ . $\sqrt{\frac{d}{e^{t-1}}} = x$ . $\sqrt{\frac{d}{t-1}} - \frac{e}{t}xx = v$ . $\frac{z^{n+\frac{n}{n}}}{a} = t = \frac{d}{a} A DGa$ .	Fig. 5. 1. $\frac{d2^{n+1}}{e^{-1}f2^{n}} = y$ . $2^{n} = x$ . $\frac{d}{e^{1+x}} = v$ . $\frac{1}{n} S = t = \frac{e^{GDB}}{n}$ 2. $\frac{d2^{2n+1}}{e^{-1}f2^{n}} = y$ . $2^{n} = x$ . $\frac{d}{e^{-1}fx} = v$ . $\frac{d}{n}Z^{n} - \frac{e^{C}}{n!}S = t$ . 3. $\frac{d2^{2n+1}}{e^{-1}f2^{n}} = y$ . $2^{n} = x$ . $\frac{d}{e^{-1}fx} = v$ . $\frac{d}{n}Z^{n} - \frac{e^{C}}{n!}S^{n} - \frac{e^{C}}{n!}S^{n} + \frac{e^{C}}{n!}S^{n}$

[200]

4de ... rnr ... ! ... ADDRS TDR 4dein V2 Forma tertia. d . / -1-fr.

Fig.6,7,8. 1. 
$$\frac{1}{2}$$
 Vertual = y.  $\frac{1}{2n}$  = XX. Vertex = V.  $\frac{1}{nH}$   $\frac{1}{nEx}$  = S = t =  $\frac{\pi}{nF}$  in aGD1, Vertin APDE TDB.  
Vel ffc,  $\frac{1}{2n}$  = X.  $\sqrt{ix+exx} = V. \frac{8de}{nH}$  in  $S-\frac{1}{2}$  Vu  $\frac{fv}{4e}$   $+ \frac{fv}{4eex}$  = t =  $\frac{8de}{nH}$  in aGDA  $+ \frac{fv}{4eex}$ .  
2.  $\frac{d}{2r_{11}^{\mu}}$   $\sqrt{ertin}$  = y.  $\frac{1}{2n}$  = XX.  $\sqrt{irtexx} = V. \frac{2d}{n}$  S = t =  $\frac{2d}{n}$  APDB, feu  $\frac{2d}{n}$  aGDB.  
Yel fic,  $\frac{1}{2n}$  = X.  $\sqrt{irtexx} = V. \frac{2d}{nF}$  in  $S-\frac{1}{2}$  Vu  $-\frac{fv}{2e}$  = t =  $\frac{4de}{nT}$  × aGDB.  
3.  $\frac{d}{2r_{11}^{\mu}}$   $\sqrt{ertin}$  = y.  $\frac{1}{2n}$  = X.  $\sqrt{irtexx} = V. \frac{d}{nF}$  in  $S-\frac{1}{2}$  ×  $-a$ GDB vel BDFK.  
4.  $\frac{d}{2r_{11}^{\mu}}$   $\sqrt{ertin}$  = y.  $\frac{1}{2n}$  = X.  $\sqrt{irtexx} = V. \frac{d}{nS}$  = t =  $\frac{d}{n}$  ×  $-a$ GDB vel BDFK.  
Forma quarta.

2.  $\frac{d}{2!} = y \cdot \frac{1}{2^n} = y \cdot \frac{1}{2^n} = xx$ .  $\sqrt{f + exx} = y \cdot \frac{2d}{ne}$  in  $s - xy = t = \frac{2d}{ne}$  in POD, vel in AODGa. Fig. 6. I.  $\frac{d}{Z\sqrt{e+fz_n}} = y \cdot \frac{1}{z_n} = x_x$ .  $\sqrt{F+exx} = v$ .  $\frac{4d}{n^f}$  in  $\frac{1}{2}xv \div s = t = \frac{4d}{n^f}$  in PAD vel in aGDA. Vel fic,  $\frac{1}{2\eta} = x$ .  $\sqrt{fx-fexx} = v$ .  $\frac{8de}{nff}$  in  $s - \frac{1}{2}xv - \frac{fv}{4e} = t = \frac{8de}{nff}$  in aGDA. Ddd

Vel fic, 
$$\frac{1}{z_{\eta}} = x$$
,  $\sqrt{ix+exx} = v$ .  $\frac{4d}{\pi i}$  in  $\frac{1}{2}xv \div s = t = \frac{4d}{n^{2}}$  in aDGa.

3. Z +1

[201]


	[203]	
4 <u>dfgs</u> = t. S = t.	∆DBA.	Forma
ff ac GDB c ff c ff t f	₫GDB <u>⊣</u>	t = t, xv + rode
$\frac{\det f Y - 2df f}{4n eg - n}$ $t = \frac{d}{n} in$ $\frac{df}{2ng} S$	.8dg #cgff in	-2defg v-l-10deff
$4\frac{dee\xi r-1-2}{n}$ $\frac{d}{n} S = 1$ $\frac{d}{n} V = \frac{d}{24ng}$	$\frac{v}{dt} = t = \frac{1}{4}$	-2dff XV. 4ncgg mffg -8degg XX -2dffg XX 24ncg3 -
x = v. y = r. y =	14dgxv-2df 1cg-nff 1cg-1-2d 4ncg	36defg S
-fxgx -fxgx fx+- c+fx+- c+fx	$v = V \cdot \frac{8 dgs}{44}$	XX = V.
₹. V <del>g+</del> ₹. V <del>g+</del> =x. V <del>g</del>	try - fx - gxx =	+fx+8 +fx-1-8
$y \cdot \begin{cases} Z_n = \\ y \cdot Z_n = \\ -y \cdot Z^n \end{cases}$ $= y \cdot Z^n \cdot Z^n$	x.Ve-l-fy = x. Ve-	- x. Ve-
$+gz^{2n} =$	a. = y.z"= = y. z"=	$= y \cdot z^{a} =$
et fzu- "Vet fi "Vet fi "Vet fi	ma očtav dz <sup>n-1</sup> -f <sup>-f</sup> z <sup>n-1</sup> -g <sup>2</sup> n dz <sup>2</sup> n <sup>-1</sup>	$\frac{dz^{3n-1}}{fz^{n-1}-gz^{2n}}$
$\frac{1}{2} dz$	Forr P. Ve-	4 ×
Fig. 6,7	Fig. 6.	Ddd 2

Forma feptima.



204]

In Tabulis hifce, feries Curvarum cujufq; formæ utrinq; in infinitum continuari poteft. Scilicet in Tabula prima, in numeratoribus arearum formæ tertiæ & quartæ, numeri coefficientes initialium terminorum (2, -4, 16, -96, 868, &c.) generantur multiplicando numeros -2, -4, -6, -10, &c.in fe continuo, & fubfequentium terminorum coefficientes ex initialibus derivantur multiplicando ipfos gradatim, in Forma quidem tertia, per  $-\frac{3}{2}, -\frac{7}{6}, -\frac{2}{8}, -\frac{11}{10}$  &c. in quarta vero per  $-\frac{1}{2}, -\frac{3}{4}, -\frac{5}{6}, -\frac{7}{8}, -\frac{9}{10}$ , &c. Et Denominatorum coefficientes 3, 15, 105, &c. prodeunt multiplicando numeros 1, 3, 5, 7, 9, &c. in fe continuo.

In fecunda vero Tabula, feries Curvarum formæ primæ, fecundæ, quintæ, fextæ, nonæ & decimæ ope folius divifionis, & formæ reliquæ ope Propofitionis tertiæ & quartæ, utrinq; producuntur in infinitum.

Quinetiam hæ feries mutando fignum numeri variari folent. Sic enim, e. g. Curva  $\frac{d}{z}\sqrt{e+fz^*} = y_2$ evadit  $\frac{d}{z_{z}^{i_{1}}+1}\sqrt{f+ez^*}$ .

## PROP. IX. THEOR. VIII.

Sit ADIC Curva quævis Abfeiffam habens Fig. 9.4 AB=z & Ordinatam BD=y, & fit AEKC Curva alia cujus Ordinata BE æqualis eft prioris areæ ABC

A DB ad unitatem applicatæ, & A F L C Curva tertia cujus Ordinata B Fæqualis eft fecundæ areæ A E B ad unitatem applicatæ, & A G M C Curva quarta cujus Ordinata B Gæqualis eft tertiæ areæ A F B ad unitatem applicatæ, & A H N C Curva quinta cujus Ordinata B Hæqualis eft quartæ areæ A G B ad unitatem applicatæ, & fic deinceps in infinitum. Et funto A, B, C, D, E, &c. Areæ Curvarum Ordinatas habentium y, zy, z<sup>2</sup>y, z<sup>3</sup>y, z<sup>4</sup>y, & Abfciffam communem z.

Detur Abfciffa quævis AC=t, fitq; BC=t-z =x, & funto P, Q, R, S, T areæ Curvarum Ordinatas habentium x, xy, xxy, x<sup>3</sup>y, x<sup>4</sup>y & Abfciffam communem x.

Terminenter autem hæ areæ omnes ad Absciffam totam datam AC, nec non ad Ordinatam positione datam & infinite productam CI : & erit arearum sub initio positarum prima ADIC=A=P, fecunda AEKC=tA-B=Q.Tertia AFLC= $\frac{ttA-2tB+C}{2} = \frac{1}{2}R$ . Quarta AGMC =  $\frac{t_3A-3ttB+3tC-D}{6} = \frac{t}{6}S$ . Quinta AHNC =  $\frac{t_4A-4t_3B+-6ttC-4tD+E}{24} = \frac{1}{24}T$ .

C 0-

# [207]

## COROL.

Unde fi Curvæ quarum Ordinatæ funt y, zy, z'y, z'y, &c. vel y, xy, x'y, x'y, &c. quadrari poffunt, quadrabuntur etiam Curvæ ADIC, AEKC, AFLC, AGMC, &c. & habebuntur Ordinatæ BE, BF, BG, BH areis Curvarum proportionales.

# SCHOLIUM.

Quantitatum fluentium fluxiones effe primas, fecundas, tertias, quartas, aliafq; diximus fupra. Hæ fluxiones funt ut termini ferierum infinitarum convergentium. Ut fi z<sup>n</sup> fit quantitas fluens & fluendo evadat  $\overline{z} \rightarrow 0^{n}$ , deinde refolvatur in feriem convergentem  $z^{n} \rightarrow n0z^{n-1} \rightarrow \frac{n+n}{2} \text{ ooz } n^{-2} \rightarrow \frac{n^{3}-3n}{6} + \frac{2n}{6}0^{3}z^{n-3}$  $\rightarrow \inftyc$ . terminus primus hujus feriei  $z^{n}$  erit quantitas illa fluens, fecundus  $n0z^{n-1}$  erit ejus incrementum primum feu differentia prima cui nafcenti proportionalis eft ejus fluxio prima, tertius  $\frac{nn+n}{2}0z^{n-2}$ erit ejus incrementum fecundum feu differentia fecunda cui nafcenti proportionalis eft ejus fluxio fecunda, quartus  $\frac{n^{3}-3nn+2n}{6}0^{3}z^{n-3}$  erit ejus incrementum tertium feu differentia tertia cui nafcenti fluxio tertia proportionalis eft, & fic deinceps in infinitum.

' Exponi

Exponi autem poffunt hæfluxiones per Curvarum Ordinatas BD, BE, BF, BG, BH, &c. Ut fi Ordinata BE  $(=\frac{ADB}{T})$  fit quantitas fluens, erit ejus fluxio prima ut ordinata BD. Si BF  $(=\frac{AEB}{T})$ fit quantitas fluens, erit ejus fluxio prima ut Ordinata BE & fluxio fecunda ut Ordinata BD. Si BH  $(=\frac{AGB}{T})$  fit quantitas fluens, erunt ejus fluxiones, prima, fecunda, tertia & quarta, ut Ordinatæ BG, BF, BE, BD refpective.

Et hinc in æquationibus quæ quantitates tantum duas incognitas involvunt, quarum una est quantitas uniformiter fluens & altera est fluxio quælibet quantitatis alterius fluentis, inveniri potest fluens illa altera per quadraturam Curvarum. Exponatur enim fluxio ejus per Ordinatam BD, & fi hæc fit fluxio prima, quæratur area  $ADB = BE \times I$ , fi fluxio lecunda, quæratur area  $AEB = BF \times I$ , fi fluxio tertia, quæratur area  $AFB = BG \times I$ , &c. & area inventa erit exponens fluentis quæsitæ.

Sed & in æquationibus quæ fluentem & ejus fluxionem primam fine altera fluente, vel duas ejufdem fluentis fluxiones, primam & fecundam, vel fecundam & tertiam, vel tertiam & quartam, &c. fine alterutra fluente involvunt : inveniri poffunt fluentes per quadraturam Curvarum. Sit æquatio aav = av - vv, exiftente v = BE, v = BD, z = AB & z = 1, & æquatio illa complendo dimensiones fluxionum, evadet aav = avz + vvz, feu  $\frac{aav}{av + vv} = z$ . Jam fluat v uniformiter & fit [209]

fit ejus fluxio v=1 & erit  $\frac{aa}{av+vv}=z$ , & quadrando Curvam cujus Ordinata eft  $\frac{aa}{av+vv}$  & Abfeiffa v, habebitur fluens z. Adhæc fit æquatio aav=av+vv exiftente v=BF, v=BE, v=BD & z=AB & per relationem inter v & v feu BD & BE invenietur relatio inter A B & BE ut in exemplo fuperiore. Deinde per hanc relationem invenietur relatio inter AB & BF quadrando Curvam AEB.

Æquationes quæ tres incognitas quantitates involvunt aliquando reduci poffunt ad æquationes quæ duas tantum involvunt, & in his cafibus fluentes invenientur ex fluxionibus ut fupra. Sit æquatio  $a-bx^m=cxy_ny-(-dy^{2n}yy)$ . Ponatur  $y^ny=v$  & erit  $a-bx^mcxv-(-dvv)$ . Hæc æquatio quadrando Curvam cujus Abfciffa eft x & Ordinata v dat aream v, & æquatio altera  $y^ny=v$  regrediendo ad fluentes dat  $\frac{1}{n!+1}y^{n+1}=v$ . Unde habetur fluens y.

Quinetiam in æquationibus quæ tres incognitas involvunt & ad æquationes quæ duas tantum involvunt reduci non poffunt, fluentes quandoq; prodeunt per quadraturam Curvarum. Sit æquatio  $a x^m + b x^{n}|^p = r e x^{r-1} y^s + s e x^r y y^{s-1} - f y y^t$ , exiftente x = I. Et pars pofterior  $r e x^{r-1} y^s + s e x^r y y^{s-1} - f y y^t$ , regrediendo ad fluentes, fit  $e x_r y^s - \frac{f}{t+1} y^{t+1}$ , quæ proinde eft ut area Curvæ cujus Abfciffa eft x & Ordinata  $a x^m + b x^{n}|^p$ , & inde datur fluens y.

Eee

Sit

[ 210 ]

Sit æquatio  $x \times ax^m + bx^n p = \frac{dy y_{n-1}}{\sqrt{c+ty_n}}$ . Et fluens cujus fluxio eft  $x \times ax^m + bx^n p$  erit ut area Curvæ cujus Abfeiffa eft x & Ordinata eft  $ax^m + bx^n p$ . Item fluens cujus fluxio eft  $\frac{dy y_{n-1}}{\sqrt{c+ty_n}}$  erit ut area Curvæ cujus Abfeiffa eft  $y & Ordinata \frac{dy y_{n-1}}{\sqrt{c-ty_n}}$ , id eft (per Cafum I. Formæ quartæ Tab. I.) ut area  $\frac{2d}{nf}\sqrt{c-ty_n}$ . Pone ergo  $\frac{2d}{nf}\sqrt{c-ty_n}$  æqualem areæ Curvæ cujus Abfeiffa eft  $x & Ordinata ax^m + bx^n p$ & habebitur fluens y.

Et nota quod fluens omnis quæ ex fluxione primæ colligitur augeri poteft vel minui quantitate quavis non fluente. Quæ ex fluxione fecunda colligitur augeri poteft vel minui quantitate quavis cujus fluxio fecunda nulla eft. Quæ ex fluxione tertia colligitur augeri poteft vel minui quantitate quavis cujus fluxio tertia nulla eft. Et fic deinceps in infinitum.

Postquam vero fluentes ex fluxionibus collectæ funt, fi de veritate Conclusionis dubitatur, fluxiones fluentium inventarum vicissim colligendæ funt & cum fluxionibus sub initio propositis comparandæ. Nam si prodeunt æquales Conclusio recte se habet :

# 

bet : fin minus, corrigendæ funt fluentes fic, ut earum fluxiones fluxionibus fub initio propofitis æquentur. Nam & Fluens pro lubitu affumi potest & affumptio corrigi ponendo fluxionem fluentis affumptæ æqualem fluxioni propositæ, & terminos homologos inter se comparando.

FINIS.

Et his principiis via ad majora sternitur.

## ERRATA.

1.11

## BOOK I. Of Opticks.

Part 1. p.3. 1.20. Properties which, ib.p.5. 1.5. and that C, p.6. 1.9. DE, p.21. 1.23. are two Rays, p.27. 1.6. in the Margin put Fig.14 & 15. p.30.1.7. MN, 1.9. M, p. 44. 1.15. as was proposed, p.52. 1. 17. a paper Circle, p.57. 1.ult. emerging, p.60. 1.25. contain with the, p.64. 1.18. and 14th, p.65. 1.13. at the, p.66. 1.3. semicircular, p.67. 1.25. Center, 1.31.  $\frac{1}{48}$  Inches, p.68. 1.8. to 16, 1.9. or  $5\frac{1}{3}$ , p.71. 1.1. bifett, p.72. 1.13. falls, 1.20. being. Part II. p.86. 1.5. lelopipede, p.89. 1.9. made by, p.93. 1.18. to  $77\frac{1}{3}$ , 1.28,29, by the third Axiom of the first Part of this Book, the Laws, p.105. 1.5. fee reprefented, p. 144.1. 24,  $\frac{1}{9}$ ,  $\frac{1}{16}$ ,  $\frac{1}{10}$ ,  $\frac{1}{9}$ ,  $\frac{1}{16}$ ,  $\frac{1}{9}$ , p. 118, 119. for Lib.1. Lib.2. write Part 1. Part 2. p.122. 1.9. indico, p. 130. 1.19. to the Angle, p.132. 1.6. by the brightnefs, p.135. 1.14. For if in the, 1.16. first Part you, p.136. 1.26. first Part, 1.27. lights, p.137. 1.20. green, accordingly as, p. 138. 1.21. Prop.6. Part. 2. p.139. 1.5. on which, p.142.1.17. XT which have been, p.143. 1.7. purple, 1.16. feveral Lights, 1.24. of white.

### BOOK. II.

P.5. 1. Sincely the, p. 7. 1.9. y, z denote, 1.28. them divers, p. 10. 1. 24. 1000 to 1024, p.11. 1.11. oliquities, I. p.17. 1.4.  $14\frac{1}{3}$  to 9, p. 25. 1. 11.  $10\frac{4}{5}$ , p.31. 1. 12. more compounded, p.55. 1.3. fizes reflect, 1.24. and therefore their Colours arife, p.65. 1.5. corpufcles can, p.71. 1.17. given breadth, p.84. 1. 4. are to thole, p. 96. 1. 24. Observation of this Part of this Book, p.103. 1.17. mas to the thickness, p. 105. 1. 19. of this rehite Ring, p.107. 1.20. become equal to the third of those.

#### Enumeratio Linearum.

p.143. 1.20. datas signis suis, p. 144. 1.27. respiciunt, p. 146. 1.5. sunt Asymptoto, p. 154. 1.13. cx-1-d dat Ordinatam y =, 1.14. qua generatur.

### Quadratura Curvarum.

p.168.1.24.*reAa AB*, p.176.1.ult.  $+ \frac{\theta}{\lambda n}$  fZ<sup>n</sup>, p.183.1.13.*a*,*b*,*c*, Sc. *e*,*f*,*g*, Sc. *k*, *l*, *m*, Sc. p. 185. 1.4. *in* Z0-1, p.188. 1.14. Z0  $\pm$  *no*, p. 190. 1. 19. *vel*  $\frac{1}{\chi \theta - \tau} = 1 \pm 2n\chi$ . p. 192. 1. 18. *g*Z<sup>p</sup> + 2n. p.193. 1.11. *aSp*-*f*-*b*R $\tau/\omega$ .

P29 212. Quadr: Tab.I. H K 3 1 C E  $\mathbf{P}_{\mathbb{N}}$ 2 ( C.J B Ъ A D А B G D d 6 4 5 P 0 B А N S N Å B Á Ваа D Ė à K м H  $\mathcal{S}$ 7 0 9 0 P D Ŕ B· N BC B A a A 1 A







---

``

۲.

ر ر

2-10-47

۸ .

